

題名:Mechanoelectrical feedback regulates the arrhythmogenic activity of pulmonary veins.

作者:陳亦仁

Chang SL; Chen YC; Chen YJ; Wangcharoen W; Lee SH; Lin CI; Chen SA

貢獻者:臨床醫學研究所

上傳時間:2009-08-21T08:58:43Z

摘要:BACKGROUND: Atrial fibrillation is commonly associated with dilated pulmonary veins. Stretch has been shown to have mechano-electrical effects. OBJECTIVE: To investigate whether stretch can increase the arrhythmogenic activity of the pulmonary veins. METHODS: The transmembrane action potentials were recorded from rabbit pulmonary veins before and after stretch (100 and 300 mg). Gadolinium and streptomycin (stretch-activated ion channel blockers) were each perfused into the pulmonary veins under a 300-mg stretch. RESULTS: Stretch (0, 100 and 300 mg) force dependently increased the incidence of spontaneous activity (22%, 48% and 83%; $p<0.05$), mean (standard deviation (SD)) firing rates of spontaneous activity (1.7 (0.2), 2.1 (0.3) and 3 (0.2) Hz; $p<0.05$) and incidence of early post-depolarisations (9%, 26% and 61%; $p<0.05$) and delayed post-depolarisations (0%, 4% and 30%; $p<0.05$) in 23 pulmonary veins. In the seven preparations with spontaneous activity after the 300-mg stretch, gadolinium (1, 3 and 10 $\mu\text{mol/l}$) decreased the incidence of spontaneous activity by 43%, 29% and 14%, respectively ($p<0.05$), and decreased the firing rate from 2.9 (0.1) Hz to 0.8 (0.4), 0.3 (0.1) and 0.1 (0.1) Hz, respectively ($p<0.05$). Streptomycin (10 and 40 $\mu\text{mol/l}$) decreased the incidence of spontaneous activity by 71% and 29%, respectively ($p<0.05$), and decreased the firing rate from 2.9 (0.1) Hz to 1.6 (0.4) and 0.5 (0.3) Hz, respectively ($p<0.05$). CONCLUSION: Stretch is an important factor in the electrical activity of the

pulmonary vein. Stretch-induced arrhythmogenic activity of the pulmonary vein may contribute to the genesis of atrial fibrillation.