

# **Consumption of purple sweet potato leaves decreases**

## **lipid peroxidation and DNA damage in humans**

Chiao-Ming Chen, Ya-Ling Lin, C-Y Oliver Chen, Ching-Yun Hsu, Ming-Jer Shieh  
PhD, Jen-Fang Liu

**Hsu;MJ;Hsu;CY;Chen;BC;Chen;MC;Ou;George;Lin;CH**

### **Abstract**

Consumption of polyphenols is associated with reduced risk of chronic diseases, possibly via a variety of biomechanisms, including antioxidation and anti-inflammation. Purple sweet potato leaves (PSPL) commonly consumed in Asia possess polyphenols. In this study, we aim to investigate antioxidant effect of 200 g/d PSPL containing 902 mg polyphenols in a clinical trial. This randomized, crossover clinical study included 16 healthy adults (7 M, 9 F; aged 20-22 y). After a 1-wk run period, subjects were assigned randomly to receive either PSPL or low polyphenol diet (LPD) for 2 wks, followed by a 2-wk washout period before crossing over to the alternate diet. Fasting blood and 24-h urine samples were collected from each subject at day 0, 7 and 14 of each phase. Our data showed PSPL consumption enhanced urinary total phenol excretion by 24.5% at day 14 as compared to day 0, while the LPD decreased total phenol content in plasma and urine by 3.3 and 16.3%, respectively ( $p \leq 0.05$ ). Low-density lipoprotein lag time and glutathione concentration in erythrocytes at day 14 was significantly enhanced by 15.0 and 33.3% by PSPL as compared to day 0, respectively, while their values were not altered by the LPD. Urinary 8-hydroxy-deoxyguanosine (8-OHdG) excretion decreased significantly by PSPL consumption by 36.7% at day 7 as compared to day 0, yet unchanged by the LPD ( $p \leq 0.05$ ). In conclusion, our results suggest that polyphenols in 200 g PSPL were bio-available and could enhance antioxidant defense and decrease oxidative stress in young healthy people.