

Correlation of Morphological Alterations and Functional Impairment of the Tension-Free Vaginal Tape Obturator Procedure

Yang, JM ; Yang, SH ; Huang, WC
Jenn-Ming Yang; Shwu-Huey Yang; Wen-Chen Huang

Abstract

PURPOSE: We explored the morphological features associated with functional impairment in patients undergoing the tension-free vaginal tape obturator procedure. **MATERIALS AND METHODS:** We retrospectively reviewed the records of 98 women who underwent the tension-free vaginal tape obturator procedure alone or with concomitant pelvic surgery. Postoperative assessment included a symptom questionnaire, ultrasound cystourethrography and a cough stress test. During followup the measures of postoperative functional impairment included a positive cough stress test, new onset voiding dysfunction and the worsening or progression of urge symptoms. **RESULTS:** Median followup was 22 months. During followup 11 women had a positive cough stress test, 22 had voiding dysfunction and 12 had worsening or new onset urge symptoms. Failure was associated with 4 variables on multiple logistic regression analysis, including absent urethral encroachment at rest (OR 16.63, 95% CI 1.87-147.85, $p = 0.01$), bladder neck funneling (OR 8.27, 95% CI 1.99-34.26, $p < 0.01$), a urethral location of less than the 50th percentile (OR 6.01, 95% CI 1.43-25.25, $p = 0.01$) and a resting tape angle of less than 165 degrees (OR 5.21, 95% CI 1.15-23.54, $p = 0.03$). A resting tape distance of less than 12.0 mm (OR 3.00, 95% CI 1.44-6.26, $p < 0.01$) and urethral encroachment at rest (OR 2.86, 95% CI 1.30-6.30, $p < 0.01$) were the variables predictive of postoperative voiding dysfunction. Bladder neck funneling was the only risk factor for postoperative urge symptoms ($p < 0.01$). **CONCLUSIONS:** The tension-free vaginal tape obturator procedure achieves its effectiveness in a process of biological reaction and mechanical interaction between the tape and urethra. When this mechanical interaction is too great or too little, there is functional impairment after the procedure.