

Protective effects of *Ginkgo biloba* extract on the ethanol-induced gastric ulcer in rats

Sheng-Hsuan Chen, Yu-Chih Liang, Jane CJ Chao, Li-Hsueh Tsai, Chun-Chao Chang, Chia-Chi Wang, Shiann Pan

Sheng-Hsuan Chen, Chun-Chao Chang, Shiann Pan, Department and Research Center of Gastroenterology, Taipei Medical University, 252 Wu-Hsing Street, Taipei 110, Taiwan, China
Yu-Chih Liang, Graduate Institute of Biomedical Technology, Taipei Medical University, 250 Wu Hsing Street, Taipei 110, Taiwan, China
Jane CJ Chao, School of Nutrition and Health Sciences, Taipei Medical University, 250 Wu Hsing Street, Taipei 110, Taiwan, China
Li-Hsueh Tsai, Department of Physiology, Taipei Medical University, 250 Wu Hsing Street, Taipei 110, Taiwan, China
Chia-Chi Wang, Department of Gastroenterology, Min-Sheng Hospital, 168 Ching-Kuo Road, Tao-Yuan City 330, Taiwan, China
Supported by the grants of the National Science Council, No. NSC-89-2315-B-038-002 and Min-Sheng Hospital, No. 93 MSH-TMU-05

Correspondence to: Dr. Sheng-Hsuan Chen, Department of Gastroenterology, Taipei Medical University Hospital, 252 Wu-Hsing Street, Taipei 110, Taiwan, China. shchen@tmu.edu.tw
Telephone: +886-2-27372181-3903 Fax: +886-2-27363051
Received: 2004-10-15 Accepted: 2004-12-01

Abstract

AIM: To evaluate the preventive effect of *Ginkgo biloba* extract (GbE) on ethanol-induced gastric mucosal injuries in rats.

METHODS: Female Wistar albino rats were used for the studies. We randomly divided the rats for each study into five subgroups: normal control, experimental control, and three experimental groups. The gastric ulcers were induced by instilling 1 mL 50% ethanol into the stomach. We gave GbE 8.75, 17.5, 26.25 mg/kg intravenously to the experimental groups respectively 30 min prior to the ulcerative challenge. We removed the stomachs 45 min later. The gastric ulcers, gastric mucus and the content of non-protein sulfhydryl groups (NP-SH), malondialdehyde (MDA), c-Jun kinase (JNK) activity in gastric mucosa were evaluated. The amount of gastric juice and its acidity were also measured.

RESULTS: The findings of our study are as follows: (1) GbE pretreatment was found to provide a dose-dependent protection against the ethanol-induced gastric ulcers in rats; (2) the GbE pretreatment afforded a dose-dependent inhibition of ethanol-induced depletion of stomach wall mucus, NP-SH contents and increase in the lipid peroxidation (increase MDA) in gastric tissue; (3) gastric ulcer induced by ethanol produced an increase in JNK activity in gastric mucosa which also significantly inhibited by pretreatment with GbE; and (4) GbE alone had no inhibitory effect on gastric secretion in pylorus-ligated rats.

CONCLUSION: The finding of this study showed that GbE

significantly inhibited the ethanol-induced gastric lesions in rats. We suggest that the preventive effect of GbE may be mediated through: (1) inhibition of lipid peroxidation; (2) preservation of gastric mucus and NP-SH; and (3) blockade of cell apoptosis.

© 2005 The WJG Press and Elsevier Inc. All rights reserved.

Key words: *Ginkgo biloba* extract; Ethanol; Gastric ulcer; c-Jun kinase

Chen SH, Liang YC, Chao JCJ, Tsai LH, Chang CC, Wang CC, Pan S. Protective effects of *Ginkgo biloba* extract on the ethanol-induced gastric ulcer in rats. *World J Gastroenterol* 2005; 11(24): 3746-3750

<http://www.wjgnet.com/1007-9327/11/3746.asp>

INTRODUCTION

Ginkgo biloba, a member of the family Ginkgoaceae, was cultivated in China in the mid-1700s. Originating in the southeastern China some 200 million years ago, it has been living to an age of 1,000 years and it is the last remaining member of its order. Extract from the leaves of *Ginkgo biloba* (GbE) has been used as a traditional Chinese pharmacopeia for centuries in the treatment of asthma and bronchitis^[1]. It is used medically today as a standardized preparation GbE (EGb 761) which contains 240 mg/g flavonoids (ginkgo-flavone glycosides) and 60 mg/g terpenoids (ginkgolides and bilobalides). Those are the most important active ingredients in the extract. The flavonoids act as free radical scavengers, especially for oxygen-derived free radicals, such as OH·, O₂⁻, RO·, and ROO·, and to neutralize ferryl ion-induced peroxidation^[2,3]. The terpenoids is known as an antagonist of platelet-activating factor, which implicates in the processes of platelet aggregation and arterial thrombosis, acute inflammation, allergic reactions and cardiovascular insufficiency^[4,5]. Recently GbE is used to improve cardiovascular circulation and to lessen cerebrovascular insufficiency in western countries clinically^[6].

On the other hand, ethanol is well-known as a damaging agent to gastric mucosa in animal and clinical studies. At concentrations greater than 400 mL/L, it causes marked mucosal hyperemia, necrosis, edema and mucosal or submucosal hemorrhage^[7,8]. The formation of lesions may be mediated by oxygen-derived free radicals^[9,10].

GbE is well-known as a strong free radical scavenger. The gastric mucosal injury after ethanol treatment may be mediated by free radicals. Only a few published studies

showed the GbE protective effects on the ethanol-induced gastric mucosal lesion^[11]. Thus, the present study was intended to evaluate the possible mechanisms of GbE protective effect on the ethanol-induced gastric ulcer in rats.

MATERIALS AND METHODS

Animals and the experimental design

Female Wistar albino rats (Animal Center of National Taiwan University, Taipei, Taiwan) about 180–200 g were used for the study. We maintained the animals in air conditional room with 14/10 h light/dark cycles, fed them with regular chows, and allowed them free access to tap water. Food was deprived but free access to tap water was allowed 36 h before the experiment to ensure an empty stomach.

We purchased *Ginkgo biloba* extract (GbE, Cerenin® ampule, 3.5 mg GbE/mL) from DR. Willmar Schwabe Karlsruhe F.R.G. (Germany), and used normal saline as a vehicle.

The animals were divided into five subgroups for each study. GbE 8.75, 17.5, 26.25 mg/kg were given intravenously to the experimental groups respectively 30 min prior to the ulcerative challenge (50% ethanol, 1 mL) by orogastric gavage to the stomach of fasted rats. Each control and experimental group consisted of five rats.

We sacrificed animals under ether anesthesia 45 min after treatment with ethanol, and removed their stomachs which were opened along the greater curvature and examined for lesions developed in the glandular portion under dissecting microscope (×10) with a square grid. The numbers of ulcer lesions (U. No.) in the glandular portion of the stomach were noted. The ulcer area (mm²) were measured and expressed as the ulcer index (U.I.). We calculated the protective ratio (%) according to the following formula:

$$\text{Preventive ratio (\%)} = (a-b)/a \times 100$$

a: the ulcer index of the control group

b: the ulcer index of the experimental group

Determination of gastric mucus

The glandular stomach was removed and weighted. We transferred the glandular segments immediately to 0.1% Alcian blue solution in 0.16 mol/L sucrose solution with 0.05 mol/L sodium acetate to pH 5.0 and stained for 2 h at room temperature. After having rinsed with sucrose solution, we extracted the dye complexed with the gastric mucus with 0.05 mol/L magnesium chloride solutions. The aliquot of magnesium chloride solution (4 mL) was further extracted with equal volume of diethyl ether and centrifuged (3 600 r/min, 10 min). Then, we calculated the quantity of Alcian blue extracted/g (net) of glandular tissue.

Estimation of non-protein sulfhydryl groups (NP-SH) and malondialdehyde (MDA) contents

After the rats were killed, we opened the glandular stomachs and rinsed them in ice-cold saline, then stored them rapidly in a dry ice bath until analyzed. For the determination of NP-SH, the tissues were homogenized in ice-cold 50% (g/L) aqueous TCA and centrifuged. We determined the NP-SH by measuring the supernatants and 5,5'-dithiobis (2-nitrobenzoic acid) in phosphate buffer (pH 8.0). We read the absorbance

(412 nm) 5 min after it is being incubated.

To determine the MDA, we incubated the supernatants with *N*-methyl-2-phenylindole and we read absorbance at 586 nm according to the manufacturer's instructions (BIOXYTECH LPO-586 kits, OXIS International Inc., Portland, USA).

Determination of anti-secretory activity in pylorus-ligated rats

We fasted animals for 36 h. The following procedures were carried out under ether anesthesia. We gave GbE intravenously 15 min before pylorus ligation, and collected gastric juice 3 h after pylorus ligation. After its volume was measured and expressed in mEq/L, the gastric juice was centrifuged at 3 000 r/min for 10 min. We also determined the total acidity of gastric juice by titrating it with 0.01 mol/L NaOH to pH 7.0.

Western blotting and JNK kinase activity assay

Gastric mucosa was homogenized in Gold lysis buffer (40 mmol/L Tris-NaOH pH 7.5, 500 mmol/L NaCl, 0.1% NP-40, 6 mmol/L EDTA, 6 mmol/L EGTA, 10 mmol/L β-glycerophosphate, 10 mmol/L NaF, 10 mmol/L PNPP, 300 μmol/L sodium orthovanadate, 1 mmol/L benzamidine, 2 μmol/L PMSF, 10 mg/mL aprotinin, 1 μg/mL leupeptin, and 1 mmol/L DTT) and centrifuged. We collected the supernatant as total tissue lysate. Equal amounts of total tissue lysate (50 μg) were resolved by SDS-polyacrylamide gel electrophoresis (PAGE), and transferred onto Immobilon-P membrane (Millipore, Bedford, MA) as described previously^[12]. We also incubated the membrane with an anti-JNK1 antiserum (Transduction Laboratories, Lexington, KY). The membranes were subsequently probed with anti-mouse IgG antibody conjugated to horseradish peroxidase (Santa Cruz Biotechnology) and visualized using enhanced chemiluminescence's kits (ECL, Amersham). For kinase assay, we immunoprecipitated equal amounts of total tissue lysate (400 μg) with JNK1 specific antibody and protein A/G-PLUS agarose for 15 h at 4 °C. Kinase assay was carried out in 45 μL of kinase buffer (40 mmol/L Tris-NaOH pH 7.5, 500 mmol/L NaCl, 0.1% NP-40, 6 mmol/L EDTA, 6 mmol/L EGTA, 10 mmol/L β-glycerophosphate, 10 mmol/L NaF, 10 mmol/L PNPP, 300 μmol/L sodium orthovanadate, 1 mmol/L benzamidine, 2 μmol/L PMSF, 10 μg/mL aprotinin, 1 μg/mL leupeptin, and 1 mmol/L DTT) containing 5 μmol/L cold ATP, 10 μCi [γ-³²P] ATP (5 000 Ci/mmol, Amersham), and 1 μg GST-c-Jun fusion protein (Santa Cruz Biotechnology) as substrate, and incubated for 20 min at 25 °C. We mixed each sample with 8 μL of 5× Laemmli's loading buffer to stop the reaction, heated for 10 min at 100 °C, and subjected to 8% SDS-PAGE. The gels were dried, visualized by autoradiography^[12].

Statistical analysis

Results were expressed as mean ± SE for each experiment. We analyzed the data with Student's *t*-test. If a *P* value was less than 0.05, the differences were considered statistically significant.

RESULTS

Effect of GbE on ethanol-induced gastric mucosal damage

Table 1 Effect of *Ginkgo biloba* extract on the induction of gastric ulcers by 50% ethanol in rats¹

Treatment and dose	n	Mucosal lesions ²		Preventive ratio (%)
		U. Number	U.I. (mm ²)	
Control (NS)	5	0	0	
NS + 50% EtOH (1 mL/rat)	5	10±4	20.31±4.23	
GbE (8.75 mg/kg)+50% EtOH	5	8±3	15.25±3.21	24.91
GbE (17.5 mg/kg)+50% EtOH	5	3±3 ^a	8.83±3.59 ^a	56.52
GbE (26.25 mg/kg)+50% EtOH	5	2±2 ^a	6.37±4.22 ^a	68.64

¹NS, normal saline; GbE, *Ginkgo biloba* extract; EtOH, ethanol; U. No., ulcer number; U.I., ulcer index. ²Each value represents the mean±SE. Significantly different from vehicle with ethanol treatment (^a*P*<0.05).

Table 1 shows all the rats that received 1 mL 50% ethanol administration had induced gastric mucosal damages. GbE (8.75-26.25 mg/kg, intravenously) suppressed the ethanol-induced gastric lesions including ulcer number (U. No.) and the ulcer index (U.I.) in a dose-dependent manner. As shown in Table 1, normal saline did not have any protective effect on gastric mucosal damage when given in the volume of 0.5-1.5 mL to rats intravenously. The maximal effect was obtained when GbE (26.25 mg/kg) was given 30 min prior to ethanol induction, resulting in a preventive ratio of 68.64% (*P*<0.05).

Effect of GbE on ethanol-induced gastric mucus

There was a significant decrease in the gastric mucus after treatment with 1 mL 50% ethanol. As shown in Table 2, pretreatment of GbE (8.75-26.25 mg/kg) significantly protected the decline in the gastric mucus levels which were induced by ethanol.

Table 2 Effect of *Ginkgo biloba* extract on the induction of changes in gastric wall mucus by gavages with 50% ethanol¹

Treatment and dose	n	Gastric wall mucus (μg Alcian blue/g wet tissue, mean±SE)
Control (NS)	5	442.21±28.36
NS+50% EtOH (1 mL/rat)	5	356.87±30.67 ^a
GbE (8.75 mg/kg)+50% EtOH	5	348.87±32.37
GbE (17.5 mg/kg)+50% EtOH	5	395.27±25.91
GbE (26.25 mg/kg)+50% EtOH	5	410.82±29.16 ^c

¹NS, normal saline; GbE, *Ginkgo biloba* extract; EtOH, ethanol. Significantly different from the control (NS) (^a*P*<0.05). Significantly different from the ethanol treatment (^c*P*<0.05).

Effect of GbE on ethanol-induced gastric mucosal NP-SH and MDA contents

Ethanol (50%, 1 mL) treatment significantly reduced the NP-SH concentration in the gastric mucosa as compared with control in rats. As shown in Table 3, GbE pretreatment significantly prevented the decrease in NP-SH concentrations in the dose of 8.75-26.25 mg/kg. However, during the GbE treatment alone there was no change in the concentration of NP-SH in gastric mucosa. Moreover, pretreatment of GbE significantly prevented lipid peroxidation induced by ethanol, the MDA concentrations were 209.27±10.48 or 118.82±8.26 in the rats that were treated with ethanol alone or combined with GbE, respectively (Table 4).

Table 3 Effect of *Ginkgo biloba* extract on the levels of NP-SH in the stomachs of rats treated by gavages with 50% ethanol¹

Treatment and dose	n	NP-SH concentrations (μg/100 mg /wet tissue, mean±SE)
Control (NS)	5	12.21±1.03
NS+50% EtOH (1 mL/rat)	5	2.82±1.32 ^a
GbE (8.75 mg/kg)+50% EtOH	5	5.03±1.25 ^c
GbE (17.5 mg/kg)+50% EtOH	5	5.66±1.82 ^c
GbE (26.25 mg/kg)+50% EtOH	5	11.32±2.01 ^c

¹NS, normal saline; GbE, *Ginkgo biloba* extract; EtOH, ethanol; NP-SH, non-protein sulfhydryls. Significantly different from the control (normal saline, ^a*P*<0.05). Significantly different from the ethanol treatment (^c*P*<0.05).

Table 4 Effect of *Ginkgo biloba* extract on the lipid peroxidation in the stomachs of rats treated by gavages with 50% ethanol¹

Treatment and dose	n	Malondialdehyde concentration (nmol/g wet tissue, mean±SE)
Control (NS)	5	102.54±15.32
NS+50% ethanol (1 mL/rat)	5	209.27±10.48 ^a
GbE (8.75 mg/kg)+50% EtOH	5	212.32±9.83
GbE (17.5 mg/kg)+50% EtOH	5	125.48±11.39 ^c
GbE (26.25 mg/kg)+50% EtOH	5	118.82±8.26 ^c

¹NS, normal saline; GbE, *Ginkgo biloba* extract; EtOH, ethanol. Significantly different from the control (normal saline) (^a*P*<0.05). Significantly different from the ethanol treatment (^c*P*<0.05).

Effect of GbE in gastric acid secretion and acidity on pylorus-ligated rats

GbE was intravenously administered at a dose of 8.75-26.25 mg/kg 15 min before pylorus ligation. Control rats received the same volume of saline (0.5-1.5 mL). As shown in Table 5, GbE could not significantly decrease the gastric acid secretion and only slightly inhibited the gastric secretion and total acidity even at a dose of 26.25 mg/kg.

Effect of GbE on ethanol-induced JNK activity in gastric mucosa

We examined c-Jun kinase (JNK) activity by using an immunocomplex kinase assay to explore whether the JNK signaling pathway is activated within gastric mucosa in response to ethanol, since JNK activation plays an important role in the induction of cell apoptosis. As shown in Figure 1, JNK activity was obviously increased after treatment with ethanol (top). However, GbE (26.25 mg/kg) significantly suppressed the induction of JNK activity. Western blot analysis revealed that this JNK activation was not caused by the enhanced expression of JNK protein (bottom).

Table 5 Effect of *Ginkgo biloba* extract on the gastric secretion and acidity in pylorus-ligated rats¹

Treatment and dose	n	Volume of gastric secretion (mL) ²	Titration acidity (mEq/L) ²
Control (NS)	5	13.35±0.98	63.82±8.83
GbE (8.75 mg/kg)	5	12.67±0.81	61.98±7.34
GbE (17.5 mg/kg)	5	11.98±0.53	58.14±9.28
GbE (26.25 mg/kg)	5	10.78±0.91	56.28±8.91

¹NS, normal saline; GbE, *Ginkgo biloba* extract; EtOH, ethanol. ²Each value repressed with mean±SE.




Figure 1 Effect of GbE on ethanol-induced JNK activity in gastric mucosa of rats. JNK activity was obviously increased after treatment with ethanol (top). However, GbE (26.25 mg/kg) significantly suppressed the induction of JNK activity. Western blot analysis revealed that this JNK activation was not caused by enhanced expression of JNK protein (bottom). GbE, *Ginkgo biloba* extract; EtOH, ethanol.

DISCUSSION

In the past two decades, the protective factors of gastric mucosa were well evaluated. In 1979, Boyd *et al.*, found that the gastric mucosa contains high concentrations of reduced glutathione, the major component of the endogenous NP-SH pool^[13]. In 1981, Szabo *et al.*, showed the protective role of NP-SH in noxious-induced gastric injury^[3]. NP-SH pool might involve in scavenging oxygen-derived free radicals^[14,15] and influence the production and character of mucus^[16,17].

Mucus, which continuously coats over the gastric mucosa, is well known as a “mucous barrier” to prevent the injury of luminal acid, bacteria and noxious agents injuries^[18,19]. Mucus might implicate in scavenging oxygen-derived free radicals^[20,21]. Mucus glycoproteins and lipids bound to mucin might involve in the antiradical process^[22,23].

On the other hand, previous reports offered that ethanol-induced gastric mucosal lesions may be attributed to the possible mechanisms: (1) increase oxygen-derived free radicals^[9,10,24,25], (2) decrease the concentration of NP-SH contents in gastric mucosa^[3,15], (3) direct damage to the mucin layer or mucin synthesis^[26], and (4) causing gastric cell's apoptosis^[27-29].

GbE can remove free radicals especially oxygen-derived free radicals such as OH·, O₂⁻, RO·, and ROO·, and neutralize ferryl ion-induced peroxidation. GbE can improve the cardiovascular and cerebrovascular diseases e.g. Alzheimer's disease, cerebral insufficiency and depression^[6,30]. Furthermore, GbE has anti-inflammatory effect by decreasing the production of active oxygen and nitrogen species^[31,32]. Recent reports showed that GbE can improve duodenal ulcer healing and reverses CCl₄-induced liver fibrosis.

The results of the present study showed that pretreatment with GbE not only kept the mucus integrity (Table 2) but

also inhibited ethanol-induced depletion in the NP-SH concentrations (Table 3) and MDA production in gastric mucosa (Table 4). All those findings in our study suggest that GbE has its role of being a free radical scavenger, decreases the lipid peroxidation, and blocks the loss of mucus, NP-SH, resulting in having protective effect on ethanol-induced mucosal injury.

Apoptosis has been implicated in causing ethanol-induced gastric mucosal injury^[27-29]. Previous studies indicated that JNK kinase activity was elevated during the process of apoptosis and blocking of JNK activity was able to prevent cell apoptosis. The data from this study showed that ethanol significantly increased the JNK kinase activity, resulting in cell apoptosis and gastric mucosal damage. As shown in Figure 1, blocking of JNK activity may contribute to GbE's gastric mucosal protection.

The finding of this study further showed that GbE did not significantly decrease the gastric acid secretion (Table 5). Our finding here is similar to the report by Wang *et al.* in 2000^[11].

In conclusion, the data of this study suggest that GbE can inhibit ethanol-induced gastric lesions in rats. The possible mechanisms of GbE's antiulcer benefit may be due to its oxygen radicals scavenging by inhibition of lipid peroxidation, preventing the loss of gastric mucus and NP-SH, and blockade of ethanol-induced apoptosis. Further studies are warranted to evaluate GbE at the pharmacological effective dosage before any consideration for clinical trials.

REFERENCES

- 1 Kleijnen J, Knipschild P. *Ginkgo biloba*. *Lancet* 1992; **340**: 1136-1139
- 2 Pincemail J, Dupuis M, Nasr C, Hans P, Haag-Berrurier M, Anton R, Deby C. Superoxide anion scavenging effect and superoxide dismutase activity of *Ginkgo Biloba* extract. *Experientia* 1989; **45**: 708-712
- 3 Gardès-Albert M, Ferradini C, Sekaki A, Droy-Lefaix MT. Oxygen-centered free radicals and their interactions with EGb 761 or CP 202. In: Ferradini C, Droy-Lefaix MT, Christen Y, eds. *Advances in Ginkgo biloba extract research: Ginkgo biloba extract (EGb 761) as a free-radical scavenger*. *New York Elsevier Science* 1993: 1-11
- 4 Akiba S, Kawauchi T, Oka T, Hashizume T, Sato T. Inhibitory effect of the leaf extract of *Ginkgo biloba* L. on oxidative stress-induced platelet aggregation. *Biochem Mol Biol Int* 1998; **46**: 1243-1248
- 5 Akisu M, Kultursay N, Coler I, Huseyinov A. Platelet-activating factor is an important mediator in hypoxic ischemia brain injury in the newborn rat. Flunarizine and *Ginkgo biloba* extract reduce PAF concentration in the brain. *Biol Neonate* 1998; **74**: 439-444
- 6 Kleijnen J, Knipschild P. *Ginkgo biloba* for cerebral insufficiency. *Br J Clin Pharmacol* 1992; **34**: 352-358
- 7 Oates PJ, Hakkinen JP. Studies on the mechanism of ethanol-induced gastric damage in rat. *Gastroenterology* 1988; **94**: 10-21
- 8 Szabo S, Goldberg I. Experimental pathogenesis: drugs and chemical lesions in the gastric mucosa. *Scand J Gastroenterol Suppl* 1990; **174**: 1-8
- 9 Pihan G, Regillo C, Szabo S. Free radicals and lipid peroxidation in ethanol- and aspirin-induced gastric mucosal injury. *Dig Dis Sci* 1987; **32**: 1395-1401
- 10 Szelenyi I, Brune K. Possible role of oxygen free radicals in ethanol-induced gastric mucosal damage in rats. *Dig Dis Sci* 1988; **33**: 865-871

- 11 **Wang Q**, Zhao WZ, Ma CG. Protective effects of *Ginkgo biloba* extract on gastric mucosa. *Acta Pharmacol Sin* 2000; **21**: 1153-1156
- 12 **Liang YC**, Huang YT, Tsai SH, Lin-Shiau SY, Chen CF, Lin JK. Suppression of inducible cyclooxygenase and inducible nitric oxide synthase by apigenin and related flavonoids in mouse macrophages. *Carcinogenesis* 1999; **20**: 1945-1952
- 13 **Boyd SC**, Sasame HA, Boyd MR. High concentrations of glutathione in glandular stomach: possible implications for carcinogenesis. *Science* 1979; **205**: 1010-1012
- 14 **Videla LA**, Valenzuela A. Alcohol injection, liver glutathione and lipid peroxidation: metabolic interaction and pathological implication. *Life Sci* 1982; **31**: 2395-2407
- 15 **Li T**, Zhang XJ. Involvement of sulfhydryls in the protective mechanism of gastric mucosa. *Acta Physiol Sinica* 1990; **42**: 571-577
- 16 **Allen A**, Cunliffe WJ, Pearson JP, Sellers LA, Ward R. Studies on gastrointestinal mucus. *Scand J Gastroenterol Suppl* 1984; **93**: 101-113
- 17 **Salim AS**. Sulfhydryl-containing agents: new approach to the problem of refractory peptic ulceration. *Pharmacology* 1993; **46**: 281-288
- 18 **Garner A**, Flemstrom G, Allen A, Heylings JR, McQueen S. Gastric mucosal protective mechanisms: role of epithelial bicarbonate and mucus secretions. *Scand J Gastroenterol Suppl* 1984; **101**: 79-86
- 19 **Farre AJ**, Colombo M, Alvarez I, Glavin GB. Some novel 5-hydroxytryptamine 1A (5-HT_{1A}) receptor agonists reduce gastric acid and pepsin secretion, reduce experimental gastric mucosal injury and enhance gastric mucin in rats. *J Pharmacol Exp Ther* 1995; **272**: 832-837
- 20 **Grisham MB**, Von Ritter C, Smith BF, Lamont JT, Granger DN. Interaction between oxygen radicals and gastric mucin. *Am J Physiol* 1987; **253**(1 Pt 1): G93-96
- 21 **Cross CE**, Halliwell B, Allen A. Antioxidant protection: a function of tracheobronchial and gastrointestinal mucus. *Lancet* 1984; **1**: 328-330
- 22 **Gong DH**, Turner B, Bhaskar KR, Lamont JT. Lipid binding to gastric mucin: protective effect against oxygen radicals. *Am J Physiol* 1990; **259**(4 Pt 1): G681-686
- 23 **Hiraishi H**, Terano A, Ota S, Mutoh S, Sugimoto T, Hirada T, Razandi M, Ivey KJ. Role for mucus glycoprotein in protecting cultured gastric mucosal cells against toxic oxygen metabolites. *J Lab Clin Med* 1993; **121**: 570-578
- 24 **Terano A**, Hiraishi H, Ota S, Shiga J, Sugimoto T. Role of superoxide and hydroxyl radicals in rat gastric mucosal injury induced by ethanol. *Gastroenterol Jpn* 1989; **24**: 488-493
- 25 **Mutoh H**, Hiraishi H, Ota S, Ivey KJ, Terano A, Sugimoto T. Role of oxygen radicals in ethanol-induced damage to cultured gastric mucosal cells. *Am J Physiol* 1990; **258**(4 Pt 1): G603-609
- 26 **Slomiany A**, Morita M, Sano S, Piotrowski J, Skrodzka D, Slomiany BL. Effect of ethanol on gastric mucus glycoprotein synthesis, translocation, transport, glycosylation and secretion. *Alcohol Clin Exp Res* 1997; **21**: 417-423
- 27 **Piotrowaki J**, Piotrowaki E, Strodzka D, Slomiany A, Slomiany BL. Gastric mucosal apoptosis induced by ethanol: effect of antiulcer agents. *Biochem Mol Biol Int* 1997; **42**: 247-254
- 28 **Mizushima T**, Tsutsumi S, Rokutan K, Tsuchiya T. Suppression of ethanol-induced apoptotic DNA fragmentation by geranylgeranylacetone in cultured guinea pig gastric mucosal cells. *Dig Dis Sci* 1999; **44**: 510-514
- 29 **Hoshino T**, Takano T, Tsutsumi S, Tomisato W, Tsuchiya T, Mizushima T. Effects of prostaglandin E₂ on gastric irritant-induced apoptosis. *Dig Dis Sci* 2002; **47**: 2370-2379
- 30 **Marcocci L**, Maguire JJ, Dory-Lefaix L. The nitric oxide-scavenging properties of *Ginkgo biloba* extract Egb761. *Biochem Biophys Res Commun* 1994; **201**: 748-755
- 31 **Kim HK**, Son KH, Chang HW, Kang SS, Kim HP. Inhibition of rat adjuvant-induced arthritis by ginkgetin, a biflavone from ginkgo biloba leaves. *Planta Med* 1999; **65**: 465-467
- 32 **Yoshikawa T**, Naito Y, Kondo M. Ginkgo biloba leaf extract: review of biological actions and clinical applications. *Antioxid Redox Signal* 1999; **1**: 469-480

Science Editor Li WZ Language Editor Elsevier HK