

Serum β -carotene level, arsenic methylation capability, and incidence of skin cancer.

邱弘毅

Hsueh YM;Chiou HY;Huang YL;Wu WL;Huang CC;Yang MH;Lue LC;Chen GS and
Chen CJ

Abstract

This study presents the distribution of arsenic (As), copper (Cu), and zinc (Zn) in various seafoods-oysters (*Crassostrea gigas*), false fucus (*Hemifusus tuba*), venus clams (*Cyclina sineasis*), green mussels (*Perna viridis*), blood clams (*Arca granosa*), flounders (*Psettodes erumei*), and rock-shells (*Thais clavigera*) collected from the Putai coast of the black-foot disease (BFD) area in Taiwan. Special attention is paid to evaluate the relationships among As, Cu, and Zn and effect of body size on metal concentration in *Thais clavigera*. Maximum Zn and Cu geometric mean (GM) concentrations (GM = 615 and 376 microg/g, dry wt, respectively) are found in oysters (*Crassostrea gigas*), and the values are much higher than those of the other organisms by about 1.7-208 and 1.8-375 times, respectively. Similarly, *Thais clavigera* has a high capacity for accumulating Cu and Zn collected from the same location. One interesting point is that relatively high As concentrations (GM = 65.7 microg/g, dry wt) in *Thais clavigera* are found as compared with those in other organisms (range from GM = 2.37 to 40.2 microg/g, dry wt). The As concentrations are significantly higher in *Thais clavigera* (1.62-27.7 times) than those in other organisms ($p < 0.05$), except for the false fucus (*Hamifusus tuba*). A linear regression analysis shows a significant increase in Zn concentration with increasing Cu concentration in *Thais clavigera*. On the other hand, the As concentration is correlated with Cu and Zn concentrations ($r = 0.77$ and 0.77 , respectively; $p < 0.05$) in *Thais clavigera*. Double logarithmic plots of metal content and concentration against dry-body weight and shell length show linear relationships. The result indicates that large individuals have higher contents of Cu, Zn and As, and have slopes of 1.58, 1.38, and 1.34, respectively. In addition, metal concentrations against shell length for all animal sizes also indicate that Cu, Zn and As have slopes of 1.92, 1.18, and 1.11, respectively. In conclusion, *Thais clavigera* has a high capacity for accumulating As, Cu, and Zn and is a potential bioindicator for monitoring As, Cu and Zn.