

Antioxidative natural product protect against econazole-induced liver injuries. .(196):87-93

林永和

劉吉豐(Liu CF);林佳賢(Lin CH);林俊清(Lin CC);林永和
(Lin YH);陳金發(Chen CF);林昭光(Lin CK);林松洲(Lin
SC)

摘要

Abstract

The study objective of this research is in order to investigate the hepatoprotective and therapeutic effects of propolis ethanol extract (PEE) on acute econazole-induced liver injury. Positive control of various concentrations of PEE on liver function and the dose-response relationship of liver injury induced by various doses of econazole were firstly observed from biochemical assay of serum level of aspartate transaminase (SGOT) and serum alanine transaminase (SGPT) and histopathological microscopic examination. The hepatoprotective effects of various concentration of PEE on liver damage induced by hepatotoxic dose (300 mg/kg) of econazole were observed by the obvious decrement of SGOT and SGPT level and further confirmed by hepatohistological microscopic examination. The inhibitory effects of PEE on FeCl₂-induced (in vitro) or econazole-induced (in vivo) lipid peroxidation were investigated from the measurement of the formed malonic dialdehyde (MDA) level in the rat liver homogenate. The IC₅₀ (microM) of various concentrations of PEE in the superoxide scavenging activity in econazole (300 mg/kg)-damaged rat liver homogenate were assessed by cytochrome c reduction method and compared with that of (+)-alpha-tocopherol. It could be postulated that the hepatoprotective effect of PEE may be, at least in part, due to their inhibitory ability on membrane lipid peroxidation and free radical formation or due to their free radical scavenging ability.