

Is coronally positioned flap procedure adjunct with enamel matrix derivative or root conditioning a relevant predictor for achieving root coverage? A systemic review

Y.-F. Cheng¹, J.-W. Chen³,
S.-J. Lin⁴, H.-K. Lu^{1,2}

¹Department of Periodontology, College of Oral Medicine, Taipei Medical University, Taipei, Taiwan, ²Periodontal Clinics of Dental Department, Taipei Medical University Hospital, Taipei, Taiwan, ³Department of Pediatric Dentistry, University of Texas Health Science Center, Houston, TX, USA and ⁴Shin-Kong Wu Ho-Su Memorial Hospital, Taipei, Taiwan

Cheng Y-F, Chen J-W, Lin S-J, Lu H-K. Is coronally positioned flap procedure adjunct with enamel matrix derivative or root conditioning a relevant predictor for achieving root coverage? A systemic review. *J Periodont Res* 2007; 42: 474–485. © 2007 The Authors. Journal compilation © 2007 Blackwell Munksgaard

Background and Objective: This study is a systemic review of coronally positioned flap, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + enamel matrix derivative (EMD) for the treatment of Miller class I and II gingival recession.

Material and Methods: All studies available through the Medline database by the end of October 2005 were used. Each study provided mean clinical attachment level, keratinized tissue, probing pocket depth, gingival recession depth and root coverage percentage before and after treatment with coronally positioned flap alone, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + EMD. Effectiveness was evaluated by comparing the weighted mean average in gingival recession depth, probing pocket depth, clinical attachment level, keratinized tissue and root coverage percentage achieved with the three treatments.

Results: Seven studies for the coronally positioned flap + EMD group, four studies for the coronally positioned flap + chemical root surface conditioning group, and seven studies for the coronally positioned flap group were retrieved for this weighted mean analysis. The results of clinical attachment level, gingival recession depth, and root coverage percentage in the coronally positioned flap + EMD group were statistically significantly better than the changes in the coronally positioned flap and coronally positioned flap + chemical root surface conditioning group at 6 and 12 mo ($p < 0.001$). There was no significant difference at the 6-mo comparison among clinical attachment level, keratinized tissue, probing pocket depth, and gingival recession depth, except in the root coverage percentage for coronally positioned flap and coronally positioned flap + chemical root surface conditioning groups.

Prof. Hsein-Kun Lu, Department of Periodontology, College of Oral Medicine, Taipei Medical University and Taipei Medical University Hospital, 250 Wu-Hsing Street, Taipei 110, Taiwan
Tel: +886 227361661 ext. 5110
Fax: +886 227362295
e-mail: jackson@tmu.edu.tw

Key words: coronally positioned flap; enamel matrix derivative; root coverage

Accepted for publication October 23, 2006

Conclusion: The results suggest that root coverage by the coronally positioned flap and coronally positioned flap + chemical root surface conditioning procedures were unpredictable but became more predictable when the coronally positioned flap procedure was improved by the modification of adding EMD.

Gingival recession is defined as the location of marginal periodontal tissues apical to the cemento–enamel junction (1). An ideal periodontal plastic surgery procedure for root coverage should re-establish esthetics and function, and provide a sulcus exhibiting no bleeding on probing and a depth of ≤ 2 mm (2). The coronally positioned flap has also been shown to be effective in covering recession-type defects (3,4). However, as gingival fibroblasts tend to repopulate the root surface faster than periodontal ligament cells, healing will generally not lead to the formation of a functional periodontal ligament but instead to new attachment. In fact, there is some risk of root resorption in roots directly exposed to gingival fibroblasts during healing (5). Enamel matrix derivative (EMD; Emdogain[®]; Biora AB, Malmö, Sweden), harvested from embryonic porcine teeth, has been extensively studied in animals and humans, and has provided evidence of tissue regeneration (6–9). Numerous studies have reported that EMD promotes the formation of the acellular cementum that attaches to the dentin and alveolar bone (6,7,10,11). Three human biopsy reports revealed that true periodontal regeneration could be achieved with the topical application of EMD (12–14).

Recently, clinical studies have shown that it is possible to apply EMD adjunct to coronally positioned flap procedures (15–17) to achieve root coverage, as well as periodontal regeneration on a previously exposed root surface. Cueva *et al.* recently reported significant increases in the root coverage percentage and keratinized tissue 6 mo after surgery, in marginal tissue recessions (Miller classes I, II, and III) treated with coronally positioned flap + EMD compared with coronally positioned flap without EMD (18). However, Modica *et al.* observed that root coverage and the

clinical attachment level were slightly (but not significantly) improved when using coronally positioned flap + EMD compared with coronally positioned flap alone in the treatment of gingival recession (19). There are conflicting results regarding the clinical benefit of the combined procedure. Based on this evidence, it seems that using EMD in combination with coronally positioned flap or coronally positioned flap alone for treating Miller class I and II recession defects is still controversial.

Chemical root surface conditioning has been introduced, using a variety of agents, in order to detoxify, decontaminate and demineralize the root surface, thereby removing the smear layer and exposing the collagenous matrix of dentin and cementum (20–23). Various acids have been used for chemical root surface conditioning, including citric and phosphoric acids (24), ethylenediaminetetraacetic acid (EDTA) (25) and tetracycline hydrochloride (26). These procedures in an animal model are believed to be able to induce cementogenesis and enhance attachment either by connective tissue ingrowth and/or demineralization (27,28). However in human studies, no clinical advantages were observed (29,30). The clinical relevance of root conditioning with an acid agent in routine periodontal surgery is still questionable.

There being no explicit critical appraisal has created controversy and confusion about the positive or negative uses of these agents. Therefore, the aim of the present systematic review was to assess the efficacy of EMD and root conditioning on the root coverage with coronally positioned flap, in terms of changes in clinical attachment level, keratinized tissue, probing pocket depth, gingival recession depth, and root coverage percentage.

This systematic review follows a well-defined protocol whereby a clearly

formulated question is addressed using systematic and specific methods to identify, select, critically appraise and summarize relevant research. It may provide clinicians with an expanded and unbiased appraisal of human experimental studies. The purpose of this report was to present the results of the searched papers, and to evaluate the methods and quality of the systematic reviews, in order to facilitate clinical decision-making in the choice of coronally positioned flap alone, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + EMD for the treatment of gingival recession.

Material and methods

Strategy for data collection

Studies in the medical literature, available through the Medline database by the end of October 2005, were screened for this meta-analysis. We used the following interventions: coronally positioned flap, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + EMD, for the treatment of Miller class I and II gingival recession (Fig. 1). Only full-length original journal articles were considered; abstracts or unpublished studies were not included. Hand searching of journals for missed trials was not carried out. The search was restricted to studies published in English language journals and those conducted on human subjects. The contents of full-text articles identified during the literature search were reviewed to determine whether they met the inclusion criteria.

Criteria for including studies

The following inclusion criteria were set after detailed discussion between two reviewers:

- human trials;




Fig. 1. The flow for selecting the articles. CPF, coronally positioned flap; EMD, enamel matrix derivative; RC, chemical root surface conditioning.

- patients with a clinical diagnosis of gingival recession of > 2 mm (class I and II according to the Miller classification);
- treatment with coronally positioned flap, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + EMD;
- a randomized controlled trial, controlled clinical trial, or care series report with at least a 6- to 12-mo interval between the initial and final measurements; and
- baseline and final measurements of buccal recession depth.

Criteria for excluding studies

Exclusion criteria for these procedures included:

- lack of any of the outcome variables at the baseline or on the final visit;
- animal studies;
- abstracts;
- histological studies;
- studies with insufficient data;
- laterally moved, coronally advanced flap; and
- root coverage with the design of a semilunar coronally positioned flap.

Quality assessment

Two independent reviewers screened titles and abstracts to identify eligible studies and reviewed a full text of studies to assess their suitability for inclusion in a systematic review database. Any disagreement in the choice of studies of possible relevance was resolved by discussion among the reviewers. Three aspects were analyzed:

- the adequacy of the method of randomization;
- the existence of blinding of the examiners for the variable type of treatment; and
- the existence and treatment of lost cases.

After a preliminary evaluation of the selected articles, considerable heterogeneity was found in the study methodologies, characteristics of the included patients, types of treatments provided, outcome variables registered and results. Nevertheless, it was still possible to make a quantitative synthesis of the data and the consequent weighted mean analysis following the predetermined flow chart (Fig. 1).

Data extraction and outcome measures

For each trial, the following data were recorded: year of publication, details on the type of defects, the number of subjects, the chemical root surface modifier, details of the outcomes report and time intervals. A data extraction sheet was used to collect information that was then entered into a database. Primary outcome measures included changes in clinical attachment levels, keratinized tissue, probing pocket depths, gingival recession depths and root coverage percentage. Mean values were used for continuous data for primary variables.

Data analyses

Data from studies were combined in order to evaluate the treatment effect of coronally positioned flap, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + EMD for root coverage. Each study provided the mean clinical attachment level, keratinized tissue, probing pocket depth, gingival recession depth, and root coverage percentage before and after treatment with coronally positioned flap, coronally positioned flap + chemical root surface conditioning, or coronally positioned flap + EMD. Effectiveness was evaluated by comparing the average changes in clinical attachment level, keratinized tissue, probing pocket depth, gingival recession depth and root coverage percentage with these three treatments. Analysis was performed by using a random-effects model, and the results are expressed as weighted mean difference with 95% confidence interval. A weighted mean difference method (Weighted mean = $\Sigma[(\text{Mean}1.n1/n1 + n2 + \dots + nf) + (\text{Mean}2.n2/n1 + n2 + \dots + nf) + \dots + (\text{Mean}f.nf/n1 + n2 + \dots + nf)]$) was used to pool continuous data of the relevant outcomes. In studies where the standard deviation of the studied outcome data was not pooled, statistical analysis was carried out with SPSS software (version 12.0; SPSS, Chicago, IL, USA) to fit the models. The results of the systemic review were analyzed as weighted mean differences and stand-

ard deviations. The effect size was estimated using the reported *p*-values. A *p*-value of < 0.001 was considered statistically significant.

Results

Study characteristics

According to the flow chart shown in Fig. 1, our search provided 100 potentially relevant publications for coronally positioned flap + EMD and 317 for coronally positioned flap alone. Of these, 89 for coronally positioned flap + EMD and 291 for coronally positioned flap alone were clearly not relevant to the review because they addressed completely different research questions, involved animal research only, or they were review articles after screening titles and abstracts. Eleven studies for coronally positioned flap + EMD and 26 studies for coronally positioned flap were retrieved for detailed evaluation. After review of the full text of the coronally positioned flap + EMD articles, two studies did not meet the inclusion criteria (18,31), leaving seven trials that were appropriate for the meta-analysis (Table 1) (15-17, 19, 32-34). In 26 trials for coronally positioned flap, 11 publications were available for data abstraction. Of these, three studies used tetracycline for root conditioning and one study used EDTA for root conditioning (Table 2) (19, 33, 35-43). Fifteen articles were excluded following the exclusion criteria set up for coronally positioned flap and coronally positioned flap + etching groups (18,44-57).

Clinical attachment level

At 6 mo, the mean gains in clinical attachment level were 2.42 ± 0.70 mm (chi-square for heterogeneity: 88.925 ± 6 , $p < 0.05$) in the coronally positioned flap group, 2.22 ± 0.36 mm (chi-square for heterogeneity: 0.182 ± 2 , $p = 0.913$) in the coronally positioned flap + chemical root surface conditioning group, and 4.01 ± 0.77 mm (chi-square for heterogeneity: 1.000 ± 2 , $p = 0.607$) in the coronally positioned flap + EMD group (Table 3). At 12 mo, the mean gains in

clinical attachment level were 1.69 ± 0.15 mm (chi-square for heterogeneity: 2.613 ± 1 , $p = 0.106$) in the coronally positioned flap group, 3.10 ± 0.00 mm in the coronally positioned flap + chemical root surface conditioning group, and 3.61 ± 0.50 mm (chi-square for heterogeneity: 24.303 ± 3 , $p < 0.05$) in the coronally positioned flap + EMD group (Table 4). At 6 mo, analysis of variance measures demonstrated a statistically significant difference ($p < 0.001$) between the coronally positioned flap + EMD group compared with the coronally positioned flap group and coronally positioned flap + chemical root surface conditioning group. The change in the coronally positioned flap + EMD group was statistically significantly better than those in the coronally positioned flap + chemical root surface conditioning and coronally positioned flap groups at 6 mo. The mean gain in clinical attachment level in the coronally positioned flap group was better than that in the coronally positioned flap + chemical root surface conditioning group, but did not differ statistically significantly at 6 mo ($p = 0.111$). At 12 mo, differences in the gain in clinical attachment level of root coverage were statistically significant ($p < 0.001$) among all groups. The change in the coronally positioned flap + EMD group was better than that in the coronally positioned flap + chemical root surface conditioning group, and the change in the coronally positioned flap + chemical root surface conditioning group was better than that in the coronally positioned flap group at 12 mo.

Keratinized tissue

At 6 mo, the mean gains in keratinized tissue were -0.04 ± 0.37 mm (chi-square for heterogeneity: 87.898 ± 5 , $p < 0.05$) in the coronally positioned flap group, 0.14 ± 0.47 mm (chi-square for heterogeneity: 0.182 ± 2 , $p = 0.913$) in the coronally positioned flap + chemical root surface conditioning group, and 0.59 ± 0.20 mm (chi-square for heterogeneity: 11.164 ± 2 , $p < 0.05$) in the coronally positioned flap + EMD group (Table 3). At

12 mo, the mean gains in keratinized tissue were 0.10 ± 0.41 mm (chi-square for heterogeneity: 2.613 ± 1 , $p = 0.106$) in the coronally positioned flap group, 0.30 ± 0.00 mm in the coronally positioned flap + chemical root surface conditioning group, and 0.61 ± 0.14 mm (chi-square for heterogeneity: 53.647 ± 2 , $p < 0.05$) in the coronally positioned flap + EMD group (Table 4). The change in the coronally positioned flap + EMD group was statistically significantly better than those in the coronally positioned flap and coronally positioned flap + chemical root surface conditioning groups at 6 mo ($p < 0.001$) and 12 mo ($p < 0.001$). The mean gain in keratinized tissue in the coronally positioned flap + chemical root surface conditioning group was better than that in the coronally positioned flap group, but they did not differ statistically significantly at 6 mo ($p = 0.018$) or 12 mo ($p = 0.005$).

Probing depth

At 6 mo, the probing pocket depth mean values were 1.04 ± 0.38 mm (chi-square for heterogeneity: 212.014 ± 4 , $p < 0.05$) in the coronally positioned flap group, 1.09 ± 0.14 mm (chi-square for heterogeneity: 0.182 ± 2 , $p = 0.913$) in the coronally positioned flap + chemical root surface conditioning group, and 1.16 ± 0.15 mm (chi-square for heterogeneity: 11.444 ± 3 , $p < 0.05$) in the coronally-positioned flap + EMD group (Table 3). At 12 mo, the probing pocket depth mean values were 0.98 ± 0.02 mm (chi-square for heterogeneity: 2.613 ± 1 , $p = 0.106$) in the coronally positioned flap group, 1.4 ± 0.00 mm in the coronally positioned flap + chemical root surface conditioning group, and 1.41 ± 0.33 mm (chi-square for heterogeneity: 24.739 ± 4 , $p < 0.05$) in the coronally-positioned flap + EMD group (Table 4). The probing pocket depths of the three groups were all < 2 mm at 6 and 12 mo.

Gingival recession depth

The gingival recession depth in the coronally positioned flap + EMD

Table 1. Characteristics of the included studies for the coronally positioned flap plus enamel matrix derivative (CPF + EMD) group

| Reference | Year | Study description | Defects | Subjects | Interventions | Cleaning of root surface | Duration | Outcomes |
|----------------|------|---|--|---|-------------------------------------|---|---------------------------------|---|
| Abbas (32) | 2003 | Clinical trial | Miller class I recession of at least 4 mm in depth | Six patients with six sites | CPF + EMD | A polishing paste and rubber cup, 24% ethylenediaminetetraacetic acid (EDTA) gel, pH 6.7, for 2 min | 12 mo | Mucogingival surgery in combination with the application of EMD resulted in predictable root coverage and gain of clinical attachment while maintaining shallow pockets |
| Berlucchi (15) | 2002 | Random assignment | Gingival recession depth > 2 mm Miller class I or II | 14 patients with 26 sites | CPF + CTG + EMD; CPF + EMD | EDTA 24% for 2 min | 6 mo | EMD for the treatment of Miller class I or II gingival recessions displayed good clinical results in combination with CPF or CPF + CTG |
| Berlucchi (16) | 2005 | Clinical trial | Gingival recession depth \geq 2 mm; Miller class I or II | 30 patients with 30 sites | CPF + EMD | Scaled and planed by ultrasonic instruments and Gracey curets; EDTA gel 24% for 2 min | 12 mo | CPF + EMD gave excellent results when treating Miller class I or II recession, with 91.7% root coverage obtained at 12 mo and a mean attachment gain of 3.23 mm |
| Hägwald (33) | 2002 | A blinded, split-mouth, placebo-controlled, randomized design | \geq 3-mm buccal recession defects Miller class I or II | 24 patients with 2 paired defects | CPF + vehicle solution CPF + EMD | Hand instruments and a rubber cup; conditioning with 24% EDTA gel (PrefgelA, BIORA) for 2 min | 1 and 3 wk, and 3, 6, and 12 mo | There was no clear benefit of combining Emdogain with this surgical technique |
| McGuire (34) | 2003 | Randomized, single-center, split-mouth protocol | Miller class II in incisors or bicuspids | 17 patients 34 sites | CPF + CTG CPF + EMD | Conditioning with 24% EDTA | 12 mo | The addition of EMD to the coronally advanced flap resulted in root coverage similar to the CTG |
| Modica (19) | 2000 | Randomized, controlled, split-mouth protocol | Miller class I or II | 12 patients 24 sites | CPF; CPF + EMD | Conditioning with 24% EDTA | 12 mo | EMD did not seem to improve significantly the clinical outcomes of gingival recession treated by means of CPF |
| Nemcovsky (17) | 2004 | Randomized, controlled, split-mouth protocol | Miller class I or II in anterior or premolar teeth | 70 patients 30 sites with EMD 40 sites with CTG | CGF; CPF + EMD | Conditioning with 24% EDTA | 6 and 12 mo | The CTG procedure was superior to the coronally positioned flap with the addition of EMD in percentage of coverage and increase in width of keratinized tissue |

Table 2. Characteristics of the 11 included studies for coronally positioned flap (CPF) and coronally positioned flap plus chemical root surface conditioning (CPF + RC) groups

| Reference | Year | Study description | Defects | Subjects | Interventions | Cleaning of root surface | Duration | Outcomes |
|---------------|------|---|--|-------------------------------------|--------------------------------------|---|---------------------------------|--|
| Amarante (35) | 2000 | A blinded, split-mouth, randomized design | ≥ 3-mm buccal recession; Miller class I or II in cuspids or bicuspid | 20 patients 40 sites | CPF; CPF + membrane | Polished with a rubber cup and an abrasive paste | 3 and 6 mo | CPF offered a predictable, simple and convenient approach as a root coverage procedure in Miller class I and II recession defects Combining CPF with a bioabsorbable membrane did not seem to improve the results |
| Córtes (36) | 2004 | A blinded, split-mouth, randomized design | Miller class I gingival recession (≥ 3.0 mm) | 13 patients 26 sites | CPF; CPF + ADM | Curets and washed with a saline solution | 6 mo | Both techniques provided significant root coverage in class I gingival recessions; however, a greater keratinized tissue thickness could be expected with ADM |
| Gurgan (37) | 2004 | Clinical trial | ≥ 1-mm buccal recession defects; Miller class I | 24 patients 24 sites | CPF | None | 1, 6, 12, and 60 mo | After the CPF procedure, the MGJ was observed to move back to its original position in the 60-mo follow-up |
| Hägewald (33) | 2002 | A blinded, split-mouth, placebo-controlled, randomized design | ≥ 3-mm buccal recession defects Miller class I or II | 24 patients with two paired defects | CPF + vehicle solution; CPF + EMD | Hand instruments and a rubber cup; conditioning with 24% EDTA gel (PrefgelA, BIORA) for 2 min | 1 and 3 wk, and 3, 6, and 12 mo | There was no clear benefit to combine EMD with this surgical technique |
| Leknes (38) | 2005 | Randomized, split-mouth protocol | Miller class I or II in cuspids or bicuspid | 20 patients 40 sites | CPF; CPF + biodegradable membrane | Polished with a rubber cup and an abrasive paste | 6 and 12 mo, and 6 years | The CPF procedure offered a simple and reliable treatment alternative as a root coverage procedure. Placement of a biodegradable membrane underneath the flap did not improve either the short- or long-term results |
| Lims (39) | 2003 | Randomized, split-mouth protocol | Miller class I or II in cuspids or bicuspid | 10 patients 20 sites | CPF; CPF + tr-ePTFE | Scaled and planed; conditioned with 50 mg/ml tetracycline for 3 min | 6 mo | The amount of root coverage obtained with CPF was greater than that observed with GTR |

Table 2. Continued

| Reference | Year | Study description | Defects | Subjects | Interventions | Cleaning of root surface | Duration | Outcomes |
|-----------------|------|---|---|---|--------------------------|---|----------|--|
| Modica (19) | 2000 | Randomized, controlled, split-mouth protocol | Miller class I or II | 12 patients 24 sites | CPF; CPF + EMD | None | 12 mo | EMD did not seem to improve significantly the clinical outcomes of gingival recession treated by means of CPF |
| Pini Prato (40) | 2005 | Clinical trial | ≥ 2-mm buccal recession defects Miller class I | 60 patients 60 sites | CPF | None | 6 mo | The location of the gingival margin relative to the cemento-enamel junction following CPF procedure seemed to affect CRC |
| Silva (41) | 2004 | Randomized, controlled, split-mouth protocol | Miller class I in anterior or premolar teeth | 11 patients 22 sites | CPF; CPF + SCTG | None | 6 mo | Both surgical approaches were effective in addressing root coverage, but CPF + SCTG was better in increasing KT |
| Thromelli (42) | 1996 | Randomized, controlled, split-mouth protocol | Class I or II recession defects | 11 patients 22 sites | CPF CPF + fibrin glue | Tetracycline in sterile water (100 mg/ml) for about 4 min | 6 mo | There were no clinical or statistically significant differences between the treatments |
| Woodyard (43) | 2004 | Randomized, controlled, blinded, clinical investigation | Miller class I or II buccal recession defects of ≥ 3 mm treated | 24 subjects; 12 for CPF and 12 for CPF + ADM | CPF; CPF + ADM | Tetracycline in sterile water (100 mg/ml) for about 4 min | 6 mo | Treatment with a CPF plus an ADM allograft significantly increased gingival thickness when compared with a CPF allograft alone |

ADM, acellular dermal matrix allograft; EMD, enamel matrix derivative; KT, keratinized tissue; MGJ, mucogingival junction; tr-ePTFE, titanium-reinforced expanded polytetrafluoroethylene; SCTG, subepithelial connective tissue graft.

Table 3. Mean (\pm SD) of clinical attachment level (CAL), keratin thickness (KT), probing pocket depth (PD) and gingival recession depth (GRD) for 6 mo of follow-up

| | | 6-mo follow-up | | | | |
|-----|-----------|----------------|----------|--------------------------------------|-------------------------|-------------------------------|
| | | Study <i>n</i> | Subjects | Baseline weighted mean \pm SD (mm) | 6 mo mean \pm SD (mm) | Difference mean \pm SD (mm) |
| CAL | CPF | 7 | 147 | 6.67 \pm 3.69 | 4.25 \pm 4.23 | 2.42 \pm 0.70 |
| | CPF + RC | 3 | 33 | 4.63 \pm 0.24 | 2.41 \pm 0.12 | 2.22 \pm 0.36 |
| | CPF + EMD | 3 | 42 | 5.26 \pm 0.76 | 1.25 \pm 0.03 | 4.01 \pm 0.77 |
| KT | CPF | 7 | 147 | 2.68 \pm 0.47 | 2.64 \pm 0.49 | -0.04 \pm 0.37 |
| | CPF + RC | 3 | 33 | 2.47 \pm 0.72 | 2.57 \pm 0.78 | 0.14 \pm 0.47 |
| | CPF + EMD | 3 | 55 | 1.43 \pm 0.37 | 2.02 \pm 0.34 | 0.59 \pm 0.20 |
| PD | CPF | 7 | 147 | 1.19 \pm 0.14 | 1.04 \pm 0.38 | 0.15 \pm 0.30 |
| | CPF + RC | 3 | 33 | 1.30 \pm 0.29 | 1.09 \pm 0.14 | 0.22 \pm 0.22 |
| | CPF + EMD | 4 | 72 | 1.49 \pm 0.26 | 1.16 \pm 0.15 | 0.33 \pm 0.17 |
| GRD | CPF | 7 | 147 | 3.36 \pm 0.36 | 0.80 \pm 0.42 | 2.56 \pm 0.43 |
| | CPF + RC | 4 | 69 | 3.62 \pm 0.29 | 1.17 \pm 0.25 | 2.46 \pm 0.48 |
| | CPF + EMD | 5 | 108 | 3.91 \pm 0.35 | 0.62 \pm 0.36 | 3.29 \pm 0.43 |

SD, between studies; *, $p < 0.001$.
 CPF, coronally positioned flap; CPF + EMD, coronally positioned flap plus enamel matrix derivative; CPF + RC, coronally positioned flap plus chemical root surface conditioning.

Table 4. Mean (\pm SD) of clinical attachment level (CAL), keratin thickness (KT), probing pocket depth (PD) and gingival recession depth (GRD) for 12 mo of follow-up

| | | 12-mo follow-up | | | | |
|-----|-----------|-----------------|----------|--------------------------------------|--------------------------|-------------------------------|
| | | Study <i>n</i> | Subjects | Baseline weighted mean \pm SD (mm) | 12 mo mean \pm SD (mm) | Difference mean \pm SD (mm) |
| CAL | CPF | 2 | 31 | 9.54 \pm 4.33 | 7.85 \pm 4.18 | 1.69 \pm 0.15 |
| | CPF + RC | 1 | 36 | 5.50 \pm 0.00 | 2.40 \pm 0.00 | 3.10 \pm 0.00 |
| | CPF + EMD | 4 | 89 | 5.37 \pm 0.72 | 1.76 \pm 0.44 | 3.61 \pm 0.50 |
| KT | CPF | 2 | 31 | 2.87 \pm 0.37 | 2.97 \pm 0.04 | 0.10 \pm 0.41 |
| | CPF + RC | 1 | 36 | 2.40 \pm 0.00 | 2.70 \pm 0.00 | 0.30 \pm 0.00 |
| | CPF + EMD | 4 | 102 | 1.86 \pm 0.50 | 2.47 \pm 0.44 | 0.61 \pm 0.14 |
| PD | CPF | 2 | 31 | 1.30 \pm 0.00 | 0.98 \pm 0.02 | 0.31 \pm 0.02 |
| | CPF + RC | 1 | 36 | 1.60 \pm 0.00 | 1.40 \pm 0.00 | 0.20 \pm 0.00 |
| | CPF + EMD | 5 | 119 | 1.57 \pm 0.29 | 1.41 \pm 0.33 | 0.16 \pm 0.29 |
| GRD | CPF | 2 | 31 | 3.18 \pm 0.57 | 1.37 \pm 0.04 | 1.82 \pm 0.53 |
| | CPF + RC | 1 | 36 | 3.90 \pm 0.00 | 1.00 \pm 0.00 | 2.90 \pm 0.00 |
| | CPF + EMD | 5 | 119 | 3.91 \pm 0.42 | 0.72 \pm 0.40 | 3.16 \pm 0.41 |

SD, between studies; *, $p < 0.001$.
 CPF, coronally positioned flap; CPF + EMD, coronally positioned flap plus enamel matrix derivative; CPF + RC, coronally positioned flap plus chemical root surface conditioning.

group decreased from 3.91 \pm 0.35 to 0.62 \pm 0.36 mm (chi-square for heterogeneity: 21.537 \pm 4, $p < 0.05$) at 6 mo (Table 3) and decreased from 3.91 \pm 0.42 to 0.72 \pm 0.40 mm (chi-square for heterogeneity: 24.739 \pm 4, $p < 0.05$) at 12 mo (Table 4). In the coronally positioned flap group, the gingival recession depth decreased from 3.36 \pm 0.36 to 0.80 \pm 0.42 mm (chi-square for heterogeneity: 76.469 \pm 5, $p < 0.05$) at 6 mo (Table 3) and decreased from 3.18 \pm

0.57 to 1.37 \pm 0.04 mm (chi-square for heterogeneity: 2.613 \pm 1, $p = 0.106$) at 12 mo (Table 4). In the coronally positioned flap + chemical root surface conditioning group, gingival recession depth decreased from 3.62 \pm 0.29 to 1.17 \pm 0.25 mm (chi-square for heterogeneity: 27.290 \pm 3, $p < 0.05$) at 6 mo (Table 3) and decreased from 3.90 \pm 0.00 to 1.00 \pm 0.00 mm at 12 mo (Table 4). The change in the coronally positioned flap + EMD group was statistically significantly

better ($p < 0.001$) than those in the coronally positioned flap and coronally positioned flap + chemical root surface conditioning groups at 6 and 12 mo. At 6 mo, the mean reduction in the coronally positioned flap group was better than that in the coronally positioned flap + chemical root surface conditioning group, but this was not statistically significant ($p = 0.112$). At 12 mo, the change in the coronally positioned flap + chemical root surface conditioning group was statisti-

cally significantly better ($p < 0.001$) than that in the coronally positioned flap group.

Root coverage percentage

The results in the coronally positioned flap group were $74.12 \pm 15.80\%$ (chi-square for heterogeneity: 88.952 ± 6 , $p < 0.05$) root coverage percentage at 6 mo and $54.16 \pm 0.00\%$ root coverage percentage at 12 mo. In the coronally positioned flap + chemical root surface conditioning group, the results were $60.88 \pm 5.12\%$ (chi-square for heterogeneity: 0.182 ± 2 , $p = 913$) root coverage percentage at 6 mo and $79.00 \pm 0.00\%$ root coverage percentage at 12 mo. In the coronally positioned flap + EMD group, the results were $84.33 \pm 7.72\%$ (chi-square for heterogeneity 21.537 ± 4 , $p < 0.05$) root coverage percentage at 6 mo and $84.42 \pm 8.75\%$ (chi-square for heterogeneity 25.000 ± 5 , $p < 0.05$) root coverage percentage at 12 mo (Table 5). The root coverage percentage of the coronally positioned flap + EMD group was statistically significantly better ($p < 0.001$) than those of the coronally positioned flap and coronally positioned flap + chemical root surface conditioning groups at 6 and 12 mo. At 6 mo, the root coverage percentage of the coronally positioned flap group was statistically significantly better ($p < 0.001$) than that of the coronally positioned flap + chemical root surface conditioning group. At 12 mo, the root coverage percentage of the coronally positioned flap + chemical root surface conditioning group was statistically significantly better ($p < 0.001$) than that of the coronally positioned flap group.

Discussion

Clinically, the present analysis demonstrated that all three groups are useful in treating Miller's class I and II recession defects. All three groups achieved considerable root coverage and gains in clinical attachment, and maintained the amount of keratinized tissue and shallow probing pocket depths. The application of EMD to denuded root surfaces treated with the coronally positioned flap procedure significantly increased the percentage of root coverage and the attachment level compared with coronally positioned flap alone and the coronally positioned flap + chemical root surface conditioning procedure. In the present study, the coronally positioned flap and coronally positioned flap + chemical root surface conditioning groups resulted in root coverage percentage values ranging from 55 to 75%. The mean root coverage percentage of coronally positioned flap + EMD-treated sites ranged from 71.7 to 95.1%. This implies that coronally positioned flap alone for root coverage may be technique-sensitive, and the success of root coverage is overwhelmingly influenced by the condition of the surgical sites, such as the soft-tissue thickness (15,44). The coronally positioned flap procedure, combined with EMD for root coverage, showed improved and more consistent results.

The average root coverage of coronally positioned flap plus EMD amounted to $84.33 \pm 7.72\%$ after 6 mo and $84.42 \pm 8.75\%$ at 12 mo. The outcome of coronally positioned flap + EMD was better than coronally positioned flap alone after 6 mo ($74.12 \pm 15.80\%$) and 12 mo

($54.16 \pm 0.00\%$); the outcomes were also better than coronally positioned flap + chemical root surface conditioning after 6 mo ($60.88 \pm 5.12\%$) and 12 mo ($79.00 \pm 0.00\%$). The amount of root coverage obtained was quite stable between 6 and 12 mo in the coronally positioned flap + EMD group for root coverage. This suggests that root coverage procedures in the coronally positioned flap alone and coronally positioned flap + chemical root surface conditioning procedures were unpredictable. They became more predictable when the coronally positioned flap procedure was improved by the modification of adding EMD. In the present study, we did not discuss the relationship between the thickness of the flap and the amount of root coverage after coronally positioned flap alone or coronally positioned flap in combination with chemical root surface conditioning or EMD. Neither this nor the other studies detected any significant impact of pre-operative clinical parameters (tissue thickness, recession width, papilla height and width, vestibular bone height, anatomic factors, smoking, or others) on root coverage procedures. More evidence is required to determine which pre-operative clinical parameters can influence the recession reduction following coronally positioned flap surgery.

Root surface conditioning is a prerequisite of the EMD protocol on the premise that root surface conditioning removes the smear layer and allows the EMD to precipitate onto a surface free of organic remnants. The use of EDTA seems to improve the quality as well as quantity of the available root surface before EMD use, by removing the smear layer and exposing the collagen fibers (58). However, three quasi-

Table 5. Mean (\pm SD) of root coverage percentage for 6 and 12 mo of follow-up

| | 6-mo follow-up | | | 12-mo follow-up | | |
|-----------|----------------|----------|-------------------|-----------------|----------|------------------|
| | Study <i>n</i> | Subjects | RCP (%) | Study <i>n</i> | Subjects | RCP (%) |
| CPF | 7 | 147 | 74.12 ± 15.80 | 1 | 11 | 54.16 ± 0.00 |
| CPF + RC | 3 | 33 | 60.88 ± 5.12 | 1 | 36 | 79.00 ± 0.00 |
| CPF + EMD | 3 | 55 | 84.33 ± 7.72 | 6 | 149 | 84.42 ± 8.75 |

SD, between studies; *, $p < 0.001$.

CPF, coronally positioned flap; CPF + EMD, coronally positioned flap plus enamel matrix derivative; CPF + RC, coronally positioned flap plus chemical root surface conditioning; RCP, root coverage rate percentage.

experimental studies examined the effects of a solution containing 24% EDTA at a pH of between 7.0 and 7.2. The EDTA was applied to root surfaces for either 2 or 3 min, and the study duration was 6 mo. In all three studies, there were no differences in probing pocket depth, clinical attachment level, gingival recession depth, or probing bone levels between EDTA treatment and control root surfaces (59–61). In Mariotti's systematic review, the use and application of citric acid, tetracycline, or EDTA to modify the root surface provided no clinical benefit to the patient with respect to reduction of probing depths or gain in clinical attachment levels (62). It was confirmed in the present study that clinical outcomes for root coverage do not depend on the use of root conditioning. There were no statistically significant differences between coronally positioned flap alone group and the coronally positioned flap + chemical root surface conditioning group in the amount of root coverage and increase of clinical attachment level at 6 mo. Obviously, chemical root surface conditioning used alone therefore cannot be considered as beneficial for root coverage (63). Although the root coverage and increase of clinical attachment level in the coronally positioned flap + chemical root surface conditioning group was statistically significantly better than that of the coronally positioned flap group at 12 mo, it was justified by only one study that was included in our study. It needs more evidence to verify the above point.

The postoperative measurement period of 6 mo is sufficient to determine the long-term results of the method studied because it has been shown that a stable tissue relationship exists after the first postoperative month following root coverage procedures, and that was the reason why we decided to examine data at 6 and 12 mo on a short-term basis (64). In order to provide a definitive comparison of the results of coronally positioned flap alone and coronally positioned flap + chemical root surface conditioning procedures for root coverage, changes in clinical results between these two procedures were

distinguished in this analysis. There was no significant difference at the 6-mo comparison time-point among clinical attachment level, keratinized tissue, probing pocket depth and gingival recession depth, except for root coverage percentage. However, there were certain contrariwise differences between coronally positioned flap and coronally positioned flap + chemical root surface conditioning in terms of clinical attachment level, probing pocket depth, gingival recession depth and root coverage percentage at the 12-mo follow-up. We cannot draw any firm conclusions about the respective data owing to the limited number of studies included in the 12-mo follow-up. More evidence is required to determine definitively whether root conditioning improves the results of root coverage for the coronally positioned flap procedure.

It is still not clear which biological mechanism is involved in the regeneration of periodontal tissue during healing in root coverage. Finally, histological reports are needed to gain greater insights into the tissue-healing process and prove that EMD is really responsible for improving the percentage of regenerated vs. repaired tissues compared with other techniques.

From this systemic analysis, it can be concluded that root coverage manipulated by the coronally positioned flap and coronally positioned flap + chemical root surface conditioning procedures were unpredictable. The results can be modified by adding EMD to increase significantly the clinical outcomes of gingival recession.

References

1. The American Academy of Periodontology. *Glossary of periodontal terms*, 4th edn. Chicago, IL, USA: The American Academy of Periodontology, 2001:44.
2. Wennstrom JL, Zucchelli G. Increased gingival dimensions. A significant factor for successful outcome of root coverage procedures? A 2-year prospective clinical study. *J Clin Periodontol* 1996;**23**:770–777.
3. Harris RJ, Harris AW. The coronally positioned pedicle graft with inlaid margins: a predictable method of obtaining root coverage of shallow defects.

4. Allen EP, Miller PD. Coronal positioning of existing gingiva: short term results in the treatment of shallow marginal tissue recession. *J Periodontol* 1989;**60**:316–319.
5. Becker W, Becker BE, Berg L. New attachment after treatment with root isolation procedures: report for treated class III and class II furcations and vertical osseous defects. *Int J Periodont Restor Dent* 1988;**8**:9–23.
6. Hammarström L, Heijl L. Periodontal regeneration in a buccal dehiscence model in monkeys after application of enamel matrix derivative. *J Clin Periodontol* 1997;**24**:669–677.
7. Heijl L. Periodontal regeneration with enamel matrix derivative in one human experimental defect. A case report. *J Clin Periodontol* 1997;**24**:693–696.
8. Sculean A, Reich E. Treatment of intrabony periodontal defects with an enamel matrix protein derivative (Emdogain): a report on 32 cases. *Int J Periodont Restor Dent* 1999;**19**:157–164.
9. Heden G. A case report study of 72 consecutive Emdogain-treated intrabony periodontal defects: clinical and radiographic findings after 1 year. *Int J Periodont Restor Dent* 2000;**21**:127–139.
10. Hammarström L. Enamel matrix, cementum development and regeneration. *J Clin Periodontol* 1997;**24**:658–668.
11. Gestrelus S, Andersson C, Johansson AC *et al.* Formulation of enamel matrix derivative for surface coating. Kinetics and cell colonization. *J Clin Periodontol* 1997;**24**:678–684.
12. Yukna RA, Mellonig JT. Histologic evaluation of periodontal healing in humans following regenerative therapy with enamel matrix derivative. A 10-case series. *J Periodontol* 2000;**71**:752–759.
13. Windisch P, Sculean A, Klein F *et al.* Comparison of clinical, radiographic, and histometric measurements following treatment with guided tissue regeneration or enamel matrix proteins in human periodontal defects. *J Periodontol* 2002;**73**:409–417.
14. Cochran DL, King Schoolfield J, Velasquez-Plata D, Mellonig JT, Jones A. The effect of enamel matrix proteins on periodontal regeneration as determined by histological analyses. *J Periodontol* 2003;**74**:1043–1055.
15. Berlucchi I, Francetti L, Del Fabbro M, Testori T, Weinstein RL. Enamel matrix proteins (Emdogain) in combination with coronally advanced flap or subepithelial connective tissue graft in the treatment of shallow gingival recessions. *Int J Periodont Restor Dent* 2002;**22**:583–593.
16. Berlucchi I, Francetti L, Del Fabbro M, Basso M, Weinstein RL. The influence of

- anatomical features on the outcome of gingival recessions treated with coronally advanced flap and enamel matrix derivative: a 1-year prospective study. *J Periodontol* 2005;**76**:899–907.
17. Nemcovsky CE, Artzi Z, Tal H, Kozlovsky A, Moses O. A multicenter comparative study of two root coverage procedures: coronally advanced flap with addition of enamel matrix proteins and subpedicle connective tissue graft. *J Periodontol* 2004;**75**:600–607.
 18. Cueva MA, Boltchi FE, Hallmon WW, Nunn ME, Rivera-Hidalgo F, Rees T. A comparative study of coronally advanced flaps with and without the addition of enamel matrix derivative in the treatment of marginal tissue recession. *J Periodontol* 2004;**75**:949–956.
 19. Modica F, Pizzo MD, Rocuzzo M, Romagnolo R. Coronally advanced flap for the treatment of buccal gingival recessions with and without enamel matrix derivative. A split-mouth study. *J Periodontol* 2000;**71**:1693–1698.
 20. Hanes PJ, O'Brien NJ, Garnick JS. A morphological comparison of radicular dentin following root planning and treatment with citric acid or tetracycline HCL. *J Clin Periodontol* 1991;**18**:660–668.
 21. Hanes RJ, Poison AM, Frederick GT. Initial wound healing attachments to demineralized dentin. *J Periodontol* 1988;**59**:176–183.
 22. Hanes PJ, Poison AM, Frederick GR. Citric acid treatment of periodontitis affected cementum. A scanning electron microscope study. *J Clin Periodontol* 1991;**18**:567–575.
 23. Hanes PJ, Poison AM, Ladenheim S. Cell and fiber attachment to demineralized dentin front normal root surfaces. *J Periodontol* 1985;**56**:752–765.
 24. Register AA, Burdick FA. Accelerated reattachment with cementogenesis to dentin demineralized in situ. 1. Optimum range. *J Periodontol* 1975;**46**:646–655.
 25. Lasho DJ, O'Leary TJ, Kafrawy AH. A scanning electron microscope study of the effects of various agents on instrumented periodontally involved root surfaces. *J Periodontol* 1983;**54**:210–220.
 26. Labahn R, Fahrenbach WH, Clark SM, Lie T, Adams DF. Root dentin morphology after different modes of citric acid and tetracycline hydrochloride conditioning. *J Periodontol* 1992;**63**:303–309.
 27. Garrett JS, Crigger M, Egelberg J. Effects of citric acid on diseased root surfaces. *J Periodont Res* 1978;**13**:155–163.
 28. Willey R, Steinberg AD. Scanning electron microscopic studies of root dentin surfaces treated with citric acid, elastase, hyaluronidase, pronase and collagenase. *J Periodontol* 1984;**55**:592–596.
 29. Lowenguth RA, Blieden TM. Periodontal regeneration: root surface demineralisation. *Periodontol* 2000 1993;**1**:54–68.
 30. Egelberg J. *Periodontics, the scientific way*, 2nd edn. Malmö, Sweden: OdontoScience, 1995;191–226.
 31. McGuire MK, Cochran DL. Evaluation of human recession defects treated with coronally advanced flaps and either enamel matrix derivative or connective tissue. Part 2: Histological evaluation. *J Periodontol* 2003;**74**:1126–1135.
 32. Abbas F, Wennström J, Van der Weijden V, Schneiders T, Van der Velden U. Surgical treatment of gingival recessions using emdogain gel: clinical procedure and case reports. *Int J Periodont Restor Dent* 2003;**23**:607–613.
 33. Hägewald S, Spahr A, Rompola E, Haller B, Heijl L, Bernimoulin JP. Comparative study of Emdogain and coronally advanced flap technique in the treatment of human gingival recessions. A prospective controlled clinical study. *J Clin Periodontol* 2002;**29**:35–41.
 34. McGuire MK, Nunn M. Evaluation of human recession defects treated with coronally advanced flaps and either enamel matrix derivative or connective tissue. Part 1: Comparison of clinical parameters. *J Periodontol* 2003;**74**:1110–1125.
 35. Amarante ES, Leknes KN, Skavland J, Lie T. Coronally positioned flap procedures with or without a bioabsorbable membrane in the treatment of human gingival recession. *J Periodontol* 2000;**71**:989–998.
 36. Côrtes AQ, Martins ã, Francisco H et al. Coronally positioned flap with or without acellular dermal matrix graft in the treatment of class I gingival recessions: a randomized controlled clinical study. *J Periodontol* 2004;**75**:1137–1144.
 37. Gurgan CA, Oruc AM, Akkaya M. Alterations in location of the mucogingival junction 5 years after coronally repositioned flap surgery. *J Periodontol* 2004;**75**:893–901.
 38. Leknes KN, Amarante ES, Price DE, Bøe OE, Skavland RJ, Lie T. Coronally positioned flap procedures with or without a biodegradable membrane in the treatment of human gingival recession. A 6-year follow-up study. *J Clin Periodontol* 2005;**32**:518–529.
 39. Lins LH, Lima AF, Sallum AW. Root coverage: comparison of coronally positioned flap with and without titanium-reinforced barrier membrane. *J Periodontol* 2003;**74**:168–174.
 40. Pini Prato GP, Baldi C, Nieri M et al. Coronally advanced flap: the post-surgical position of the gingival margin is an important factor for achieving complete root coverage. *J Periodontol* 2005;**76**:713–722.
 41. da Silva RC, Joly JC, de Lima AF, Tatakis DN. Root coverage using the coronally positioned flap with or without a subepithelial connective tissue graft. *J Periodontol* 2004;**75**:413–419.
 42. Trombelli L, Scabbia A, Wikesjö UM, Calura G. Fibrin glue application in conjunction with tetracycline root conditioning and coronally positioned flap procedure in the treatment of human gingival recession defects. *J Clin Periodontol* 1996;**23**:861–867.
 43. Woodyard JG, Greenwell H, Hill M, Drisko C, Iasella JM, Scheetz J. The clinical effect of acellular dermal matrix on gingival thickness and root coverage compared to coronally positioned flap alone. *J Periodontol* 2004;**75**:44–56.
 44. Baldi C, Pini-Prato G, Pagliaro U et al. Coronally advanced flap procedure for root coverage. Is flap thickness a relevant predictor to achieve root coverage? A 19-case series. *J Periodontol* 1999;**70**:1077–1084.
 45. Pini-Prato G, Baldi C, Pagliaro U et al. Coronally advanced flap procedure for root coverage. Treatment of root surface: root planning versus polishing. *J Periodontol* 1999;**70**:1064–1076.
 46. Clauser C, Nieri M, Franceschi D, Pagliaro U, Pini-Prato G. Evidence-based mucogingival therapy. Part 2: Ordinary and individual patient data meta-analyses of surgical treatment of recession using complete root coverage as the outcome variable. *J Periodontol* 2003;**74**:741–756.
 47. Coatoam G. Laterally moved, coronally advanced flap: a modified surgical approach for isolated recession-type defects. *J Periodontol* 2004;**75**:1734–1741.
 48. Zucchelli G, Cesari C, Amore C, Montebugnoli L, De Sanctis M. Laterally moved, coronally advanced flap: a modified surgical approach for isolated recession-type defects. *J Periodontol* 2004;**75**:1734–1741.
 49. Harris RJ. A comparison of 2 root coverage techniques: guided tissue regeneration with a bioabsorbable matrix style membrane versus a connective tissue graft combined with a coronally positioned pedicle graft without vertical incisions. Results of a series of consecutive cases. *J Periodontol* 1998;**69**:1426–1434.
 50. Cafiero C, Coraggio F, Vaia E, de Paoli S. Guided tissue regeneration versus coronally repositioned flap in the treatment of recession with double papillae. *Int J Periodont Restor Dent* 1998;**18**:444–453.
 51. Pini Prato G, Pagliaro U, Baldi C et al. Coronally advanced flap procedure for root coverage. Flap with tension versus flap without tension: a randomized controlled clinical study. *J Periodontol* 2000;**71**:188–201.

52. Saletta D, Pini Prato G, Pagliaro U, Baldi C, Mauri M, Nieri M. Coronally advanced flap procedure: is the interdental papilla a prognostic factor for root coverage? *J Periodontol* 2001;**72**:760–766.
53. Thompson BK, Meyer R, Singh GB, Mitchell W. Desensitization of exposed root surfaces using a semilunar coronally positioned flap. *Gen Dent* 2000;**48**:68–71.
54. Trombelli L, Tatakis DN, Scabbia A, Zimmerman GJ. Comparison of mucogingival changes following treatment with coronally positioned flap and guided tissue regeneration procedures. *Int J Periodont Restor Dent* 1997;**17**:448–455.
55. Harris RJ, Harris AW. The coronally positioned pedicle graft with inlaid margins: a predictable method of obtaining root coverage of shallow defects. *Int J Periodont Restor Dent* 1994;**14**:228–241.
56. Tarnow D. Solving restorative esthetic dilemmas with the semilunar coronally positioned flap. *J Esthet Dent* 1994;**6**:61–64.
57. Miller PD Jr, Binkley LH Jr. Root coverage and ridge augmentation in Class IV recession using a coronally positioned free gingival graft. *J Periodontol* 1986;**57**:360–363.
58. Blomlof J, Blomlof L, Lindskog S. Effect of different concentrations of EDTA on smear removal and collagen exposure in periodontitis-affected root surfaces. *J Clin Periodontol* 1997;**24**:534–537.
59. Blomlöf L, Jonsson B, Blomlöf J, Lindskog S. A clinical study of root surface conditioning with an EDTA gel. II. Surgical periodontal treatment. *Int J Periodont Restor Dent* 2000;**20**:567–573.
60. Blomlöf L, Bergman E, Forsgårdh Å *et al.* A clinical study of root surface conditioning with an EDTA gel. I. Nonsurgical periodontal treatment. *Int J Periodont Restor Dent* 2000;**20**:561–565.
61. Mayfield L, Söderholm G, Norderyd O, Attström R. Root conditioning using EDTA gel as an adjunct to surgical therapy for the treatment of intraosseous periodontal defects. *J Clin Periodontol* 1998;**25**:707–714.
62. Mariotti A. Efficacy of chemical root surface modifiers in the treatment of periodontal disease. A systematic review. *Ann Periodontol* 2003;**8**:205–226.
63. Wennström JL. Mucogingival therapy. *Ann Periodontol* 1996;**1**:671–701.
64. Caffesse R, Guinard E. Treatment of localized recessions, part IV. Results after three years. *J Periodontol* 1980;**51**:167–170.