

# **Antimicrobial Activity of Antibiotics in Combination with Natural flavonoids against Clinical Extended-Spectrum B-Lactamase (ESBL)-producing *Klebsiella pneumoniae* .**

秦一平

**Rong-Dih Lin;Yi-Ping Chin;Mei-Hsien Lee**

摘要

## **Abstract**

Extended-spectrum beta-lactamases (ESBLs) are plasmid-mediated class A enzymes commonly found in the family Enterobacteriaceae, mainly in *Klebsiella pneumoniae*. Flavonoids have also been reported to possess antimicrobial activity. In this study, the in vitro activities of 18 antibiotics and 12 flavonoids against 20 ESBL-producing *K. pneumoniae* isolates were evaluated. All of these isolates were susceptible to imipenem and cefmetazole, but were resistant to ampicillin, ampicillin/sulbactam, aztreonam, cefazolin, cefoperazone, cefotaxime, ceftazidime, ceftriaxone, cefuroxime, piperacillin and ticarcillin. Susceptibilities to amikacin, amoxicillin/clavulanate, ceftazidime, ciprofloxacin and gentamicin were variable. Myricetin, a flavonol, inhibited ESBL-producing *K. pneumoniae* isolates at a high minimum inhibitory concentration (MIC) (MIC<sub>90</sub>) value 256 mg/mL), but exhibited significant synergic activity against ESBL-producing *K. pneumoniae* in separate combination with amoxicillin/clavulanate, ampicillin/sulbactam and ceftazidime. Because of the low-toxic nature of flavonoids, the combination of antibiotics and flavonoids is a potential new strategy for developing therapies for infections caused by ESBL-producing bacteria in the future.