

2. Engberding, R., Reinbach, R., Frommelt, T. The influence of nicardipine on left ventricular hemodynamics and compliance in patients with coronary heart disease. *J. Cardiovasc. Pharm.* 1990; **16**: 20-25.
3. Mielcarek, J. Inclusion complexes of nifedipine and other 1,4-dihydropyridine derivatives with cyclodextrins. IV. The UV study on photochemical stability of the inclusion complexes of nisoldipine, nimodipine, nitrendipine and nicardipine with β -cyclodextrin in the solution. *Acta Poloniae Pharm.* 1995; **52**: 459-463.
4. Islam, M.S., Narurkar, M.M. The effect of 2-hydroxypropyl- β -cyclodextrin on the solubility, stability and dissolution rate of famotidine. *Drug Dev. Ind. Pharm.* 1991; **17**: 1229-1239.
5. Bekers, O., Uijtendaal, E.V., Beijnen, J.H., Bult, A., Underberg, W.J.M. Cyclodextrins in the pharmaceutical field. *Drug Dev. Ind. Pharm.* 1991; **17**: 1503-1509.
6. Duchene, D., Wouessidjewe, D. Pharmaceutical uses of cyclodextrins and derivatives. *Drug Dev. Ind. Pharm.* 1990; **16**: 2487-2499.
7. Bekers, O., Beijnen, J.H., Vis, B.J., Suenaga, A., Otagiri, M., Bult, A., Underberg, W.J.M. Effect of cyclodextrin complexation on the chemical stability of doxorubicin and daunorubicin in aqueous solution. *Int. J. Pharm.* 1991; **72**: 123-130.
8. Choudhury, S., Nelson, K.F. Improvement of oral bioavailability of carbamazepine by inclusion in 2-hydroxypropyl- β -cyclodextrin. *Int. J. Pharm.* 1992; **85**: 175-180.
9. Yamamoto, M., Yoshida, A., Hirayama, F., Uekema, K. Some physicochemical properties of branched β -cyclodextrins and their inclusion characteristics. *Int. J. Pharm.* 1989; **49**: 163-171.
10. Chao, S.-H. Comparative photostability studies of *o*-nitrophenyl and *m*-nitrophenyl 1,4-dihydropyridine calcium channel blocker in alcoholic solvents. Master Thesis; Taipei Medical College, Taipei, Taiwan 1998.
11. Higuchi, T., Connors, K.A. Phase solubility techniques. *Anal. Chem. Instr.* 1965; **4**: 117-212.
12. Erden, N., Celebi, N. A study of the inclusion complex of naproxen with β -cyclodextrin. *Int. J. Pharm.* 1988; **48**: 83-89.
13. Hassan, M.A., Suleiman, M.S., Najib, N.M. Improvement of the in vitro dissolution characteristics of famotidine by inclusion in β -cyclodextrin. *Int. J. Pharm.* 1990; **58**: 19-24.




Fig. 4. Dissolution profile of intact NCDP and 3 physical mixtures (1:1, 1:2 and 1:3) in pH 7.5 phosphate buffer. The results are expressed as mean \pm SD.