

present in tea is a promising antioxidant which may contribute to the reduced incidence of coronary and stroke mortality by the increased intake of green or black tea.

ACKNOWLEDGEMENTS

This study was supported by a grant from the National Science Council of the R.O.C, no. NSC 89-2314-B-038-012.

REFERENCES

- Cross, C.E., Halliwell, B., Borish E.T., Dryor, W.A., Ames, B.N., Saul, R.L., McCord, J.M., Harman, D. Oxygen radicals and human disease. *Ann. Intern. Med.* 1987; **107**: 526-545.
- Ambrosio, G., Backer, L.C., Hutchins, G.M., Weisman, H.F., Weisfeldt, M.L. Reduction in experimental infarction size by recombinant human superoxide dismutase: insights into the pathophysiology of reperfusion injury. *Circulation*. 1986; **74**: 1424-1433.
- Jayakumari, N., Ambikakumari, V., Balakrishnan, K.G., Subramonia, Iyer K. Antioxidant status in relation to free radical production during stable and unstable angina syndromes. *Atherosclerosis* 1992; **94**: 183-190.
- Flaherty, J.T. Myocardial injury mediated by oxygen free radicals. *Am. J. Med.* 1991; **91**(suppl. 3C): S79-S85.
- Chan, P., Niu, C.S., Tomlinson, B., Hong, C.T., Chen, J.P., Hong, C.Y. Effect of trilinolein on superoxide dismutase activity and left ventricular pressure in isolated rat hearts subjected to hypoxia and normoxic perfusion. *Pharmacology*. 1997; **55**: 252-258.
- Goldhaber, J.I., Weiss, J.N. Oxygen free radicals and cardiac reperfusion abnormalities. *Hypertension*. 1992; **20**: 118-127.
- Das, D.K., Engelman, R.M., Rousou, J.A., Breyer, R.H., Otani, H., Lemeshow, S. Pathophysiology of superoxide radical as potential mediator of perfusion injury in pig heart. *Basic Res. Cardiol.* 1986; **81**: 155-156.
- Shlafer, M., Myers, C.L., Adkins, S. Mitochondrial hydrogen peroxide generation and activities of glutathione peroxidase and superoxide dismutase following global ischemia. *J. Mol. Cell. Cardiol.* 1987; **19**: 1195-1206.
- Bolli, R. Oxygen-derived free radicals and myocardial reperfusion injury. An overview. *Cardiovasc. Drug Ther.* 1991; **5**: 249-268.
- Renaud, S., de Lorgeril, M. Wine, alcohol, platelets, and the French paradox for coronary heart disease. *Lancet* 1992; **339**: 1523-1526.
- Hertog, M.G., Feskens, E.J., Hollman, P.C., Katan, M.B., Kromhout, D. Dietary antioxidant flavonoids and risk of coronary heart disease: Zutphen Elderly Study. *Lancet*. 1993; **342**, 1007-1011.
- Hong, C.Y., Lai, L.J., Yeh, S.F. Linoleate-rich triacylglycerol in *Panax pseudoginseng* improves erythrocyte deformability in vitro. *Planta. Med.* 1993; **59**: 323-326.
- Chan, P., Cheng, J.T., Tsao, C.W., Niu, C.S., Hong, C.Y. The *in vitro* antioxidant activity of lipid-related natural substances as measured by enhanced chemiluminescence. *Life Sci.* 1996; **59**: 2067-2073.
- Chan, P., Hong, C.Y., Cheng, J.T. Myocardial protective effect of medicinal plant antioxidant trilinolein isolated from *Panax pseudoginseng*. *Life Sci.* 1997; **61**: 1999-2006.
- Chan, P.H., Chen, S.F., Yu, A.C.H. Induction of intracellular superoxide radical formation by arachidonic acid and by polyunsaturated fatty acids in primary astrocytic cultures. *J. Neurol. Chem.* 1998; **50**: 1185-1193.
- Bradford, M.M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 1976; **72**: 248-254.
- Chomczynski, P., Sacchi, N. Single-step of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 1987; **162**: 156-159.
- Herrin, D.L., Schmidt, G.W. Rapid, reversible staining of Northern blots prior to hybridization. *Biotechnology* 1988; **6**: 196-198.
- Sambrook, J., Fritsch, E.F., Maniatis, T. "Molecular cloning-a laboratory manual", 2nd ed.; Cold Spring Harbor Laboratory Press: Cold Spring Harbor NY. 1989.
- Steinberg, D. A critical look at the evidence for the oxidation of LDL in atherogenesis. *Atherosclerosis*. 1997; **131**(suppl.), S5-S7.