

- pendent mitogen-activated protein kinase pathway regulated by oxidant stress. *J. Biol. Chem.* 1997;272: 17810-4.
16. Li YM, Mitsuhashi T, Wojciechowicz D, Shimizu N, Li J, Stitt A, He C, Banerjee D, Vlassara H. Molecular identity and cellular distribution of advanced glycation end product receptors: relationship of p60 to OST-48 and p90 to 80K-H membrane proteins. *Proc. Natl. Acad. Sci. USA.* 1996;93:11047-52.
 17. Lin CH, Lin YF, Chang MC, Wu CH, Ho YS, Lee HM. Advanced glycosylation end products induced inducible nitric oxide synthase expression in C6 glioma cells: Involvement of a p38 MAP kinase-dependent mechanism. *Life. Sci.* 2001;69:2503-15.
 18. Loske C, Neumann A, Cunningham AM, Nichol K, Schinzel R, Riederer P, and Munch G. Cytotoxicity of advanced glycation end products is mediated by oxidative stress. *J. Neural. Transm.* 1998;105:1005-15.
 19. Martinez J, Sanchez T, Moreno JJ. Regulation of prostaglandin E₂ production by the superoxide radical and nitric oxide in mouse peritoneal macrophages. *Free Radic. Res.* 2000;32:303-11.
 20. Mohamed AK, Bierhaus A, Schiekofer S, Tritschler H, Ziegler R, Nawroth PP. The role of oxidative stress and NF-kappa B activation in late diabetic complications. *Biofactors.* 1999;10:157-67.
 21. Munch G, Schinzel R, Loske C, Wong A, Durancy N, Li JJ, Vlassara H, Smith MA, Perry G, and Riederer P. Alzheimer's disease-synergistic effects of glucose deficit, oxidative stress and advanced glycation end products. *J. Neural. Transm.* 1998;105:439-61.
 22. Rojas A, Caveda I, Romay C, Lopez E, Valdes S, Padron J, Glaria L, Martinez O, and Delgado R. Effects of advanced glycation end products on the induction of nitric oxide synthase in murine macrophages. *Biochem. Biophys. Res. Commun.* 1996;225:358-62.
 23. Satoh H, Togo M, Hara M, Miyata T, Han K, Maekawa H, Ohno M, Hashimoto Y, Kurokawa K, and Wanatabe T. Advanced glycation end products stimulate mitogen activated protein kinase and proliferation in rabbit vascular smooth muscle cells. *Biochem. Biophys. Res. Commun.* 1997;239:111-5.
 24. Schmidt AM, Weidman E, Lalla E, Yan SD, Hori O, Cao R, Brett JG, and Lamster IB. Advanced glycation end products (AGEs) induce oxidant stress in the gingiva: a potential mechanism underlying accelerated periodontal disease associated with diabetes. *J. Periodontal. Res.* 1996;31:508-15.
 25. Schmidt AM, Yan SD, Brett J, Mora R, Nowygrod R, and Stern D. Regulation of human mononuclear phagocyte migration by cell-surface-binding proteins for advanced glycation end products. *J. Clin. Invest.* 1993;91: 2155-68.
 26. Scivittaro V, Ganz MB, and Weiss MF. AGEs induce oxidative stress and activate protein kinase C-beta(II) in neonatal mesangial cells. *Am. J. Physiol.* 2000;278: F676- 83.
 27. Simm A, Munch G, Seif F, Schenk O, Heidland A, Richter H, Vamvakas S, and Schinzel R. Advanced glycation end products stimulate the MAP-kinase pathway in tubulus cell line LLC-PK1. *FEBS Lett.* 1997;410:481-4.
 28. Smedsrød B, Melkko J, Araki N, Sano H, Horiuchi S. Advanced glycation end products are eliminated by scavenger-receptor-mediated endocytosis in hepatic sinusoidal Kupffer and endothelial cells. *Biochem. J.* 1997;322 (Pt 2):567-73.
 29. Smith MA, Richey PL, Sayre LM, Beckman JS, and Perry G. Widespread peroxynitrite-mediated damage in Alzheimer disease. *J. Neurosci.* 1997;17:2653-7.
 30. Wautier JL, Zoukourian C, Chappéy O, Wautier MP, Guillausseau PJ, Cao R, Hori O, Stern D, and Schmidt AM. Receptor-mediated endothelial cell dysfunction in diabetic vasculopathy. Soluble receptor for advanced glycation end products blocks hyperpermeability in diabetic rats. *J. Clin. Invest.* 1996;97:238-43.
 31. Yamagishi S, Yamamoto Y, Harada S, Hsu CC, Yamamoto H. Advanced glycosylation end products stimulate the growth but inhibit the prostacyclin-producing ability of endothelial cells through interactions with their receptors. *FEBS Lett.* 1996;384:103-6.