

Physiological Levels of Glutamine Reduces ICAM-1 Expression in Endothelial Cell Activated by Preeclamptic Plasma

Chun-Sen Hsu, M.D., Szu-Yuan Chou, M.D., So-Jung Liang, M.D., Chun-Yaw Chang, M.D., Sung-Ling Yeh, Ph.D.

Department of Obstetrics and Gynecology, Taipei Medical University-Municipal Wan Fang Hospital, and Institute of Nutrition and Health Sciences, Taipei Medical University, Taipei, Taiwan

Running title: Glutamine effect on CAM expression in preeclampsia

This study was supported by research grant 93TMU-WFH-10 from the Taipei Medical University-Municipal Wan-Fang Hospital, Taipei, Taiwan.

Corresponding author:

Sung-Ling Yeh, PhD

School of Nutrition and Health Sciences

Taipei Medical University

250 Wu-Hsing Street, Taipei, Taiwan 110

Tel: 8862-27361661 ext. 6551-115

E-mail: sangling@tmu.edu.tw

Synopsis

Women with preeclampsia had lower glutamine levels than normal pregnant women.

Administering GLN at physiologic levels reduces endothelial cell ICAM-1 expression induced by preeclamptic plasma.

Abstract

OBJECTIVE: The purpose of our study was intended to investigate whether plasma glutamine (GLN) concentration would be depleted in pregnant women with preeclampsia, and whether administering GLN comparable to physiologic levels would decrease cellular adhesion molecule expression in human umbilical vein endothelial cells (HUVEC) induced by plasma from preeclamptic women.

STUDY DESIGN: We assessed plasma GLN levels from blood samples collected from 20 women with preeclampsia and 10 normal pregnant women. HUVEC were cultured in medium-199 containing fetal calf serum, antibiotics, growth factor and with different concentrations (0, 300, 500 uM) of GLN for 24 hr. We stimulated those cells for 1.5-6 hr with sera from patients with preeclampsia, and then determined the expression of intercellular cell adhesion molecule (ICAM)-1 and vascular cell adhesion molecule (VCAM)-1 on endothelial cells by flow cytometry.

RESULTS: Women with preeclampsia had significantly lower plasma GLN concentrations as compared with the normal pregnant women. We found that there were no differences in VCAM-1 expression in HUVEC among various GLN concentrations at each time point. However, ICAM-1 expression in HUVEC were significantly lower in 500 uM GLN group than 0 and 300uM groups at 3, 4.5, and 6hr.

CONCLUSION: This study showed that plasma from women with preeclampsia had significantly lower GLN levels than those from normal pregnant women, and that administering GLN at physiologic levels reduces HUVEC ICAM-1 expression induced by preeclamptic plasma.

Keywords: glutamine; preeclampsia; human umbilical vein endothelial cells; adhesion molecules.

Introduction

Preeclampsia is a pregnancy-specific syndrome in a previously healthy women characterized by increased blood pressure, proteinuria and/or pathologic edema appearing after 20 weeks of gestation.¹ The mechanisms leading to this syndrome have not been clarified. Recent studies suggest that endothelial activation/dysfunction is a central pathogenic feature in women with preeclampsia.^{2,3} Previous reports revealed that circulating levels of intracellular adhesion molecules (ICAM)-1 and vascular cell adhesion molecule (VCAM)-1 are increased in women with preeclampsia.⁴⁻⁶ Soluble CAM levels were associated with the severity of the disease, and have been thought to be markers of endothelial activation in this disease.⁴⁻⁶ ICAM-1 and VCAM-1 which belong to adhesion molecules of immunoglobulin gene superfamily are expressed by activated endothelium, and are important in the adhesion of monocytes, lymphocytes and neutrophils to activated endothelium.^{7,8} Previous reports showed that ICAM-1 and VCAM-1 expressions in human umbilical endothelial cells (HUVEC) was induced by plasma from preeclamptic patients.^{6,9}

Glutamine (GLN) is the most abundant free amino acid in plasma and tissue pool. It is (1) substrate for protein synthesis and precursors for nucleic acid biosynthesis, (2) an important carrier of nitrogen and carbon, (3) a precursor for gluconeogenesis in the liver, (4) a role in the acid-base balance homeostasis in the kidney and the liver. Being synthesized by body cells, GLN is formerly classified as a nonessential amino acid.¹⁰⁻¹² Over the past 15 years, GLN has been found to have great interest because of its roles in being a critical substrate for enterocytes and rapid proliferating immune cells, and having immuno-enhancing properties.¹⁵⁻¹⁸ Previous reports showed that a relatively GLN-deficient state occurs during catabolic process, and can be corrected following GLN supplementation.^{19,20} GLN is considered as an essential amino acid

if the patients have certain catabolic, inflammatory and infectious conditions.²¹⁻²³ A study done by Fukatsu et al.²⁴ showed that compared with conventional total parenteral nutrition, GLN-supplemented parenteral nutrition reduced ICAM-1 expression in intestinal homogenates. Another study done by Arndt et al.²⁵ demonstrated that administering GLN reduced leukocyte adhesion and transmigration in indomethacin-induced intestinal inflammation in rats.

To our best knowledge, the GLN effect on CAM expression in preeclampsia has not been investigated. In this study, we are aiming to test the hypotheses that a chronic inflammation during preeclampsia may deplete plasma GLN, and that administering GLN comparable to physiological levels may decrease CAM expression in HUVEC induced by preeclamptic plasma.

MATERIALS AND METHODS

Patients and sample information

The pregnant women were recruited from Taipei Medical University-Wan Fang Medical Center. Women with single pregnancies but without chronic diseases were included in this study. Normal pregnant controls (n = 10) were one without hypertensive and proteinuric conditions, gave birth at between 37-42 weeks and their pregnancy was not complicated with fetal growth retardation or other fetal or maternal problems. Preeclampsia (n = 20) was diagnosed if a patient had: (1) the reading of blood pressure was > 140/90 mmHg on at least two occasions occurring after the 20th week of gestation, and (2) more than 300 mg protein in a 24 hr urine collection or 1+ proteinuria detected on reagent strip on two occasions more than 4 h apart.⁵ The pregnant controls had mean gestational age of 37.6 ± 0.9 weeks (range 36-41 weeks) and preeclamptic women had 37.3 ± 1.9 weeks (range 34-40 weeks). Of those 20

preeclamptic patients, we collected 5 plasma samples for in vitro study from patients with severe preeclampsia, which was defined as blood pressure being higher than 160/110 mmHg at two occasions 6 h apart and having proteinuria corresponding to > 2+ detected on reagent strip on two occasions more than 4 h apart.⁵ The protocol was approved by the TMU-WFMC Ethical Committee, and all subjects provided written informed consent prior to their participation in this study.

HUVEC isolation and culture

HUVECs were isolation as described elsewhere.²⁶ Briefly, we cannulated and washed the umbilical vein with PBS and then perfused with PBS containing 0.1% collagenase for 10 min at 37°C in 5% CO₂. Primary cells were collected and plated into 75 cm² tissue culture flasks with medium-199 (M-199) containing 20% fetal calf serum (FBS), 20mM NaHCO₃, 25mM HEPES, antibiotics (100 U/ml penicillin and 100 µg/ml streptomycin), 10 IU/ml heparin sodium and 15mg/L endothelial cell growth factor at 37°C in 5% CO₂ /95% humidity. Cells were serially passaged 2-3 for experimental assay. HUVEC (1×10⁵ cells/well) from second subcultures were grown on 24-well plate. Every 48-72 h the medium was replaced until the cells reached confluence and were detached with trypsin-ethylenediaminetetra-acetic acid (EDTA). When the cells reached a confluent monolayer, they were then incubated in M-199 (with 20% FBS) with different concentrations of GLN (0, 300, 500 uM) for 24 hr. Subsequently, the cells were washed twice with PBS and cultured with various concentrations of GLN (without FBS) and stimulated with 100uL plasma from severe preeclamptic women. Then, we analyzed ICAM-1 and VCAM-1 expression on HUVEC by flow cytometry.

Measurements and analytical procedures

Plasma amino acid analysis

We took blood samples from the healthy pregnant and preeclamptic women at the time they came to the clinic for obstetric care between 34-36 weeks gestational age. We also collected 6 age-matched nonpregnant women as normal control. Antecubital venous blood was drawn from fasting women into EDTA tubes. Blood was centrifuged at 300g for 10 min at 4°C, and plasma was stored at -70°C until assay. We analyzed plasma amino acids with standard ninhydrin technology (Beckman Instruments, model 6300, Palo Alto, CA), after deproteinization of the plasma with 5% salicylic acid.²⁷

ICAM-1 and VCAM-1 expression of HUVECs

HUVECs surface expression of ICAM-1 and VCAM-1 were measured after stimulated by preeclamptic plasma for 1.5, 3, 4.5 and 6 hr, because the CAM expression reached a peak at 6 hr after stimulation. After removing supernatant, we washed the cells twice with PBS, resuspended them in 100 µl of M-199 (FBS free and containing 5mM EDTA), and incubated for a further 30min at 4°C with fluorescein-conjugated mouse anti-human VCAM-1(CD 106) and phycoerythrin conjugated mouse anti-human ICAM-1(CD 54). The cells suspension was collected into tube and suspended in 500µL of PBS (containing 0.3 ml of 350mM formaldehyde). The fluorescence intensity of 5000 cells population was counted and analyzed with flow cytometry (Coulter, Miami, FL, USA).

Measurement of IL-8 and NO in culture medium

We determined IL-8 concentrations by enzyme-linked immunosorbent assay. The human IL-8 enzyme-linked immunosorbent assay kit was purchased from BioSource International (Camarillo, CA, USA). NO concentrations were determined with a commercial kit (R&D Systems Inc., Minneapolis, MN, USA). NO is a highly unstable molecule, and is converted to stable nitrite and nitrate ions in aqueous solution. After conversion of nitrate to nitrite using nitrate reductase, nitrite

concentrations were measured using the Griess reagent. Procedures are described in the manufacturer's instruction.

Statistical analysis

Data were presented with mean \pm SD. Differences among groups were analyzed by ANOVA using Duncan's test. The differences were considered significant if p value was smaller than 0.05.

RESULTS

Plasma GLN levels

Plasma GLN levels in normal pregnant group were significantly higher than those in the preeclampsia groups (normal pregnant 509.5 ± 52.1 vs. preeclampsia 349.8 ± 55.1 $\mu\text{mol/L}$, $p < 0.05$), but were lower than the normal non-pregnant women (570.5 ± 46.4 $\mu\text{mol/L}$, $p < 0.05$).

CAM expression in HUVEC induced by plasma from preeclamptic women

As shown in Fig. 1, ICAM-1 expression in HUVEC were significantly lower at 500 μM than at 0, 300 μM GLN after incubation for 3, 4.5, and 6 hr ($p < 0.05$, power $> 80\%$). There were no differences among various GLN concentrations at 0 and 1.5 hr after incubation. No differences were observed in HUVEC VCAM-1 expression among various GLN concentrations at each time point (Fig. 2).

IL-8 and NO production from endothelial cells

No differences were found in IL-8 production among various GLN levels at different time points after incubation (Fig. 3). No difference in NO levels were also observed among various GLN concentrations at different time points after incubation (Fig. 4).

Discussion

In this study, we observed that pregnant women had lower plasma GLN levels than non-pregnant controls, possibly result from hemodilution during the pregnancy. The finding of this study also showed that plasma from women with preeclampsia had significantly lower GLN levels than that from normal pregnant women. This finding was compatible with the previous reports that plasma GLN was reduced during a catabolic conditions such as inflammatory, infection, and injury.^{17,21-23} To understand whether physiologic levels of GLN may reduce CAM expression on endothelial cell stimulated by preeclamptic plasma, we pretreatment the endothelial cells with or without GLN including: 0 uM, 300 uM (similar to preeclamptic patients) and 500 uM (similar to normal pregnant women). The findings showed that administering GLN at levels similar to physiological condition reduced HUVEC ICAM-1 expression, but that VCAM-1 did not alter when same concentration of GLN was administered. Since the ICAM-1 and VCAM-1 receptors expressed on immune cells differ,²⁸ the GLN effect on different CAM expression may vary. ICAM-1 is the ligand counterpart of integrins on polymorphonuclear neutrophils, reduced ICAM-1 expression may decrease polymorphonuclear neutrophil-endothelium interaction and thus attenuate tissue injury.⁷

Preeclampsia is a multiple system disorder. Redman et al.²⁹ suggest that preeclampsia is an excessive maternal inflammatory response to pregnancy, and that the endothelial dysfunction is a part of a more generalized intravascular inflammatory reaction. Cytokines which are peptides produced by cells of the immune system that act as mediators of the immune response and the response to inflammation. Previous studies have shown that some cytokines were associated with the initiation of preeclampsia.^{30,31} IL-8 which is a potent chemo-attractant for leukocytes to

implicate inflammation, initiates the acute inflammatory cascade and is considered as an early marker of the inflammatory process.³² Previous report revealed that an increased IL-8 level can increase permeability in endothelial cells from preeclampsia.³¹ Study by Fowler et al.³³ showed that exogenous NO suppresses IL-8 gene expression in activated endothelial cells by inhibiting NF-κB binding to DNA. Since ICAM-1 is regulated transcriptionally by NF-κB, inhibition of NF-κB may result in reduced ICAM-1 expression.⁹ An in vitro study by Huang et al.³⁴ showed that GLN decreases LPS-induced IL-8 production in Caco-2 cells. In this study, we found that pretreatment with 500 μM GLN reduced ICAM-1 expression in vascular endothelial cells 3, 4.5 and 6 hr after stimulation with preeclamptic plasma. However, no differences were observed in IL-8 and NO levels between various GLN concentrations at different time points. Wang et al.³⁵ showed that maternal plasma from pregnancies with umbilical placental vascular disease did not affect endothelial cell expression of NO synthase. A recent study by Wang et al.³¹ revealed that no difference in IL-8 production was found between endothelial cells from normal and preeclamptic pregnancies. Those studies showed that endothelial cell IL-8 and NO production may not be responsible for endothelial cell activation in preeclampsia. Based on the results of this study, we found that GLN did not influence endothelial cell IL-8 and NO production, and that the GLN effect to reduce endothelial ICAM-1 expression was independent of IL-8 and NO production.

GLN which is considered as the main fuel for rapid proliferating cells, can modulate immune response and has antioxidant effect under metabolic stress. Hong et al.³⁶ found that GLN supplemented nutrition protects the liver during hepatic injury by preserving glutathione (GSH) stores. An in vitro study by Babu et al.³⁷ also found that GLN can prevent liver from damage possibly mediated via GSH

synthesis. GSH is a major antioxidant and acts as a vital component in host defense. GLN was found to be rate limiting for GSH synthesis, and the availability of GLN is critical to generate GSH stores.³⁸ Previous study showed that preeclampsia increased oxidative stress, and that plasma from women with preeclampsia had significantly higher malondialdehyde and lipid peroxide levels than those of normal pregnant women.⁹ Lipid peroxide and oxidative stress up-regulate endothelial NF-kB activity and ICAM-1 expression.⁹ Based on the findings of this study, we suggest that the GLN antioxidant property may be partly implicated in reducing HUVEC ICAM-1 expression. Whether other mechanisms are also involved in the favorable effect of GLN on preclamptic plasma-induced CAM expression requires further investigation to clarify.

In summary, our study revealed that plasma from women with preeclampsia had significantly lower GLN levels than normal pregnant women. Administering GLN at levels similar to physiological condition reduces HUVEC ICAM-1 expression stimulated by preeclamptic plasma.

Acknowledgments

The authors wish to thank Ms Chiu-Li Yeh for her technical assistance, and professor Winston W. Shan for editing comment.

References

1. Davey DA, MacGillivray I: The classification and definition of the hypertensive disorders and of pregnancy. *Am J Obstet Gynecol* 1988;158:892-898

2. Roberts JM, Taylor RN, Musci TJ, et al: Preeclampsia: an endothelial cell disorder. *Am J Obstet Gynecol* 1989;161:1200-1204
3. Wang Y, Adair CD, Coe L, et al: Activation of endothelial cells in preeclampsia: increased neutrophil-endothelial adhesion correlates with up-regulation of adhesion molecule P-selectin in human umbilical vein endothelial cells isolated from preeclampsia. *J Soc Gynecol Investig* 1998;5:237-243
4. Coata G, pennacchi L, Bini V, et al: Soluble adhesion molecules: marker of preeclampsia and intrauterine growth restriction. *J Matern-Fetal Neont Med* 2002;12:28-34
5. Djurovic S, Schjetlein R, Wisloff F, et al: Increased levels of intercellular adhesion molecules and vascular cell adhesion molecules in preeclampsia. *Br J Obstet Gynecol* 1999;104:466-470
6. Heyl W, Handt S, Reister F, et al: The role of soluble adhesion molecules in evaluating endothelial cell activation in preeclampsia. *Am J Obstet Gynecol* 1999;180:68-72
7. Carlos T, Harian J: Leukocyte-endothelial adhesion molecules. *Blood* 1994;54:2068-2101
8. Jang Y, Linoff AM, Plow EF, et al: Cell adhesion molecules in coronary artery disease. *J Am Coll Cardiol* 1994;24:1591-1601
9. Takacs P, Kauma SW, Sholley MM, et al: Increased circulating lipid peroxides in severe preeclampsia activate NF- κ B and upregulate ICAM-1 in vascular endothelial cells. *FASEB* 2001;15:279-281
10. Smith RJ, Willmore DW: Glutamine nutrition and requirements. *J Parent Enter*

Nutr 1990;14:94S-99S

11. Souba WW: Glutamine: a key substrate for the splanchnic bed. *Ann Rev Nutr* 1991;11:285-308
12. Smith RJ: Glutamine metabolism and its physiologic importance. *J Parenter Enter Nutr* 1990;14:40S-44S
13. Alverdy JA, Aloys E, Weiss-Carrington P, et al: The effect of glutamine enriched TPN on gut immune cellularity. *J Surg Res* 1992;52:34-38
14. Ardawi MSM, Newsholme EA: Metabolism in lymphocytes and its importance in the immune response. *Eaasys Biochem* 1983;21:1-44
15. Calder PC: Glutamine and the immune system. *Clin Nutr* 1994;13:2-8
16. Heberer M, Babst R, Juretic A, et al: Role of glutamine in the immune response in critical illness. *Nutrition* 1996;12 (Suppl):S71-S72
17. Gismondo MR, Drago L, Fassina MC, et al: Immunostimulating effect of oral glutamine. *Dig Dis Sci* 1998;43:1752-1754
18. Wilmore DW, Shabert JK: Role of glutamine in immunologic responses. *Nutrition* 1998;14:618-626
19. Hammerquist F, Wernerman J, Al R, et al: Addition of glutamine to total parenteral nutrition after elective abdominal surgery spares glutamine in muscle, counteracts the fall in muscle protein synthesis and improves nitrogen balance. *Ann Surg* 1989;209:455-461
20. Gianotti L, Alexander JW, Gennari R, et al: Oral glutamine decreases bacterial translocation and improves survival in experimental gut-origin sepsis. *J Parenter*

Enter Nutr 1995;19:69-74

21. Lacey JM, Wilmore DW: Is glutamine a conditionally essential amino acid? Nutr Rev 1990;48:297-309
22. Willmore DW: The effects of glutamine supplementation in patients following elective surgery and accidental injury. J Nutr 2001;131:2543S-2549S
23. Parry-Billings M, Evans J, Calder PC, et al: Does glutamine contribute to immunosuppression after major burns? Lancet 1990;336:523-525
24. Fukatsu K, Lundberg AH, Kudsk KA, et al: Modulation of organ ICAM-1 expression during IV-TPN with glutamine. Shock 2001;15:24-28
25. Arndt H, Kullmann F, Reuss F, et al: Glutamine attenuates leukocyte-endothelial cell adhesion in indomethacin-induced intestinal inflammation in the rat. J Parenter Enter Nutr 1999;23:12-18
26. Jaffe EA, Nachman RL, Becker CG, et al: Culture of human endothelial cells derived from umbilical veins. Clin Invest 1973;52:2745-2756
27. Smith RJ, Panico K: Automated analysis of o-phthalaldehyde derivatives of amino acids in physiological fluids of reverse phase high performance liquid chromatography. J Liq Chromatogr 1985;8:1783-1795
28. Fickling SA, Whitley GS, Nussey SS: The cell adhesion molecule, VCAM-1, is selectively elevated in serum in pre-eclampsia: does this indicate the mechanism of leukocyte activation? Br J Obstet Gynaecol 1995;102:173-174
29. Redman CW, Sacks GP, Sargent IL: Preeclampsia: an excessive maternal inflammatory response to pregnancy. Am J Obstet Gynecol 1999;180:499-506

30. Ellis J, Wennerholm UB, Bengtsson A, et al: Levels of dimethylarginines and cytokines in mild and severe preeclampsia. *Acta Obstet Gynecol Scand* 2001;80:602-608
31. Wang Y, Gu Y, Zhang Y, et al: Evidence of endothelial dysfunction in preeclampsia: decreased endothelial nitric oxide synthase expression is associated with increased cell permeability in endothelial cells from preeclampsia. *Am J Obstet Gynecol* 2004;190:817-824
32. Baggiolini M: Chemokines in pathology and medicine. *J Intern Med* 2001;250:91-104
33. Fowler AA, Fisher BJ, Sweeney LB, et al: Nitric oxide regulates interleukin-8 gene expression in activated endothelium by inhibiting NF-kappa B binding to DNA: effects on endothelial function. *Biochem Cell Biol* 1999;77:201-208
34. Huang Y, Li N, Liboni K, et al: Glutamine decreases lipopolysaccharide-induced IL-8 production in Caco-2 cells through a non-NF-kappa B p50 mechanism. *Cytokine* 2003;22:77-83
35. Wang X, Athayde N, Trudinger B: Maternal plasma from pregnant women with umbilical placental vascular disease does not affect endothelial cell mRNA expression of nitric oxide synthase. *J Soc Gynecol Investig* 2004;11:149-153
36. Hong RW, Rounds JD, Helton WS, et al: Glutamine preserves liver glutathione after lethal hepatic injury. *Ann Surg* 1992;215:114-119
37. Babu R, Eaton S, Drake DP, et al: Glutamine and glutathione counteract the inhibitory effects of mediators of sepsis in neonatal hepatocytes. *J Pediatr Surg* 2001;36:282-286

38. Welbourne TC: Ammonia production and GLN incorporation into GSH in the functioning rat kidney. *Can J Biochem* 1979;57:233-237

Figure legends

Fig. 1. Intercellular adhesion molecule-1 (ICAM-1) expression (%) in HUVECs with various GLN concentrations at different incubation times. ICAM-1 expression was significantly lower at 500 uM than at 0 and 300 uM GLN after incubation for 3, 4.5, and 6 h. * $p < 0.05$ compared with 0, 300 uM GLN at the same time point.

Fig. 2. Vascular cellular adhesion molecule-1 (VCAM-1) expression (%) in HUVECs with various GLN concentrations at different incubation times. No differences were found among various GLN concentrations at each time point.

Fig. 3. Effect of pretreatment with different concentrations of GLN on IL-8 production from HUVECs after stimulation with preeclamptic plasma. There were no differences among various GLN concentrations at each time point.

Fig. 4. Effect of pretreatment with different concentrations of GLN on nitric oxide (NO) released from HUVECs after stimulation with preeclamptic plasma. There were no differences among various GLN concentrations at each time point.




Fig. 1




Fig.2




Fig. 3




Fig. 4