

行政院國家科學委員會補助專題研究計畫成果報告

PPAR 配位体之抗癌化及抗發炎作用機制

計畫類別：個別型計畫 整合型計畫
計畫編號：NSC 90 - 2320 - B - 038 - 050 -
執行期間：90 年 8 月 1 日至 91 年 7 月 31 日

計畫主持人：梁 有 志
共同主持人：林 仁 混
計畫參與人員：劉正典, 潘憲, 林時宜, 何元順,
洪苓方, 蔡淑慧.

本成果報告包括以下應繳交之附件：

- 赴國外出差或研習心得報告一份
- 赴大陸地區出差或研習心得報告一份
- 出席國際學術會議心得報告及發表之論文各一份
- 國際合作研究計畫國外研究報告書一份

執行單位：台北醫學大學醫學系

中 華 民 國 91 年 10 月 18 日

行政院國家科學委員會專題研究計畫成果報告

PPAR 配位体之抗癌化及抗發炎作用機制

Study on the molecular mechanisms of anti-carcinogenic and anti-inflammatory effects of peroxisome proliferator-activated receptor (PPAR) ligands

計畫編號：NSC 90-2320-B-038-050

執行期限：90 年 8 月 1 日至 91 年 7 月 31 日

主持人：梁 有 志 執行機構及單位名稱：台北醫學大學醫學系

共同主持人：林 仁 混 執行機構及單位名稱：台大醫學院生化所

計畫參與人員：劉正典, 潘憲, 林時宜, 何元順, 洪苓方, 蔡淑慧.

執行機構及單位名稱：台北醫學大學醫學系

一、中文摘要

花生四烯酸的多種前列腺素衍生物已知可經由活化 peroxisome proliferator-activated receptor (PPAR)或其他路徑引起細胞凋亡。本研究計畫試驗了多種 eicosanoids 對細胞增殖的抑制能力。結果發現 prostaglandin J₂ (PGJ₂)的代謝衍生物 15-deoxy- $\Delta^{12,14}$ -PGJ₂ (15d-PGJ₂)及 PGA₁, 可有效抑制胃腫瘤細胞株 AGS 的生長, 且造成此細胞凋亡。過渡表現 PPAR_α 或 PPAR_γ 的細胞, 並不會增加 15d-PGJ₂ 及 PGA₁ 所造成的細胞凋亡, 顯示 15d-PGJ₂ 及 PGA₁ 引起細胞凋亡非經由活化 PPAR 的路徑。然而 15d-PGJ₂ 及 PGA₁ 會依濃度及時間的增加, 提昇 JNK 及 Caspase-3 的活化; 若細胞先轉殖 dominant-negative c-Jun N-terminal kinase (DN-JNK)則可減少 JNK 及 Caspase-3 的活性且降低細胞凋亡的數目, 這些結果顯示 15d-PGJ₂ 及 PGA₁ 可活化 JNK, 且活化 JNK 與造成細胞凋亡有相關性。

關鍵詞：前列腺素、細胞凋亡、JNK

ABSTRACT

Cyclopentenone prostaglandins (CyPGs) derivatives of arachidonic acid have been suggested to exert growth-inhibitory activity through peroxisome proliferator-activated receptor (PPAR)-dependent and -independent mechanisms. Here we examined that various eicosanoids on the inhibition of cell proliferation, and found that the terminal derivative of PGJ₂ metabolism, 15-deoxy- $\Delta^{12,14}$ -PGJ₂ (15d-PGJ₂), and PGA₁ markedly inhibited growth and induced apoptosis in AGS gastric carcinoma cells. There were no significant increase in DNA-fragmentation in the cells with overexpression of PPAR_α or PPAR_γ plasmid, indicating that 15d-PGJ₂ and PGA₁ induced apoptosis through PPAR-independent pathway. Moreover, 15d-PGJ₂ and PGA₁ activated the c-Jun N-terminal kinase (JNK), and the caspase-3 activity in dose- and time-dependent manners. To further examine the role of JNK signaling cascades in apoptosis induced by 15d-PGJ₂ and PGA₁, we transfected dominant-negative (DN) mutants of JNK into the cells to analyze the apoptotic characteristics of cells expressing DN-JNK plasmid following exposure to 15d-PGJ₂ and PGA₁. Expression of DN-JNK proteins repressed both of endogenous JNK and caspase-3 activity, and subsequently inhibited apoptosis induced by 15d-PGJ₂ and PGA₁. These results suggest that CyPGs, such as 15d-PGJ₂ and PGA₁, activates the JNK signaling pathway, and that JNK activation may be involved in 15d-PGJ₂ and PGA₁ induced cell death.

Key words: prostaglandins; apoptosis; JNK

INTRODUCTION

The c-Jun N-terminal kinases (JNKs) members of mitogen-activated protein kinase (MAPKs), are classic stress-activated protein kinase (1), can be activated by proinflammatory cytokines (2, 3) and environmental stress such as UV light (4), γ -irradiation (5), heat shock, osmotic stress (6), shear stress (7), growth factor withdrawal (8), ceramide (9), protein synthesis inhibitor (2), and treatment with chemotherapy drugs including paclitaxel (10), adriamycin, vinblastine (11), and etoposide (12). Recently, JNK had play an important mediator of apoptosis signaling (13). For example, JNK activation was associated with apoptosis in PC-12 cells deprived of NGF (14). In addition, nonneuronal embryonic fibroblasts with targeted disruptions of the *jnk1* and *jnk2* genes, were able to inhibit apoptosis (15). These reports suggested that activation of JNK can be associated with induction of apoptosis, however, the mechanism(s) and functional role for JNK in the regulation of apoptosis are not clear (4).

Prostaglandins are oxygenated metabolites of arachidonic acid (AA), and divided into two groups, conventional prostaglandins and cyclopentenone prostaglandins (CyPGs) according to their mechanisms of action (16). The conventional eicosanoids, such as PGE₂ and PGD₂, bind to cell surface receptor and transduces signal into cells. The CyPGs, such as the A and J series of PGs, lack cell surface receptors but directly enter the cells and exert their actions in the nuclei, including induction of cell growth arrest and cell differentiation (16). Several proteins have been identified to be induced by the CyPGs, such as heat shock proteins (17), γ -glutamylcysteine synthetase (18), collagen (19), gadd 45 (20), and heme oxygenase (21). Arachidonic acid is first converted to PGH₂ by cyclooxygenase and subsequently converted to one of several related products, including PGD₂, PGE₂, PGF₂, PGI₂ and thromboxane A₂, through the action of specific PG synthases. 15d-PGJ₂ is derived from PGD₂ and involves sequential conversion of PGD₂, PGI₂, Δ^{12} -PGJ₂ and 15d-PGJ₂ (16). 15d-PGJ₂ has been shown to bind to and activate peroxisome proliferator-activated receptor γ (PPAR γ) and is therefore an intracellular target of this cyclopentenone prostaglandin (22).

The CyPGs have been demonstrated that possess potent antiproliferative and antitumor activities, but their mechanisms of action are complex and not well understood. Recently it was shown that PGD₂ metabolites, such as PGJ₂, Δ^{12} -PGJ₂, and 15d-PGJ₂, were potential inducers of intracellular oxidative stress that mediated the cytotoxic effects in human neuroblastoma cells (23). Uchida and colleagues showed that the 15d-PGJ₂-induced accumulation of p53 resulted in the activation of a death-inducing caspase cascade mediated by Fas and the Fas ligand in neurons (24). Another studies showed that de novo gene transcription was necessary for 15d-PGJ₂-induced apoptosis in breast cancer cells (25). The current investigation was undertaken to study the effects of arachidonic acid and its metabolites on human gastric cells death. The results demonstrated that 15d-PGJ₂ and PGA₁ induced apoptosis, activated JNK and caspase-3 activity in AGS cells. Moreover, the cells were decrease in apoptosis with overexpression of DN-JNK plasmid, indicating that induction of apoptosis by 15d-PGJ₂ and PGA₁ may be mediated the activation of JNK.

MATERIALS AND METHODS

Materials

PGA₁, PGJ₂, and LTB₄ were purchased from Caymen Chemical (Ann Arbor, MI), and arachidonic acid, PGD₂, PGE₂, PGF₂, PGH₂, TBX₂ and 15d-PGJ₂ were purchased from Biomol (Plymouth Meeting, PA).

Cell Culture

The human AGS gastric carcinoma cell lines (CCRC 60102) were obtained from Food Industry Research and Development Institute, Hsinchu, Taiwan, and cultured in DMEM/F12 containing 10% heat-inactivated fetal bovine serum (Gibco BRL, Grand Island, NY). For all assays, cells were plated in 6-cm dishes at 5×10^6 cells per dish and allowed to grow for 24 h.

Western Blot

Equal amounts of total cellular protein (50 μ g) were resolved by SDS-polyacrylamide gel electrophoresis (PAGE), and transferred onto Immobilon-P membrane (Millipore, Bedford, MA) as described previously (26). The membrane was then incubated with an anti-PARP or JNK antiserum (Santa Cruz Biotechnology, Santa Cruz, CA) and subsequently probed with anti-mouse antibody conjugated to horseradish peroxidase (Santa Cruz Biotechnology) and visualized using enhanced chemiluminescence kits (Amersham, Arlington, IL).

Kinase Assay

Equal amounts of total cellular protein (200 μ g) were immunoprecipitated with JNK1 or ERK1 specific antibody (Santa Cruz Biotechnology) and protein A/G-PLUS agarose for 12 h at 4 °C. Kinase assay was performed in kinase buffer with Gst-c-Jun fusion protein (for JNK, Santa Cruz Biotechnology) or myelin basic protein (for ERK, Sigma) as substrates as previously described (27).

Plasmids and Transfection

The DN-JNK plasmid was constructed from human JNK1 with double point mutant of Thr183 to Ala and Tyr185 to Phe (28). The PPAR γ expression plasmid and PPAR α expression plasmid were generously provided by Professor Christopher K. Glass (California University) and Professor Bart Staels (Université de Lille), respectively (29, 30). AGS cells were seeded in 6-cm dishes, and transfected with the mock, DN-JNK, PPAR γ , or PPAR α expression plasmid using LipofectAMINETM 2000 (Gibco) for 48 h (31). After transfection, the cells were treated with 15d-PGJ₂ or PGA₁, and collected cells for DNA fragmentation, caspase activity, or kinase assay.

DNA Fragmentation

After treatment with tested drugs, the cells were washed with PBS and detected the DNA fragmentation by Cell Death Detection ELISA kits according to the manufacturer's instructions (Roche Diagnostics GmbH, Mannheim, Germany).

Caspase-3 Activity Assay

Briefly, cells were washed with cold PBS twice, and extracted the cytosolic fraction with extraction buffer (12.5 mM Tris, pH7.4, 2 mM DTT, 0.125 mM EDTA, 5% glycerol, and 1 μ g/ml leupeptin, and 1 μ g/ml aprotinin) by repeated cycles of freezing and thawing. Cell lysate (150 μ g) were incubated with 10 μ M fluorescence substrate, Ac-DEVD-AMC (Calbiochem, La Jolla, CA) in assay buffer (50 mM Tris, pH7.4, 1 mM EDTA and 10 mM EGTA) at 37°C for 30 min in dark. The fluorescence intensity of the cleaved substrate was measured using a fluorescence spectrophotometer (Hitachi F-3000) by Ex 380 nm/Em 460 nm (32).

Flow Cytometry

SubG1 population of the cell cycle was analyzed by flow cytometry as described previously (26). Briefly, cells were trypsinized, washed once with PBS, and fixed in 75% ethanol for 1 h at -20°C. Fixed cells were then washed with PBS, incubated with 0.5 ml PBS containing 0.05% RNase and 0.5% Triton X-100 for 30 min at 37°C, and stained with propidium iodide. The stained cells were analyzed using a FACscan laser flow cytometer (Becton Dickinson, San Jose, CA).

RESULTS

15d-PGJ₂ and PGA₁ Inhibited the Growth of Human AGS Gastric Carcinoma Cells

To examine whether CyPGs inhibit growth of AGS cells, we screened a large number of arachidonic acid and its metabolites, and found that PGA₁, the PGD₂ itself and its metabolites, such as PGJ₂ and 15d-PGJ₂ significantly inhibited the growth of AGS cells (Fig. 1A). Among the classes of PGs examined, 15d-PGJ₂ and PGA₁ were most effective. 10 μ M 15d-PGJ₂ and 20 μ M PGA₁ reduced the cell number (Fig. 1) by 80% and 50%, respectively. The terminal metabolites of PGJ₂, 15d-PGJ₂, represented the most potent growth inhibitor. PGD₂ and PGJ₂, the precursors of 15d-PGJ₂, were less effective than 15d-PGJ₂. The inhibitory order was 15d-PGJ₂ > PGA₁ > PGJ₂ > PGD₂. Arachidonic acid, PGE₂, PGF_{2 α} , PGH₂, LTB₄, and TXB₂, were ineffective to inhibit the

growth at the concentration of 50 μM . Thus, a reactive α , β -unsaturated carbonyl group in the cyclopentane ring, such as 15d-PGJ₂, PGA₁, and PGJ₂ (Fig. 1B), which renders this portion of the molecule able to form Michael adducts with cellular nucleophiles and covalently modify specific protein (33, 34), is essential for apoptosis.

We next examined the effects of different concentrations of 15d-PGJ₂ and PGA₁ on the induction of AGS cells apoptosis. As shown in Fig. 2A, 15d-PGJ₂ and PGA₁ induced DNA fragmentation in dose- and time- (Data not shown) dependent manners. DNA fragmentation was approximately 75% and 68% after 12 h incubation with 10 μM 15d-PGJ₂ and 20 μM PGA₁, respectively. Moreover, we measured the numbers of cells with a subG₁ DNA content as a measure of apoptosis and observed an increase in the apoptosis cells (Fig. 2B). Previous studies have indicated that 15d-PGJ₂ activates caspase-3 in a variety of cancer cell types (35-37), so this parameter was monitored as a measure of the rate and extent of apoptosis. As shown in Fig. 2C, caspase-3 activity was detected in AGS cells with 10 μM 15d-PGJ₂ and 20 μM PGA₁ at 8 h. The substrate of caspase-3, poly(ADP ribose) polymerase (PARP) also cleaved in 15d-PGJ₂ (Fig. 2D) and PGA₁ (Data not shown). These results indicated that 15d-PGJ₂ and PGA₁ inhibited the growth through induction of apoptotic cell death and the caspase-3 pathway.

15d-PGJ₂ and PGA₁ Induced Apoptosis Through PPAR-Independent Pathway

Since 15d-PGJ₂ or PGA₁ was a potent agonist of PPAR _{α} and PPAR _{γ} . Activation of PPAR _{α} or PPAR _{γ} can promote cells apoptosis in other cells (38). To examine this possibility, AGS cells were transfected with mock, PPAR _{α} , or PPAR _{γ} expression plasmid, and treated with 15d-PGJ₂ or PGA₁. The DNA fragmentation was measured and shown the results in Fig. 3. 15d-PGJ₂ and PGA₁ induced DNA fragmentation in mock-expression AGS cells, however, it was equally effective in PPAR _{α} or PPAR _{γ} -overexpression AGS cells. Treatment of mock-expression AGS cells with 10 μM 15d-PGJ₂ and 10 μM PGA₁ resulted in induction of DNA fragmentation by 75% and 28%, respectively. However, similar effects of 15d-PGJ₂ and PGA₁ were observed in PPAR _{α} or PPAR _{γ} -overexpression AGS cells. These results indicated that overexpression of PPAR _{α} or PPAR _{γ} did not enhance the apoptosis, and induction of apoptosis may be mediated PPAR-independent pathway in the cells with 15d-PGJ₂ or PGA₁.

15d-PGJ₂ and PGA₁ Activated the JNK and Caspase-3 Activity

Since JNK is a classic stress-activated protein kinase and plays an important role in drugs inducing apoptosis. To examine whether activity of JNK participated in the 15d-PGJ₂ and PGA₁-induced apoptosis, we determined the kinase activity by an immunocomplex kinase assay as described in "Materials and Methods." As shown in Fig. 4, the activation of JNK was observed at the earliest measured time point of 1.5 h at a concentration of 10 μM 15d-PGJ₂ and 20 μM PGA₁. The activation of JNK by 15d-PGJ₂ or PGA₁ was a time- and dose-dependent manner (Fig. 4).

Dominant-Negative Mutants of JNK Blocked Apoptosis Induced by 15d-PGJ₂ or PGA₁

To assess directly the involvement of JNK in apoptosis by 15d-PGJ₂ or PGA₁, dominant-negative (DN) mutants of JNK plasmid were used. As shown in Fig. 5, overexpression of DN-JNK significantly inhibited 15d-PGJ₂ and PGA₁-induced DNA fragmentation and caspase-3 activity. To verify that overexpression of DN-JNK efficiently decreased endogenous JNK activity, we determined the total JNK activity in both mock and DN-JNK expression cells by immunocomplex kinase assay. As shown in Fig. 5C, transfection of DN mutant of JNK exactly resulted in the reduction of JNK activity about > 50% induced by 15d-PGJ₂ and PGA₁. These results indicated that the JNK-mediated induction of apoptosis is important for the treatment of 15d-PGJ₂ and PGA₁ in AGS cells.

DISCUSSION

We here show that 15d-PGJ₂ and PGA₁-induced apoptosis in human AGS gastric carcinoma cells requires the activation of JNK. Activation of PPAR _{γ} has also been reported to induce apoptosis in various cell types, including choriocarcinoma cells, breast cancer cells, endothelial cells and

macrophages (39-41). Recently, Harris and Phipps showed that PGD₂ and 15d-PGJ₂ have potent antiproliferative effects through activation of PPAR_γ (42). However, here we have described a different, PPAR-independent mechanism that explains the ability of 15d-PGJ₂ and PGA₁ to induce apoptosis in AGS cells.

The specified J₂-derivatives are potential metabolites of PGD₂ that are sequentially formed from this prostaglandin. A comparison of the PG biosynthetic pathway and the induction of apoptosis profile revealed that induction of apoptosis was mediated mainly by the metabolites of PGD₂, the most active of which is the terminal metabolite of 15d-PGJ₂. The induction potency was 15d-PGJ₂ > PGJ₂ > PGD₂, indicating a gain in biological potency as the catabolism of PGD₂ proceeds. CyPGs are reactive compounds that possess a α, β-unsaturated carbonyl group in the cyclopentenone ring. These groups are very susceptible to nucleophilic addition reactions with sulfhydryl groups of cysteine residues of proteins or of the tripeptide glutathione (43). Since it has been demonstrated that PGA₁ formed a bisconjugate with thiol-containing proteins (44), and 15d-PGJ₂ formed a covalent adduct with the p50 subunit of NFκB (45), which resulted in modulation of protein functions. Based on functional evidence, a modification of IKK (46) and of the p65 subunit of NFκB (47) by 15d-PGJ₂ has also been proposed. In this study, we showed that 15d-PGJ₂ and PGA₁ activated the JNK pathway. Thus, whether 15d-PGJ₂ and PGA₁ activate JNK via interaction with JNK itself or upstream proteins of JNK deserves further study.

Activation of the JNK signaling pathway has been mechanistically implicated in regulation of apoptosis. In contrast, several reports also indicated that JNK play a role in promoting or preventing apoptosis, depending on both cell type and apoptotic-triggering stimuli (48-50). On the other hand, JNK activation may be a stress response secondary to apoptosis itself, rather than a primary mediator in apoptotic pathway (50). However, we showed that activation of JNK reached a peak at early 2 h with 15d-PGJ₂ and PGA₁, in which the apoptotic cells are still minimal (Data not shown). These results suggested that JNK activation was not a secondary response to 15d-PGJ₂ and PGA₁-induced apoptosis. In addition, 15d-PGJ₂ and PGA₁ caused a more sustained pattern of JNK activation (Fig. 4), suggesting that activation of JNK was directly linked to cell death in 15d-PGJ₂ and PGA₁ treatment cells. Since several reports have demonstrated that the duration of JNK activation is a determining factor for cell proliferation or death (4). Transient JNK activation led to cell proliferation not death, such as activation of JNK by PMA plus ionomycin.

In summary, we have found that 15d-PGJ₂ and PGA₁ represent the most potent inducers of apoptosis in AGS cells, and that JNK is directly involved in 15d-PGJ₂ and PGA₁-induced program cell death. The present study may therefore represent a first step in establishing a link between the activation of JNK and apoptosis induced by 15d-PGJ₂ and PGA₁.

REFERENCES

1. Hibi M, Lin A, Smeal T, Minden A, Karin M. Identification of an oncoprotein- and UV-responsive protein kinase that binds and potentiates the c-Jun activation domain. *Genes Develop* 1993;7:2135-48.
2. Kyriakis JM, Banerjee P, Nikolakaki E, Dai T, Rubie EA, Ahmad MF, Avruch J, Woodgett JR. The stress-activated protein kinase subfamily of c-Jun kinases. *Nature* 1994;369:156-60.
3. Raingeaud J, Gupta S, Rogers JS, Dickens M, Han J, Ulevitch RJ, Davis RJ. Pro-inflammatory cytokines and environmental stress cause p38 mitogen-activated protein kinase activation by dual phosphorylation on tyrosine and threonine. *J Biol Chem* 1995;270:7420-26.
4. Chen YR, Meyer CF, Tan TH. Persistent activation of c-Jun N-terminal kinase 1 (JNK1) in gamma radiation-induced apoptosis. *J Biol Chem* 1996;271:631-4.
5. Zanke BW, Boudreau K, Rubie E, Winnett E, Tibbles LA, Zon L, Kyriakis J, Liu FF, Woodgett JR. The stress-activated protein kinase pathway mediates cell death following injury induced by cis-platinum, UV irradiation or heat. *Curr Biol* 1996;6:606-13.
6. Galcheva-Gargova Z, Derijard B, Wu IH, Davis RJ. An osmosensing signal transduction

- pathway in mammalian cells. *Science* 1994;265:806-8.
7. Li YS, Shyy JY, Li S, Lee J, Su B, Karin M, Chien S. The Ras-JNK pathway is involved in shear-induced gene expression. *Mol Cell Biol* 1996;16:5947-54.
 8. Xia Z, Dickens M, Raingeaud J, Davis RJ, Greenberg ME. Opposing effects of ERK and JNK-p38 MAP kinases on apoptosis. *Science* 1995;270:1326-31.
 9. Verheij M, Bose R, Lin XH, Yao B, Jarvis WD, Grant S, Birrer MJ, Szabo E, Zon LI, Kyriakis JM, Haimovitz-Friedman A, Fuks Z, Kolesnick RN. Requirement for ceramide-initiated SAPK/JNK signalling in stress-induced apoptosis. *Nature* 1996;380:75-9.
 10. Srivastava RK, Mi QS, Hardwick JM, Longo DL. Deletion of the loop region of Bcl-2 completely blocks paclitaxel-induced apoptosis. *Proc Nat Acad Sci USA* 1999;96:3775-80.
 11. Fan M, Goodwin ME, Birrer MJ, Chambers TC. The c-Jun NH₂-terminal protein kinase/AP-1 pathway is required for efficient apoptosis induced by vinblastine. *Cancer Res* 2001;61:4450-8.
 12. Osborn MT, Chambers TC. Role of the stress-activated/c-Jun NH₂-terminal protein kinase pathway in the cellular response to adriamycin and other chemotherapeutic drugs. *J Biol Chem* 1996;271:30950-5.
 13. Davis RJ. Signal transduction by the JNK group of MAP kinases. *Cell* 200;103:239-52.
 14. Le-Niculescu H, Bonfoco E, Kasuya Y, Claret FX, Green DR, Karin M. Withdrawal of survival factors results in activation of the JNK pathway in neuronal cells leading to Fas ligand induction and cell death. *Mol Cell Biol* 1999;19:751-63.
 15. Tournier C, Hess P, Yang DD, Xu J, Turner TK, Nimmual A, Bar-Sagi D, Jones SN, Flavell RA, Davis RJ. Requirement of JNK for stress-induced activation of the cytochrome c-mediated death pathway. *Science* 2000;288:870-4.
 16. Fukushima M. Biological activities and mechanisms of action of PGJ₂ and related compounds: an update. *Prostaglandin Leukotri Essent Fatty Acids* 1992;47:1-12.
 17. Santoro MG, Garaci E, Amici C. Prostaglandins with antiproliferative activity induce the synthesis of a heat shock protein in human cells. *Proc Nat Acad Sci USA* 1989;86:8407-11.
 18. Ohno K, Higaki J, Takechi S, Hirata M. Specific role of an alpha,beta-unsaturated carbonyl group in gamma-glutamylcysteine synthetase induction by prostaglandin A₂. *Chem Biol Interact* 1990;76:77-87.
 19. Tasaki Y, Takamori R, Koshihara Y. Prostaglandin D₂ metabolite stimulates collagen synthesis by human osteoblasts during calcification. *Prostaglandins* 1991;41:303-13.
 20. Ohtani-Fujita N, Minami S, Mimaki S, Dao S, Sakai T. p53-Independent activation of the gadd45 promoter by Delta¹²-prostaglandin J₂. *Biochem Biophys Res Commun* 1998;251:648-52.
 21. Koizumi T, Odani N, Okuyama T, Ichikawa A, Negishi M. Identification of a cis-regulatory element for delta 12-prostaglandin J₂-induced expression of the rat heme oxygenase gene. *J Biol Chem* 1995;270:21779-84.
 22. Kliewer SA, Lenhard JM, Willson TM, Patel I, Morris DC, Lehmann JM. A prostaglandin J₂ metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 1995;83:813-9.
 23. Kondo M, Oya-Ito T, Kumagai T, Osawa T, Uchida K. Cyclopentenone prostaglandins as potential inducers of intracellular oxidative stress. *J Biol Chem* 2001;276:12076-83.
 24. Kondo M, Shibata T, Kumagai T, Osawa T, Shibata N, Kobayashi M, Sasaki S, Iwata M, Noguchi N, Uchida K. 15-Deoxy-Δ^{12,14}-prostaglandin J₂: The endogenous electrophile that induces neuronal apoptosis. *Proc Nat Acad Sci USA* 2002;99:7367-72.
 25. Clay CE, Atsumi GI, High KP, Chilton FH. Early de novo gene expression is required for 15-deoxy-Delta 12,14-prostaglandin J₂-induced apoptosis in breast cancer cells. *J Biol Chem* 2001;276:47131-5.
 26. Liang YC, Lin-Shiau SY, Chen CF, Lin JK. Inhibition of cyclin-dependent kinases 2 and 4 activities as well as induction of Cdk inhibitors p21 and p27 during growth arrest of human breast carcinoma cells by (-)-epigallocatechin-3-gallate. *J Cell Biochem* 1999;75:1-12.

27. Liang YC, Huang YT, Tsai SH, Lin-Shiau SY, Chen CF, Lin JK. Suppression of inducible cyclooxygenase and inducible nitric oxide synthase by apigenin and related flavonoids in mouse macrophages. *Carcinogenesis* 1999;20:1945-52.
28. Lin SY, Liang YC, Ho YS, Tsai SH, Pan S, Lee WS. Involvement of both extracellular signal-regulated kinase and c-Jun N-terminal kinase pathways in the 12-o-tetradecanoylphorbol-13-acetate-induced upregulation of p21^{Cip1} in colon cancer cells. *Mol Carcinogenesis* 2002; In press.
29. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature* 1998;391:79-82.
30. Chinetti G, Lestavel S, Bocher V, Remaley AT, Neve B, Torra IP, Teissier E, Minnich A, Jaye M, Duverger N, Brewer HB, Fruchart JC, Clavey V, Staels B. PPAR-alpha and PPAR-gamma activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway. *Nature Med.* 2001;7:53-8.
31. Liang YC, Tsai DC, Lin-Shiau SY, Chen CF, Lin JK. Suppression of inducible cyclooxygenase and inducible nitric oxide synthase through peroxisome proliferator-activated receptor- γ by apigenin and related-flavonoids in macrophages. *FEBS Lett* 2001;496:12-8.
32. Shiah SG, Chuang SE, Kuo ML. Involvement of Asp-Glu-Val-Asp-directed, caspase-mediated mitogen-activated protein kinase kinase 1 cleavage, c-Jun N-terminal kinase activation, and subsequent bcl-2 phosphorylation for paclitaxel-induced apoptosis in HL-60 cells. *Mol Pharmacol* 2001;59:254-62.
33. Fukushima M. Prostaglandin J2--anti-tumour and anti-viral activities and the mechanisms involved. *Eicosanoids* 1990;3:189-99.
34. Rossi A, Elia G, Santoro MG. 2-Cyclopenten-1-one, a new inducer of heat shock protein 70 with antiviral activity. *J Biol Chem* 1996;271:32192-6.
35. Rohn TT, Wong SM, Cotman CW, Cribbs DH. 15-deoxy-delta12,14-prostaglandin J2, a specific ligand for peroxisome proliferator-activated receptor-gamma, induces neuronal apoptosis. *Neuro Report* 2001;12:839-43.
36. Chattopadhyay N, Singh DP, Heese O, Godbole MM, Sinohara T, Black PM, Brown EM. Expression of peroxisome proliferator-activated receptors (PPARS) in human astrocytic cells: PPARgamma agonists as inducers of apoptosis. *J Biol Chem* 2000;61:67-74.
37. Keelan JA, Sato TA, Marvin KW, Lander J, Gilmour RS, Mitchell MD. 15-Deoxy-Delta (12,14)-prostaglandin J(2), a ligand for peroxisome proliferator-activated receptor-gamma, induces apoptosis in JEG3 choriocarcinoma cells. *Biochem Biophys Res Commun* 1999;262:579-85.
38. Chinetti G, Griglio S, Antonucci M, Torra IP, Delerive P, Majd Z, Fruchart JC, Chapman J, Najib J, Staels B. Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages. *J Biol Chem* 1998;273:25573-80.
39. Elstner E, Müller C, Koshizuka K, Williamson EA, Park D, Asou H, Shintaku P, Said JW, Heber D, Koeffler DP. Ligands for peroxisome proliferator-activated receptor and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells in vitro and in BNX mice. *Proc Nat Acad Sci USA* 2001;95:8806-11.
40. Bishop-Bailey D, Hla T. Endothelial cell apoptosis induced by the peroxisome proliferator-activated receptor (PPAR) ligand 15-deoxy-Delta12, 14-prostaglandin J2. *J Biol Chem* 1999;274:17042-8.
41. Chinetti G, Griglio S, Antonucci M, Torra IP, Delerive P, Majd Z, Fruchart JC, Chapman J, Najib J, Staels B. Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages. *J Biol Chem* 1998;273:25573-80.
42. Harris SG, Phipps RP. Prostaglandin D(2), its metabolite 15-d-PGJ(2), and peroxisome proliferator activated receptor-gamma agonists induce apoptosis in transformed, but not normal, human T lineage cells. *Immunology* 2002;105: 23-34.
43. Atsmon J, Sweetman BJ, Baertschi SW, Harris TM, Roberts LJ 2nd. Formation of thiol

- conjugates of 9-deoxy-delta 9,delta 12(E)-prostaglandin D2 and delta 12(E)-prostaglandin D2. *Biochemistry* 1990;29:3760-5.
44. Parker J, Ankel H. Formation of a prostaglandin A2-glutathione conjugate in L1210 mouse leukemia cells. *Biochem Pharmacol* 1992;43:1053-60.
 45. Cernuda-Morollon E, Pineda-Molina E, Canada FJ, and Perez-Sala D. 15-Deoxy-Delta 12,14-prostaglandin J2 inhibition of NF-kappaB-DNA binding through covalent modification of the p50 subunit. *J Biol Chem* 2001;276:35530-6.
 46. Rossi A, Kapahi P, Natoli G, Takahashi T, Chen Y, Karin M, Santoro MG. Anti-inflammatory cyclopentenone prostaglandins are direct inhibitors of IkappaB kinase. *Nature* 2000;403:103-8.
 47. Straus DS, Pascual G, Li M, Welch JS, Ricote M, Hsiang CH, Sengchanthalangsy LL, Ghosh G, Glass CK. 15-deoxy-delta 12,14-prostaglandin J2 inhibits multiple steps in the NF-kappa B signaling pathway. *Proc Nat Acad Sci USA* 2000;97:4844-9.
 48. Zanke BW, Lee C, Arab S, Tannock IF. Death of tumor cells after intracellular acidification is dependent on stress-activated protein kinases (SAPK/JNK) pathway activation and cannot be inhibited by Bcl-2 expression or interleukin 1beta-converting enzyme inhibition. *Cancer Res* 1998;58:2801-8.
 49. Nishina H, Fischer KD, Radvanyi L, Shahinian A, Hakem R, Rubie EA, Bernstein A, Mak TW, Woodgett JR, Penninger JM. Stress-signalling kinase Sek1 protects thymocytes from apoptosis mediated by CD95 and CD3. *Nature* 1997;385:350-3.
 50. Ip YT, Davis RJ. Signal transduction by the c-Jun N-terminal kinase (JNK)--from inflammation to development. *Curr Opin Cell Biol* 1998;10:205-19.

LEGENDS TO FIGURES

Fig. 1. Effect of arachidonic acid (AA) and its metabolites on the growth of AGS cells. (A). Cells were treated with 50 μ M AA, LTB₄, PGD₂, PGE₂, PGF_{2 α} , PGH₂ or TXB₂, or 20 μ M PGA₁, PGJ₂, or 10 μ M 15d-PGJ₂ for 24 h. Cell numbers were measured by counting the number of trypan blue-excluding cells. The values were expressed as mean \pm S.E. of triplicate test. *, $p < 0.05$ versus control. (B). Structures of arachidonic acid and its metabolites.

Fig. 2. Effect of 15d-PGJ₂ and PGA₁ on the DNA fragmentation, caspase-3 activity, and PARP cleavage in AGS cells. Cells were treated with different concentration of PGA₁ or 15d-PGJ₂ for 15 h. (A). DNA fragmentation was determined by the histone-associated DNA-fragments as described in "Materials and Methods." The values were expressed as mean \pm S.E. of triplicate test. *, $p < 0.05$ versus control. (B). Cells were treated with 5 μ M 15d-PGJ₂ or 10 μ M PGA₁, and subjected to DNA content analysis by flow cytometer. The percentages of total cells with subG1 apoptotic DNA are indicated. (C). Cells were treated with 10 μ M 15d-PGJ₂ or 20 μ M PGA₁ for 4 h and 8 h as indicated, and cell extracts were subjected to caspase-3 assay as described in "Materials and Methods". The values were expressed as mean \pm S.E. of triplicate test. *, $p < 0.05$ versus control. (D). Cells were treated with different concentrations of 15d-PGJ₂ for 15 h. Total cell lysates (50 μ g) were subjected to Western blot by using anti-PARP antibody. The intact (112 kD) and cleaved (86 kD) species of PARP were indicated in right of the panel.

Fig. 3. Effect of overexpression of PPAR α or PPAR γ plasmid on the induction of DNA fragmentation by 15d-PGJ₂ or PGA₁. Cells were transfected with PPAR α or PPAR γ expression plasmid for 48 h, and treated with different concentrations of 15d-PGJ₂ or PGA₁ for 15 h. DNA fragmentation was determined by the histone-associated DNA-fragments as described in "Materials and Methods." The values were expressed as mean \pm S.E. of triplicate test. *, $p < 0.05$ versus mock control.

Fig. 4. Effect of 15d-PGJ₂ and PGA₁ on the activation of JNK in AGS cells. (A). Cells were treated with 10 μ M 15d-PGJ₂ or 20 μ M PGA₁ for various times as indicated. (B). Cells were treated with different concentrations of 15d-PGJ₂ or PGA₁ for 3 h. JNK1 was immunoprecipitated from cell lysates and incubated with purified GST-c-Jun fusion protein as substrate in an in vitro kinase assay

as described in “Materials and Methods”.

Fig. 5. Effect of dominant-negative JNK on the blocking of endogenous JNK activity, DNA fragmentation, and caspase-3 activity induced by 15d-PGJ₂ or PGA₁. Cells were transfected with mock expression plasmid or DN-JNK expression plasmid for 48 h. (A). Transfected cells were treated with 10 μM 15d-PGJ₂ and 20 μM PGA₁ for 15 h. DNA fragmentation was determined by the histone-associated DNA-fragments as described in “Materials and Methods.” The values were expressed as mean ± S.E. of triplicate test. *, $p < 0.05$ versus control. (B). Transfected cells were treated with 10 μM 15d-PGJ₂ or 20 μM PGA₁ for 3 h or 8 h, and determined the caspase-3 activity (8 h), and JNK activity (3 h) as described in “Methods and Methods.” The values were expressed as mean ± S.E. of triplicate test. *, $p < 0.05$ versus control.

Fig. 1 Liu et al.

A.





Fig. 1 Liu et al.

B.

Fig. 2 Liu et al.

A.



B.





Fig. 2 Liu et al.

C.



D.





Fig. 3 Liu et al.

A.



B.




Fig. 4 Liu et al.




Fig. 5 Liu et al.

