

行政院國家科學委員會專題研究計畫 成果報告

前列環素(Prostacyclin)類似物 Beraprost 對阿黴素
(Adriamycin)所引發腎小管細胞凋亡的抑制作用之研究

計畫類別：個別型計畫

計畫編號：NSC94-2314-B-038-033-

執行期間：94年08月01日至95年07月31日

執行單位：臺北醫學大學醫學系

計畫主持人：許永和

共同主持人：陳正憲

報告類型：精簡報告

處理方式：本計畫可公開查詢

中 華 民 國 95 年 10 月 30 日

行政院國家科學委員會補助專題研究計畫 成果報告
 期中進度報告

(計畫名稱)

前列環素(Prostacyclin)類似物Beraprost 對阿黴素(Adriamycin)所引發腎小管細胞凋亡的抑制作用之研究

計畫類別： 個別型計畫 整合型計畫

計畫編號：NSC 94-2314-B-038-033-

執行期間： 94 年 8 月 1 日至 95 年 7 月 31 日

計畫主持人：許永和

共同主持人：陳正憲

計畫參與人員：許永和、陳正憲

成果報告類型(依經費核定清單規定繳交)： 精簡報告 完整報告

本成果報告包括以下應繳交之附件：

- 赴國外出差或研習心得報告一份
- 赴大陸地區出差或研習心得報告一份
- 出席國際學術會議心得報告及發表之論文各一份
- 國際合作研究計畫國外研究報告書一份

處理方式：除產學合作研究計畫、提升產業技術及人才培育研究計畫、列管計畫及下列情形者外，得立即公開查詢

涉及專利或其他智慧財產權， 一年 二年後可公開查詢

執行單位：台北醫學大學

中 華 民 國 95 年 9 月 25 日

中文摘要

阿黴素(Adriamycin)是一種抗癌的化療藥物，廣泛地應用在惡性腫瘤和淋巴瘤的治療上。大鼠經阿黴素的處理下，會很快地引發持續性的腎絲球病變，並表現出許多與人類慢性腎臟病變相同的特性，因此此系統常應用在慢性腎臟病的研究中。而阿黴素所造成的細胞毒害，細胞凋亡為其中重要的原因之一。前列環素(Prostacyclin)是由血管內皮細胞所分泌，透過細胞表面的受器作用在血小板與血管，得以抑制血小板的功能，擴張血管。另外也有許多報告顯示，前列環素對多種細胞具保護效果。本研究主要研究阿黴素在大鼠腎小管細胞中所引發細胞凋亡的分子機制與訊號路徑，以及前列環素的類似物 Beraprost 對阿黴素引起之腎小管細胞細胞凋亡的保護作用。而表現 COX-1 及 PGIS 以增加內生性前列環素之系統亦應用於此研究中。研究結果顯示，內生性前列環素增加，對阿黴素所引發的細胞凋亡具保護作用。另外前列環素對阿黴素所引發的 caspase-3 及 caspase-9 之活性、cytochrome c 的釋放、以及 Bcl-x_L 的減少均有抑制的效果。而此種抑制效果極可能是透過對轉譯因子 NF-κB 活性的抑制來達成的。本研究顯示了前列環素及 Beraprost 在阿黴素引發之病變，以及人類慢性腎臟疾病上的治療潛力。

關鍵詞：前列環素，Beraprost，阿黴素，細胞凋亡，腎小管細胞

英文摘要

Adriamycin-induced nephrosis in rats is a commonly used experimental model for pharmacological studies of human chronic renal diseases. Adriamycin-induced apoptosis of renal tubular cells has been reported in adriamycin-treated rats. In addition, prostacyclin (PGI₂) is known to have various protective effects on many kinds of cells. To investigate the protective effect of PGI₂ on cells undergoing adriamycin-induced apoptosis, the PGI₂ agonist beraprost and the selectively PGI₂ augmentation *via* adenovirus-mediated transfer of genes for cyclooxygenase-1 (COX-1) and prostacyclin synthase (PGIS) (two key enzymes of PGI₂ synthesis) to renal tubular cells were applied in this study. The PGI₂ overexpression protected rat renal tubular cells from adriamycin-induced apoptosis. Ad-COX-1/PGIS transfection was found to reduce the adriamycin-stimulated activities of caspase-3 and caspase-9, inhibit adriamycin-induced release of cytochrome *c*, elevate the expression of Bcl-x_L, and suppress the activation and translocation of nuclear factor-kappaB (NF-κB) in adriamycin-treated renal tubular cells. Our results reveal that selective augmentation of PGI₂ production can protect rat renal tubular cells from adriamycin-induced apoptosis *via* the NF-κB signaling pathway. This implies the therapeutic potential of PGI₂ and beraprost for chronic renal diseases.

Keywords: Prostacyclin (PGI₂); Beraprost; Adriamycin; Renal tubular cell; Apoptosis

報告内容

1. Introduction

Adriamycin is the anti-tumor anthracycline antibiotic of choice for the treatment of many solid malignancies and lymphomas. Rats treated with adriamycin develop heart failure as well as a self-perpetuating glomerular nephropathy. Even in the absence of continued adriamycin exposure, the glomerular damage progresses and late-onset tubular lesions are observed (Bertani et al., 1982; Scholey et al., 1989). Within a few weeks after adriamycin administration, the glomerular filtration rate declines gradually and the animals develop a nephrotic and a tubular syndrome. A long-term study of this pathological change in rats demonstrated severe renal damage with characteristic features of chronic progressive renal diseases in humans (Bertani et al., 1982; Okuda et al., 1986). Adriamycin-induced nephrosis in rats, therefore, is a common experimental model used for pharmacological studies of human chronic renal diseases. However, the correlative molecular mechanism of this model is not very well understood.

Induction of apoptosis is an important cytotoxic mechanism of adriamycin (Muller et al., 1998). The apoptosis of renal tubular cells has been reported in adriamycin-treated rats (Zhang et al., 1996). Renal tubular cell apoptosis is a key feature of tubular atrophy, which is a hallmark of chronic renal diseases (Khan et al., 1999; Schelling et al., 1998). Reactive oxygen species derived from redox activation of adriamycin is a proposed cause of adriamycin cytotoxicity (Singal et al., 2000). Previous studies have indicated that, at submicromolar concentrations, adriamycin induces apoptosis with the activation of caspases in endothelial cells and myocytes (Kotamraju et al., 2000; Sawyer et al., 1999). In mammalian cells, a major caspase activation pathway is the cytochrome *c*-initiated pathway. In this pathway, a variety of apoptotic stimuli cause cytochrome *c* release from mitochondria, which in turn induces a series of biochemical reactions that result in caspase activation and subsequent cell death (Jiang and Wang, 2004). Cytochrome *c* release is known to be regulated by Bcl-2 family proteins, including Bcl-2 and Bcl-x_L, which bind to the mitochondrial outer membrane and block cytochrome *c* efflux (Yang et al., 1997). Therefore, mitochondria-mediated apoptosis signaling plays a major role in adriamycin-induced cytotoxicity.

Prostacyclin (PGI₂), largely produced in vascular endothelial cells, acts on platelets and blood vessels through its specific cell surface receptor, thereby inhibiting platelet function, dilating blood vessels, and protecting the vascular endothelium (Moncada, 1982). PGI₂ is also known to inhibit leukocyte functions such as migration and reactive oxygen species production (Boxer et al., 1980) and inhibit mesangial cell proliferation (Mene et al., 1990). PGI₂ is produced by the cyclooxygenase (COX) system in which COX converts arachidonic acid to PGH₂, and PGH₂ is subsequently converted to PGI₂ by the action of PGI₂ synthase (PGIS) (Vane and Botting, 1995). The production of PGI₂ is executed by either COX-1 or COX-2 coupled to PGIS (Smith et al., 2000). Because of its unstable property and valuable clinical implications in vascular physiology, several synthetic analogues of PGI₂ with more stable chemical structures have been developed (Sturzebecher et al., 1986). Among these analogues, beraprost has been reported to prevent radiocontrast nephropathy in LLC-PK1 cells (Yano et al., 2005).

The possibility of protecting the heart during the administration of adriamycin by prior administration of prostacyclin has been reported (Dowd et al., 2001). We suppose that PGI₂ is able to protect kidney cells from adriamycin-induced injury. To investigate the possible effect of PGI₂ on adriamycin-induced apoptosis and its potential in a gene therapy treatment for renal tubular atrophy, an adenovirus-mediated gene transfer of PGI₂ was chosen in this study to provide a more physiologically relevant augmentation of PGI₂ production. Previous experimental work (Shyue et al., 2001) used a single gene transfer to augment PGI₂ production. Overexpression of COX-1 alone was accompanied by overproduction of PGE₂ (a key proinflammatory mediator that may contribute to vascular inflammation), whereas overexpression of PGIS alone had a minimal effect on increasing PGI₂ synthesis. In contrast, combined COX-1/PGIS gene transfer selectively augmented PGI₂ production and is thus better than the single gene transfer approach (Lin et al., 2002). Here, the adenovirus-mediated bicistronic COX-1/ PGIS gene transfer was executed in renal tubular cells, and a selective augmentation of PGI₂ production was found. The protective effect of endogenous PGI₂ on adriamycin-induced apoptosis was evaluated by analyzing the apoptosis profile of renal tubular cells. Our results reveal that the selective augmentation of endogenous PGI₂ production can reduce adriamycin-induced apoptosis. This implies the potential of combined COX-1 and PGIS gene transfer as a gene therapy for chronic renal diseases.

2. Materials and methods

2.1. Materials

Dulbecco's modified Eagle's medium (DMEM), fetal calf serum, and tissue culture reagents were from Invitrogen Corporation (PO, USA). All other chemicals of reagent grade were obtained from Sigma (MO, USA). Antibodies used in this research were purchased from BD Laboratories (CA, USA) and Santa Cruz Biotechnology (CA, USA).

2.2. Cell culture

Rat renal proximal tubular cells (NRK-52E) were purchased from Food Industry Research and Development Institute (Taiwan), and cultured in DMEM supplemented with antibiotic/antifungal solution and 10% fetal bovine serum. They were grown until the monolayer became confluent. The medium for the cultured cells was then changed to the serum-free medium, and the cells were incubated overnight before the experiment.

2.3. Preparation of replication-defective recombinant adenoviral vectors

We constructed in the replication-defective recombinant adenoviral (rAd) vector with two separate human phosphoglycerate kinase (HPGK) promoters (bicistronic) to drive COX-1 and PGIS (Ad-COX-1/PGIs), and a HPGK alone to serve as control (Ad-HPGK) as previously described (Lin et al., 2002). Replication-defective rAd vectors were generated by homologous recombination and amplified in 293 cells as described previously (Lin et al., 2002). rAd stocks were prepared by CsCl gradient centrifugation, aliquoted, and stored at -80°C. Viral titers were determined by a plaque-assay method. Two hundred ninety-three cells were infected with serially diluted viral preparations and then overlaid with low melting-point agarose after infection. Numbers of plaques formed were counted within 2 weeks.

2.4. Measurements of eicosanoids by enzyme immunoassay

Cells were sonicated in 1 ml of ice-cold buffer (0.05M Tris at pH 7.0, 0.1M NaCl, and

0.02M EDTA) and centrifuged at 55,000Xg for 1 h. The supernatant was acidified and passed through a Sep-Pak C18 cartridge. Eicosanoids were eluted with 100% methanol, dried under nitrogen gas, redissolved in a small amount of buffer, and analyzed using 6-keto-PGF_{1α}, PGE₂, and 15-d-PGJ₂ ELISA kits from R&D Systems Inc.

2.5. DAPI stain

The cells grown on slides were washed twice in phosphate buffered saline (PBS) for 1 min. The slides were overlaid with DAPI (4'-6-diamidino-2-phenylindole) (1 µg/ml) in PBS plus 0.5% 1,4-diazabicyclo[2,2,2]octane and analyzed immediately.

2.6. TUNEL Stain

Adriamycin-mediated apoptosis in NRK-52E cells was detected by enzymatic labeling of DNA strand breaks using terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick end-labeling (TUNEL). The cells grown on slides were washed twice in PBS for 1 min. Following the incubation of slides with the permeabilisation solution (0.1% Triton X-100 in 0.1% sodium citrate) for 8 min at 4°C and washing twice with PBS for 5 min, the labeling reaction was performed using 50 µl of TUNEL reagent for each sample, except the negative control, in which reagent without enzyme was added and incubated for 1 h at 37°C. Following PBS washing, slides were incubated with converter reagent for 30 min at 37°C. After washing, slides were incubated with Fast Red substrate solution for 10 min to stain cells containing labeled DNA strand breaks. TUNEL labeling was conducted using a Cell Death Detection kit (Roche; Mannheim, Germany) and performed according to the manufacturer's instructions.

2.7. Western blot analysis

For Western blot analysis, the protein concentration of each sample was measured using Bio-Rad Protein Assay Dye (Bio-Rad Laboratories, Inc., CA, USA) according to the manufacturer's directions. A total of 30 µg of NRK-52E lysate proteins were applied to each lane and analyzed by Western blotting. Cytosol cytochrome *c* was extracted by using a Cytochrome *c* Release Apoptosis Assay Kit (Calbiochem Inc., CA, USA). The antibodies of caspase-3, caspase-9, cytochrome *c*, Bcl-2 and nuclear factor-kappaB (NF-κB) were diluted to 1:1000 for assay. Peroxidase-conjugated anti-rabbit or anti-mouse IgG (1:5000 dilution) was used as the second antibody to detect caspase-3, caspase-9, cytochrome *c*, Bcl-x_L, and NF-κB bands by enhanced chemiluminescence (Amersham Biosciences Corp, NJ, USA).

2.8. Lactate dehydrogenase cytotoxicity assay

For lactate dehydrogenase (LDH) cytotoxicity assays, NRK-52E cells were plated at 10,000 cells/well in 96-well plates and grown overnight. The culture medium from cells treated with adriamycin was collected and assayed using the LDH Cytotoxicology Detection kit (Roche) according to the manufacturer's directions. Each data point was determined in triplicate.

2.9. Electrophoretic mobility shift assay (EMSA)

To prepare nuclear protein extracts, cultured NRK-52E cells were washed with cold PBS and then immediately removed by scraping in PBS. After centrifugation of the cell suspension at 1000Xg, the cell pellets were resuspended in cold buffer A (containing KCl 10 mM, EDTA 0.1 mM, DTT 1 mM, and PMSF 1 mM) for 15 min. The cells were lysed by adding 10% NP-40 and

then centrifuged at 5000Xg to obtain pellets of nuclei. The nuclear pellets were resuspended in cold buffer B (containing HEPES 20 mM, EDTA 1 mM, DTT 1 mM and PMSF 1 mM, and NaCl 0.4 mM), vigorously agitated, and then centrifuged. The supernatant containing the nuclear proteins was used for the Western blot assay or stored at -70°C until used. A double-stranded containing a high affinity sequence for NF-κB from the mouse kappa-light chain enhancer (5'AGC TTC AGA GAC TTT CCG AGA GG3') was prepared. The oligonucleotides were end-labeled with [³²P]ATP. Extracted nuclear proteins (10 μg) were incubated with 0.1 ng of ³²P-labeled DNA for 15 min at room temperature in 25 μL of binding buffer containing 1 μg of poly (dI-dC). The mixtures were electrophoresed on 5% non-denaturing polyacrylamide gels. Gels were dried and imaged by autoradiography.

2.10. Statistical analysis

Analysis of variance (ANOVA) was used to compare the levels of eicosanoids. The level of differences among groups was analyzed by Student *t* tests. *P* <0.05 was considered statistically significant.

3. Results

3.1. Increase of COX-1 and PGIS protein and prostacyclin caused by Adv-COX-1/PGIS transfection in NRK-52E cells

NRK-52E cells were transfected with Ad-COX-1/PGIS at different pfu/cell to determine the transfection quantity of Ad-COX-1/PGIS sufficient for COX-1 and PGIS over-expression. The transfected cells were lysed, and COX-1 and PGIS protein levels were determined by Western blot analysis. The transgenic COX-1 and PGIS protein levels were markedly elevated by Ad-COX-1/PGIS transfection with 20 moi (multiplicity of infection; pfu per cell), and gradually increased along raising transfection dose (Fig. 1A). The efficiency of gene expression of adenoviral administration in rat renal tubular cells was also evaluated by determining COX-1 and PGIS protein levels 1 to 3 days after administration. Compared with Adv-HPGK control, Adv-COX-1/PGIS augmented COX-1 and PGIS protein levels in a time-dependent manner (Fig. 1B). Maximal augmentation was noted at 72 h after administration. Several prostanoids in NRK-52E were measured at different times or at different transfection doses of adenoviral administration. The production of PGI₂ was typically monitored by measurement of 6-keto-prostaglandin F_{1α} (6-keto-PGF_{1α}) because 6-keto-PGF_{1α} is a stable product of the non-enzymatic hydration of PGI₂. Adv-COX-1/PGIS increased PGI₂ levels in a time- and dose-dependent manner (Fig. 2), which was similar to the COX-1 and PGIS expression patterns. However, the PGE₂ and 15-d-PGJ₂ levels were not significantly affected by Adv-COX-1/PGIS transfection. Apparently, Adv-COX-1/PGIS transfection can selectively augment the endogenous PGI₂ production in rat renal tubular cells.

3.2. Protective effect of Ad-COX-1/PGIS transfection against adriamycin-induced apoptotic injury in NRK-52E cells

To determine the cytotoxicity of adriamycin in rat renal tubular (NRK-52E) cells, the lactate dehydrogenase (LDH) released from the cytosol of damaged cells was measured. NRK-52E cells were treated with 1.5 μM, 3 μM, and 6 μM of adriamycin from 6 to 18 h. As shown in Fig. 3A, there was already significant injury to cells treated with as little as 3 μM of adriamycin for 6 h,

and the increase in adriamycin cytotoxicity paralleled the increase in its dose. TUNEL staining indicated that NRK-52E cell injury was caused by adriamycin-induced apoptosis (Fig. 3B). Subsequently, the protective effect of Ad-COX-1/PGIS was also examined using an LDH detection system. As shown in Fig. 3C, Ad-COX-1/PGIS transfection reduced the cytotoxicity induced by adriamycin treatment. This result reveals that the endogenous PGI₂ increase caused by Ad-COX-1/PGIS transfection protects rat renal tubular cells from adriamycin-induced apoptosis.

3.3. Effect of Ad-COX-1/PGIS Transfection on Adriamycin-induced Activities of Caspases in NRK-52E Cells

Caspase-dependent apoptotic signaling plays a major role in adriamycin-induced apoptotic injury (Kotamraju et al., 2000; Sawyer et al., 1999). The activities of caspase-3 and caspase-9 stimulated by adriamycin were also evaluated by monitoring the quantity of cleaved subtypes of caspase-3 and caspase-9 in NRK-52E cells. As shown in Fig. 4A, the cleaved subtypes of both caspase-3 and caspase-9 were very significantly elevated in the cells treated with adriamycin, even for 4 h. Compared with Ad-HPGK transfection, Ad-COX-1/PGIS transfection significantly reduced the quantity of cleaved subtypes of both caspase-3 and caspase-9 in adriamycin-treated NRK-52E cells (Fig. 4B). That is, adriamycin stimulates the activities of caspase-3 and caspase-9 in rat renal tubular cells, and endogenous PGI₂ increase caused by Ad-COX-1/PGIS transfection reduces these activities.

3.4. Effect of Ad-COX-1/PGIS transfection on mitochondria-mediated apoptosis signaling induced by adriamycin in NRK-52E Cells

Cytochrome *c* and Bcl-2 family proteins are supposed to be important in mitochondria-mediated apoptosis signaling. To evaluate the influence of transfection on the variation of cytochrome *c* and Bcl-x_L caused by adriamycin, NRK-52E cells transfected with Ad-HPGK or Ad-COX-1/PGIS were treated with 1.5 μM adriamycin for different periods. In Ad-HPGK transfected NRK-52E cells, released cytochrome *c* was increased significantly within 4 h (Fig. 5). However, in Ad-COX-1/PGIS transfected cells, this increase was inhibited and expression of Bcl-x_L was even elevated.

3.5. Inhibition effect of Ad-COX-1/PGIS transfection on NF-κB translocation caused by adriamycin in NRK-52E cells

NF-κB is known as an important transcriptional factor. Recent evidence has shown that NF-κB activation and translocation are pro-apoptotic in adriamycin-treated endothelial cells and cardiomyocytes (Wang et al., 2002). To evaluate the effect of Ad-COX-1/PGIS transfection on NF-κB translocation caused by adriamycin, NRK-52E cells transfected with different doses of viral vectors were treated with 3 μM adriamycin for 6 h, and NF-κB translocated in nuclei was monitored by Western blotting. As shown in Fig. 6A, adriamycin greatly elevated NF-κB in Ad-HPGK-transfected NRK-52E cell nuclei. This nuclear NF-κB elevation was totally blocked by 20 and 40 moi of Ad-COX-1/PGIS. The activation of NF-κB caused by adriamycin was also measured in terms of DNA-binding activity using the Electrophoretic Mobility Shift Assay (EMSA). NRK-52E cells were transfected with 20 moi of viral vectors for 2 days, and then incubated with 3 μM adriamycin for different time periods. The DNA-binding activity was monitored in nuclear extracts. As shown in Fig. 6B, the DNA binding activity of NF-κB

increased in NRK-52E cells treated with adriamycin for 4 and 12 h. This increase in DNA-binding activity of NF- κ B was reduced apparently by Ad-COX-1/PGIS transfection. This result reveals that NF- κ B translocation and activation present in adriamycin-treated NRK-52E cells are inhibited by PGI₂ endogenously augmented by Ad-COX-1/PGIS transfection.

4. Discussion

Results from this study indicate that Adv-COX-1/PGIS transfection is effective in augmenting COX-1 and PGIS expression in rat renal tubular (NRK-52E) cells and in reducing apoptotic death caused by adriamycin. Adv-COX-1/PGIS transfection increases PGI₂ without overproduction of other prostanoids in rat renal tubular cells. PGI₂ level augmented by Ad-COX-1/PGIS transfection presumably accounts for this protective action, as PGI₂ is a potent protective effector for many kinds of cell injury (Boxer et al., 1980; Mene et al., 1990; Moncada, 1982; Yano et al., 2004). Besides monitoring reduction in adriamycin-mediated death of NRK-52E cells, also monitored were the apoptotic signals associated with adriamycin, such as the activation of caspase-3 and caspase-9, increase in cytochrome *c* release, and decrease in Bcl-x_L. These apoptotic signals present in adriamycin-treated NRK-52E cells are reversed by Ad-COX-1/PGIS transfection. Even the activation and translocation of transcription factor NF- κ B induced by adriamycin are also inhibited by Ad-COX-1/PGIS transfection. It is supposed that endogenous selective PGI₂ augmentation caused by Ad-COX-1/PGIS transfection protects adriamycin-treated rat renal tubular cells through the inhibition of NF- κ B.

Previous studies have shown that prostacyclin synthesis can be selectively augmented by cotransfecting endothelial and cerebral cells with COX-1 and PGIS (Lin et al., 2002; Shyue et al., 2001). Our data indicate that the cotransfection of these two enzymes also selectively augments prostacyclin synthesis in rat renal tubular cells, and protects cells from adriamycin injury. COX-1/PGIS cotransfection is potentially a therapeutic strategy for reducing adriamycin-induced nephropathy. However, the etiology of adriamycin-induced nephrotoxicity is multiple and includes glomerular and interstitial inflammation and fibrosis (Bertani et al., 1982; Okuda et al., 1986; Wang et al., 2000). In the present study, we focused only on the effect of endogenous PGI₂ augmentation on the direct toxic action of adriamycin on renal tubular cells. Still unknown is whether COX-1/PGIS cotransfection is effective in adriamycin-induced nephropathy *in vivo*. Nevertheless, renal tubular cell apoptosis is a key feature of tubular atrophy, which is a hallmark of chronic renal diseases (Khan et al., 1999; Schelling et al., 1998). Our results suggest that PGI₂ could be useful in the therapy of a chronic nephropathy *in vivo*. In addition, systemic administration of PGI₂ and its analogues is associated with undesirable side effects. Prostacyclin acts locally in an autocrine and paracrine manner. Delivery of PGI₂ to the targeted vascular region would be preferable. Local administration of PGI₂ and its more stable analogs remains a challenge because of the relatively short half-life of these drugs. On the other hand, gene transfer approaches can prolong expression of PGI₂ synthetic enzymes and thereby maintain drug concentrations for a longer period than the drugs themselves. Therefore, COX-1/PGIS cotransfection has potential as an *in vivo* therapeutic approach.

Similar to the findings in endothelial cells and myocytes (Kotamraju et al., 2000; Sawyer et al., 1999), the adriamycin-induced apoptosis in NRK-52E cells is dependent on the activation of

caspase-3 and caspase-9, since the cleavage subforms of both caspase-3 and caspase-9 are markedly induced by adriamycin. Caspase-9 is considered to be a critical apoptosis regulator in a variety of cells. Activated caspase-9 cleaves procaspase-3 to its active form, which in turn, stimulates the caspase-dependent deoxyribonuclease by inactivating its inhibitor and thereby allowing chromosomal degradation (Enari et al., 1998; Sakahira et al., 1998). Notably, Ad-COX-1/PGIS transfection inhibited the adriamycin-induced activation of caspase-9 and caspase-3. On the other hand, caspase-9 is activated from procaspase-9 by cytosolic cytochrome *c* (Jiang and Wang, 2004). The mitochondrial release of cytochrome *c* is regulated by Bcl-2 family proteins, including Bcl-2 and Bcl-x_L, which bind to the mitochondrial outer membrane and block cytochrome *c* efflux (Yang et al., 1997). In our results, adriamycin markedly reduced the Bcl-x_L expression, which was reversed by Ad-COX-1/PGIS transfection. Therefore, endogenous PGI₂ augmentation likely attenuates adriamycin-induced renal tubular cell injury by acting on a site upstream to the one for Bcl-x_L expression.

In the present study, we demonstrate that adriamycin induces NF-κB activation and translocation in rat renal tubular cells. NF-κB has been reported to be involved in regulating adriamycin-induced apoptosis in various cancer cells and carcinomas (Arlt et al., 2001; Manna and Aggarwal, 1999; Somerville and Cory, 2000). Adriamycin-induced NF-κB activation in tumour cells is anti-apoptotic. Inhibition of NF-κB activation sensitizes cancer cells to adriamycin-induced apoptosis (Arlt et al., 2001; Manna and Aggarwal, 1999; Somerville and Cory, 2000). The present data provide evidence that NF-κB activation promotes adriamycin-induced apoptosis in rat renal tubular cells. The proapoptotic character of NF-κB might be due to its direct activation of apoptotic genes, including Fas ligand, Fas, *c-Myc* and p53 (Chan et al., 1999; Qin et al., 1999), or to downregulation of the activities of some anti-apoptotic factors, e.g. Bcl-x_L (Hettmann et al., 1999). The involvement of NF-κB in regulating the cell cycle can also be a mechanism for its proapoptotic effect (Qin et al., 1999). In addition, recent studies reveal that H₂O₂ is responsible for adriamycin-induced NF-κB activation in adriamycin-treated endothelial cells and cardiomyocytes (Wang et al., 2002). The ability of PGI₂ to inhibit reactive oxygen species production may be the reason our results showed Ad-COX-1/PGIS transfection inhibited adriamycin-induced NF-κB activation. However, the signal-transduction pathways involved in this inhibition remain to be determined.

In summary, Adv-COX-1/PGIS transfection can selectively augment the endogenous PGI₂ production in rat renal tubular cells. Through the inhibition of NF-κB, this endogenous PGI₂ increase enhances Bcl-x_L expression and inhibits cytochrome *c* release, reduces the activity of caspases, and eventually protects rat renal tubular cells from adriamycin-induced apoptosis. Combined COX-1/PGIS gene transfer is a potentially useful gene therapy for chronic renal diseases.

References

- Arlt, A., Vorndamm, J., Breitenbroich, M., Folsch, U.R., Kalthoff, H., Schmidt, W.E., Schafer, H., 2001. Inhibition of NF-kappaB sensitizes human pancreatic carcinoma cells to apoptosis induced by etoposide (VP16) or doxorubicin. *Oncogene*. 20, 859-868.
- Bertani, T., Poggi, A., Pozzoni, R., Delaini, F., Sacchi, G., Thoua, Y., Mecca, G., Remuzzi, G., Donati, M.B., 1982. Adriamycin-induced nephrotic syndrome in rats: sequence of pathologic events. *Lab. Invest.* 46, 16-23.
- Boxer, L.A., Allen, J.M., Schmidt, M., Yoder, M., Baehner, R.L., 1980. Inhibition of polymorphonuclear leukocyte adherence by prostacyclin. *J.*

- Chan, H., Bartos, D.P., Owen-Schaub, L.B., 1999. Activation-dependent transcriptional regulation of the human Fas promoter requires NF-kappaB p50-p65 recruitment. *Mol. Cell. Biol.* 19, 2098-2108.
- Dowd, N.P., Scully, M., Adderley, S.R., Cunningham, A.J., Fitzgerald, D.J., 2001. Inhibition of cyclooxygenase-2 aggravates doxorubicin-mediated cardiac injury in vivo. *J. Clin. Invest.* 108, 585-590.
- Enari, M., Sakahira, H., Yokoyama, H., Okawa, K., Iwamatsu, A., Nagata, S., 1998. A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD. *Nature* 391, 43-50.
- Hettmann, T., DiDonato, J., Karin, M., Leiden, J.M., 1999. An essential role for nuclear factor kappaB in promoting double positive thymocyte apoptosis. *J. Exp. Med.* 189, 145-158.
- Jiang, X., Wang, X., 2004. Cytochrome C-mediated apoptosis. *Annu. Rev. Biochem.* 73, 87-106.
- Khan, S., Cleveland, R.P., Koch, C.J., Schelling, J.R., 1999. Hypoxia induces renal tubular epithelial cell apoptosis in chronic renal disease. *Lab. Invest.* 79, 1089-1099.
- Kotamraju, S., Konorev, E.A., Joseph, J., Kalyanaraman, B., 2000. Doxorubicin-induced apoptosis in endothelial cells and cardiomyocytes is ameliorated by nitron spin traps and ebselen. Role of reactive oxygen and nitrogen species. *J. Biol. Chem.* 275, 33585-33592.
- Lin, H., Lin, T.N., Cheung, W.M., Nian, G.M., Tseng, P.H., Chen, S.F., Chen, J.J., Shyue, S.K., Liou, J.Y., Wu, C.W., Wu, K.K., 2002. Cyclooxygenase-1 and bicistronic cyclooxygenase-1/prostacyclin synthase gene transfer protect against ischemic cerebral infarction. *Circulation* 105, 1962-1969.
- Manna, S.K., Aggarwal, B.B., 1999. Lipopolysaccharide inhibits TNF-induced apoptosis: role of nuclear factor-kappaB activation and reactive oxygen intermediates. *J. Immunol.* 162, 1510-1518.
- Mene, P., Abboud, H.E., Dunn, M.J., 1990. Regulation of human mesangial cell growth in culture by thromboxane A2 and prostacyclin. *Kidney Int.* 38, 232-239.
- Moncada, S., 1982. Eighth Gaddum Memorial Lecture. University of London Institute of Education, December 1980. Biological importance of prostacyclin. *Br. J. Pharmacol.* 76, 3-31.
- Muller, I., Niethammer, D., Bruchelt, G., 1998. Anthracycline-derived chemotherapeutics in apoptosis and free radical cytotoxicity (Review). *Int. J. Mol. Med.* 1, 491-494.
- Okuda, S., Oh, Y., Tsuruda, H., Onoyama, K., Fujimi, S., Fujishima, M., 1986. Adriamycin-induced nephropathy as a model of chronic progressive glomerular disease. *Kidney Int.* 29, 502-510.
- Qin, Z.H., Chen, R.W., Wang, Y., Nakai, M., Chuang, D.M., Chase, T.N., 1999. Nuclear factor kappaB nuclear translocation upregulates c-Myc and p53 expression during NMDA receptor-mediated apoptosis in rat striatum. *J. Neurosci.* 19, 4023-4033.
- Sakahira, H., Enari, M., Nagata, S., 1998. Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis. *Nature* 391, 96-99.
- Sawyer, D.B., Fukazawa, R., Arstall, M.A., Kelly, R.A., 1999. Daunorubicin-induced apoptosis in rat cardiac myocytes is inhibited by dexrazoxane. *Circ. Res.* 84, 257-265.
- Schelling, J.R., Nkemere, N., Kopp, J.B., Cleveland, R.P., 1998. Fas-dependent fratricidal apoptosis is a mechanism of tubular epithelial cell deletion in chronic renal failure. *Lab. Invest.* 78, 813-824.
- Scholey, J.W., Miller, P.L., Rennke, H.G., Meyer, T.W., 1989. Effect of converting enzyme inhibition on the course of adriamycin-induced nephropathy. *Kidney Int.* 36, 816-822.
- Shyue, S.K., Tsai, M.J., Liou, J.Y., Willerson, J.T., Wu, K.K., 2001. Selective augmentation of prostacyclin production by combined prostacyclin synthase and cyclooxygenase-1 gene transfer. *Circulation* 103, 2090-2095.
- Singal, P.K., Li, T., Kumar, D., Danelisen, I., Iliskovic, N., 2000. Adriamycin-induced heart failure: mechanism and modulation. *Mol. Cell Biochem.* 207, 77-86.
- Smith, W.L., DeWitt, D.L., Garavito, R.M., 2000. Cyclooxygenases: structural, cellular, and molecular biology. *Annu. Rev. Biochem.* 69, 145-182.

- Somerville, L., Cory, J.G., 2000. Enhanced roscovitine-induced apoptosis is mediated by a caspase-3-like activity in deoxyadenosine-resistant mouse leukemia L1210 cells. *Anticancer Res.* 20, 3347-3355.
- Sturzebecher, S., Haberey, M., Muller, B., Schillinger, E., Schroder, G., Skuballa, W., Stock, G., Vorbruggen, H., Witt, W., 1986. Pharmacological profile of a novel carbacyclin derivative with high metabolic stability and oral activity in the rat. *Prostaglandins* 31, 95-109.
- Vane, J.R., Botting, R.M., 1995. Pharmacodynamic profile of prostacyclin. *Am. J. Cardiol.* 75, 3A-10A.
- Wang, S., Kotamraju, S., Konorev, E., Kalivendi, S., Joseph, J., Kalyanaraman, B., 2002. Activation of nuclear factor-kappaB during doxorubicin-induced apoptosis in endothelial cells and myocytes is pro-apoptotic: the role of hydrogen peroxide. *Biochem. J.* 367, 729-740.
- Wang, Y., Wang, Y.P., Tay, Y.C., Harris, D.C., 2000. Progressive adriamycin nephropathy in mice: sequence of histologic and immunohistochemical events. *Kidney Int.* 58, 1797-1804.
- Yang, J., Liu, X., Bhalla, K., Kim, C.N., Ibrado, A.M., Cai, J., Peng, T.L., Jones, D.P., Wang, X., 1997. Prevention of apoptosis by Bcl-2: release of cytochrome c from mitochondria blocked. *Science* 275, 1129-1132.
- Yano, T., Itoh, Y., Kubota, T., Sendo, T., Koyama, T., Fujita, T., Saeki, K., Yuo, A., Oishi, R., 2005. A prostacyclin analog prevents radiocontrast nephropathy via phosphorylation of cyclic AMP response element binding protein. *Am. J. Pathol.* 166, 1333-1342.
- Yano, T., Itoh, Y., Kubota, T., Sendo, T., Oishi, R., 2004. A prostacyclin analog beraprost sodium attenuates radiocontrast media-induced LLC-PK1 cells injury. *Kidney Int.* 65, 1654-1663.
- Zhang, J., Clark, J.R., Jr., Herman, E.H., Ferrans, V.J., 1996. Doxorubicin-induced apoptosis in spontaneously hypertensive rats: differential effects in heart, kidney and intestine, and inhibition by ICRF-187. *J. Mol. Cell Cardiol.* 28, 1931-1943.

Figure legends

Fig. 1. COX-1 and PGIS protein levels in NRK-52E cells transfected with Ad-COX-1/PGIS. (A) COX-1 and PGIS protein levels in NRK-52E cells transfected with Ad-COX-1/PGIS at different moi for 3 days. (B) Time course of COX-1 and PGI₂ protein levels in NRK-52E cells transfected with Ad-COX-1/PGIS at 20 moi. Protein levels in transfected cells were determined by Western blot analysis with specific antibodies.

Fig. 2. Prostanoid levels in NRK-52E cells transfected with Ad-COX-1/PGIS. (A) Time course of prostanoid levels in NRK-52E cells infected with 20 moi of adenoviral particles. (B) Prostanoid levels in NRK-52E cells transfected with Ad-COX-1/PGIS at different moi for 2 days. Adv-HPGK transfection was included as a control. A bar is mean \pm S.D. of 3 experiments. *, $P < 0.05$ compared with the expression level of 6-keto-PGF_{1 α} in control cells.

Fig. 3. Adriamycin-induced injury in NRK-52E cells. (A) Cytotoxicity induced by adriamycin in NRK-52E cells. NRK-52E cells were treated with 1.5 μ M, 3 μ M and 6 μ M of adriamycin from 6 to 18 h. The lactate dehydrogenase (LDH) released from the cytosol of damaged cells was measured to determine the cytotoxicity of adriamycin. A bar is mean \pm S.D. of 3 experiments. #, $P < 0.05$ compared with control cells at 6h. &, $P < 0.05$ compared with control cells at 12h. *, $P < 0.05$ compared with control cells at 18h. (B) Adriamycin-induced apoptosis in NRK-52E as shown by the TUNEL assay. NRK-52E cells were treated for 6 h with 3 μ M adriamycin as indicated, harvested, stained with DAPI and TUNEL, and examined by fluorescence microscopy as described under "Experimental Procedures" (original magnification, \times 100). (C) The protective

effect of Ad-COX-1/PGIS transfection against the cytotoxicity of adriamycin in NRK-52E cells. NRK-52E cells were transfected with 20 moi of Ad-COX-1/PGIS or Ad-HPGK for 2 days, and then treated with 3 μ M adriamycin from 6 to 18 h. A bar is mean \pm S.D. of 3 experiments. &, $P < 0.05$ compared with Ad-HPGK control at 12h. *, $P < 0.05$ compared with Ad-HPGK control at 18h.

Fig. 4. Western blot of cleaved caspase-9 and caspase-3 in adriamycin-treated NRK-52E cells. (A) Protein levels of cleaved caspase-9 and caspase-3 in NRK-52E cells treated with adriamycin under various conditions. NRK-52E cells were treated with 3 μ M adriamycin for 4, 6, or 12 h, or 6 h at 1.5, 3, or 6 μ M as indicated. (B) Protein levels of cleaved caspase-9 and caspase-3 in adenoviral transfected NRK-52E cells treated with adriamycin. NRK-52E cells were transfected 2 days with 20 moi of Ad-COX-1/PGIS (CP) or Ad-HPGK (H) as a control, and treated with 3 μ M adriamycin for 4, 6, or 12 h as indicated. Western blotting was carried out with the specific antibody against cleaved caspase-9 and caspase-3.

Fig. 5. The levels of released cytochrome *c* and Bcl-x_L in adenoviral transfected NRK-52E cells. NRK-52E cells were transfected 2 days with 20 moi of Ad-COX-1/PGIS (CP) or Ad-HPGK (H) as a control, and treated with 3 μ M adriamycin for 4, 6, or 12 h as indicated. Cytosol cytochrome *c* was extracted as described in methods section. Western blotting was carried out with the specific antibodies against cytochrome *c* and Bcl-x_L.

Fig. 6. The translocation and activation of NF- κ B in adenoviral transfected NRK-52E cells treated with adriamycin. (A) The levels of nuclear NF- κ B in adenoviral transfected NRK-52E cells treated with adriamycin. NRK-52E cells were transfected with Ad-COX-1/PGIS or Ad-HPGK at varied moi as indicated for 2 days, and then treated with 3 μ M adriamycin for 6 h. Nuclear protein was extracted as described in methods section. Western blotting was carried out with the specific antibody against NF- κ B p65. The quantity of nuclear NF- κ B represents the level of NF- κ B translocation. (B) The DNA-binding activity of NF- κ B in adenoviral transfected NRK-52E cells treated with adriamycin. NRK-52E cells were transfected with 20 moi of Ad-COX-1/PGIS (CP) or Ad-HPGK (H) for 2 days, and then incubated with 3 μ M adriamycin for 4 and 12 h. The nuclear proteins were extracted and analyzed by EMSA with NF- κ B binding nucleotides. The DNA-binding activity of NF- κ B is responsible for the majority of NF- κ B activity.

Figures

Fig. 1




Fig 2




Fig 3




Fig 4




Fig 5





Fig 6



計畫成果自評

本研究內容完全達到原計畫預期之目標、不但在學術上具創新性，並且顯示了前列環素在腎臟病上的臨床應用潛力。本研究結果已發表於國際期刊 *European Journal of Pharmacology* 529 (2006, Jan) 8–15。