

• 系統編號	RN9704-1299		
• 計畫中文名稱	調控鈣子釋放(Ryanodine 接受器)對肺靜脈心肌細胞之鈣子恆定、鈣子點與引發心整活性之作用		
• 計畫英文名稱	Role of Calcium Release Channel (Ryanodine Receptor) and Regulators on Calcium Homeostasis, Calcium Spark and Arrhythmogenesis of Pulmonary Vein Cardiomyocytes		
• 主管機關	行政院國家科學委員會	• 計畫編號	NSC95-2314-B038-026
• 執行機構	台北醫學大學醫學系		
• 本期期間	9508 ~ 9607		
• 報告頁數	23 頁	• 使用語言	英文
• 研究人員	陳亦仁; 陳耀昌 Chen, Yi-jen ; Chen, Yao-Chang		
• 中文關鍵字	--		
• 英文關鍵字	--		
• 中文摘要	查無中文摘要		
• 英文摘要	<p>3.1. The $[Ca^{2+}]_i$ transient and Calcium stores in PV and Atrial Cardiomyocytes Figure 1A shows the tracings of the $[Ca^{2+}]_i$ transient in the PV and LA cardiomyocytes. The PV cardiomyocytes with pacemaker activity (n = 15) had a larger amplitude of the $[Ca^{2+}]_i$ transient (R410/485, 0.36 ± 0.04, 0.25 ± 0.02, 0.24 ± 0.02, P < 0.05), and higher peak systolic (R410/485, 1.16 ± 0.05, 0.92 ± 0.03, 0.96 ± 0.02, P < 0.01) and diastolic $[Ca^{2+}]_i$ concentration (R410/485, 0.79 ± 0.02, 0.67 ± 0.02, 0.73 ± 0.01, P < 0.01) than did the PV cardiomyocytes without pacemaker activity (n = 12) 8 and LA cardiomyocytes (n = 12). In addition, the PV cardiomyocytes with pacemaker activity had a longer time to the peak of the $[Ca^{2+}]_i$ transient than did the PV cardiomyocytes without pacemaker activity and LA cardiomyocytes (46 ± 3, 35 ± 3, 34 ± 3, P < 0.05). The tauCa in the PV cardiomyocytes with pacemaker activity was shorter than that in the PV cardiomyocytes without pacemaker activity and LA cardiomyocytes (64 ± 5, 117 ± 13, 119 ± 9 ms, P<0.01). However, the amplitudes of the $[Ca^{2+}]_i$ transient, peak systolic $[Ca^{2+}]_i$, diastolic $[Ca^{2+}]_i$, time to the peak of the $[Ca^{2+}]_i$ transient, and tauCa were not significantly different between the LA cardiomyocytes and PV cardiomyocytes without pacemaker activity. The administration of caffeine induced a rapid rise in the $[Ca^{2+}]_i$ in the PV and LA cardiomyocytes (Figure 1B). The PV cardiomyocytes with pacemaker activity (n = 8) had a larger SR Ca²⁺ store than did the PV cardiomyocytes</p>		

without pacemaker activity ($n = 16$) and LA cardiomyocytes ($n = 13$, $R_{410/480}$, 0.61 ± 0.03 , 0.42 ± 0.02 , 0.40 ± 0.02 , $P < 0.05$). However, there was a similar SR Ca^{2+} store between the PV cardiomyocytes without pacemaker activity and the LA cardiomyocytes. The time constant of the caffeine-induced $[\text{Ca}^{2+}]_i$ decay was similar among the LA and PV cardiomyocytes with and without pacemaker activity.

3.2. Effect of magnesium on the $[\text{Ca}^{2+}]_i$ transients

In the PV cardiomyocytes without pacemaker and LA cardiomyocytes, magnesium (1.8 and 5.4 mM) reduced the amplitude of the $[\text{Ca}^{2+}]_i$ transient and peak systolic $[\text{Ca}^{2+}]_i$, in a concentration-dependent manner but did not change the diastolic $[\text{Ca}^{2+}]_i$. In contrast, only magnesium with a high concentration (5.4 mM) reduced the $[\text{Ca}^{2+}]_i$ transient and peak systolic $[\text{Ca}^{2+}]_i$ in the PV cardiomyocytes with pacemaker activity (Figure 2A and 2B). Moreover, as the examples show in figure 2C, magnesium (5.4 mM) could also reduce the spontaneous beating rates in the PV cardiomyocytes from 1.5 ± 0.4 Hz to 0.9 ± 0.3 Hz ($n = 3$, $P < 0.05$).