

行政院國家科學委員會專題研究計劃成果報告

篩選活化 PPAR γ 之天然化合物及其活化路徑之研究

Screening PPAR γ ligands from natural products and study on the activation pathways

計劃編號: NSC 89-2320-B-038-075-

執行期限: 89 年 9 月 1 日 至 90 年 7 月 31 日

主持人: 梁有志 助教授 台北醫學大學醫學系

內容: 與本計畫有關之研究成果 (published results)

Suppression of inducible cyclooxygenase and nitric oxide synthase through activation of peroxisome proliferator-activated receptor- γ by flavonoids in mouse macrophages. Liang, Y.C., Tsai S.H., Tsai, D.C., Lin-Shiau, S.Y. and Lin, J.K. FEBS Letters, 496: 12-18, 2001.

摘要

PPAR α (過氧化體增殖活化受體) 的活化與抗發炎作用有關。在本次計畫中，篩選 20 餘種類黃素，發現三種類黃素 apigenin, chrysin 及 kaempferol 能有效地活化 PPAR α 。在吞噬細胞中，過度表現 PPAR α ，可增強此三種類黃素之抑制 LPS 所活化的 COX-2 及 iNOS。然而在體外的競爭結合分析，發現此三種類黃素只有微弱的 PPAR α agonist 活性。有限度蛋白分解試驗，顯示此三種類黃素會改變 PPAR α 的構形，但不屬於結合 BRL49653 之 PPAR α 。這些結果顯示，此三種類黃素可作為 PPAR α 之 allosteric effectors，結合在 PPAR α 上，並活化之，但結合的位置似乎不屬於 BRL49653。

關鍵詞: 過氧化體增殖活化受體，類黃素，發炎反應，水環化酶，一氧化氮合成酶

Abstract

PPAR γ transcription factor has been implicated in anti-inflammatory response. Of the compounds tested, apigenin, chrysin, and kaempferol significantly stimulated PPAR γ transcriptional activity in a transient reporter assay. In addition, these three flavonoids strongly enhanced the inhibition of inducible cyclooxygenase (COX-2) and nitric oxide synthase (iNOS) promoter activities in LPS-activated macrophages which containing the PPAR γ expression plasmids. However, these three flavonoids exhibited weak PPAR γ agonist activities in vitro competitive binding assay. Limited protease digestion of PPAR γ suggested these three flavonoids produced a conformational change in PPAR γ and the conformation differences in the receptor

bound to BRL49653 versus these three flavonoids. These results suggested that these three flavonoids might act as allosteric effectors and were able to bind to PPAR γ and activate it, but its binding site might be different from the natural ligand BRL49653.

Key words: Peroxisome proliferator-activated receptor- γ , Flavonoids, Inflammation, Cyclooxygenase, Nitric oxide synthase.

Introduction

Peroxisome proliferator-activated receptors (PPARs) are transcription factors belonging to nuclear receptor gene family [1]. PPARs bind to specific response elements as heterodimers with the retinoid X receptor (RXR) and activate transcription in response to a variety of endogenous and exogenous ligands, including certain polyunsaturated fatty acids, arachidonic acid metabolites [2], and some antidiabetic drugs [3] and non-steroidal anti-inflammatory drugs (NSAIDs)[4]. Currently, PPARs subfamily has been defined as PPAR α , PPAR β (also called PPAR δ and NUC1) and PPAR γ . Three PPAR isoforms differ in their tissue distribution and ligand specificity [5]. PPAR α is predominantly expressed in tissues exhibiting high catabolic rate of fatty acids (heart, liver, and kidney), whereas PPAR δ expression is ubiquitous, and its physiological role is unclear. PPAR γ is expressed predominantly in adipose tissue, the adrenal gland, spleen, large colon and the immune system [6-9]. Several lines of evidence indicated that PPAR γ plays an important role in regulating adipocyte differentiation and glucose homeostasis [10]. Both PPAR α and PPAR γ have been shown that also have anti-inflammatory actions through activating by arachidonic acid metabolites. PPAR α bind and be activated by leukotriene B4 [11], and the levels are induced at the transcriptional level by anti-inflammatory glucocorticoids [12]. PPAR γ are activated by the prostaglandin D₂ metabolite 15-deoxy- $\Delta^{12,14}$ prostaglandin J₂ (15d-PGJ₂) and synthetic antidiabetic thiazolidinedione drugs (e.g. BRL49653 and ciglitizone) and resulted in negatively regulating the expression of pro-inflammatory genes, and suppressing tumor cell growth [13-16]. Furthermore, both PPAR α and PPAR γ are activated by a number of non-steroidal anti-inflammatory drugs, such as indomethacin [4]. Recently, the PPAR γ agonists have been considered to inhibit production of monocyte inflammatory cytokines and the expression of iNOS [17, 18].

The flavonoids are a diverse family of chemicals commonly found in fruits and vegetables. Flavonoids are plant polyphenolic compounds, which comprise several classes including flavonols, flavanones, flavanols and flavans. Epidemiological studies have shown that the consumption of vegetable, fruits and tea is associated with a decreased risk of cancer and cardio-vascular diseases, and flavonoids are believed to play an important role in preventing these diseases [19]. Numerous numbers of this family have anticarcinogenic [20], anti-inflammatory [21], cytostatic [22], apoptotic [18], antioxidant [23], anti-angiogenic [24] and estrogenic [25] activities. Several reports have also shown that flavonoids are potent modulators of both the expression, and activities of specific cytochrome P450 genes/proteins [26]. These data indicate that certain flavonoids have attracted attention as possible chemoprotective or chemotherapeutic agents.

NSAIDs such as aspirin, sodium salicylate, and indomethacin exert their anti-inflammatory effects in part by inhibition of I κ B kinase- β (IKK- β), thereby preventing activation by NF- κ B of genes involved in the inflammatory response [27]. However, indomethacin and several other NSAIDs (fenoprofen, ibuprofen, and flufenamic acid) are also PPAR γ ligands and block production of inflammatory cytokines in human monocytes [17]. In addition, Ricote et al. [13] also demonstrated that treatment of peritoneal macrophages with 15d-PGJ₂ or several synthetic PPAR γ ligands reduce the expression of iNOS by interferon- γ and inhibited induction of gelatinase B and scavenger receptor A gene transcription in response to phorbol ester stimulation. Recently, we reported that apigenin and related flavonoids could suppress the transcriptional activity of COX-2 and iNOS in part through inhibition of I κ B kinase activity [28]. The current study was designed to determine whether the anti-inflammatory effects of flavonoids were correlative with their activation of PPAR γ .

2. Materials and Methods

2.1. Chemicals

LPS (*Escherichia coli* 0127:B8), flavone, 5-methoxyflavone, 7,8-dihydroxyflavone, apigenin, 3-hydroxyflavone, kaempferol, morin, quercetin, myricetin, rutin, genistein, indomethacin were purchased from Sigma Chemical Co. (St Louis, MO). Chrysin, luteolin, tangeretin, galangin, fisetin, pinocembrin, naringenin, isosakuranetin, eriodictyol, hesperetin, naringin, and biochanin A were purchased from Extrasynthese Inc. (Genay, France). Mouse interferon- γ was purchased from R & D systems Inc. (Minneapolis, MN). Two kinds of tea polyphenols, (-)-epigallocatechin-3-gallate and theaflavin-3,3'-digallate were purified as previously described [29].

2.2. Cell culture

The mouse macrophage cell lines RAW264.7 (ATCC, T1B71) were cultured as previously described [28]. Thioglycollate-elicited peritoneal macrophages were obtained from specific pathogen-free female Balb/c mice as previously described [30]. For all assays except the luciferase assay, cells were plated in 60 mm dishes at 5×10^6 cells/dish and allowed to grow for 18-24h. Treatment with vehicle (0.1% DMSO), test compounds and/or LPS or IFN- γ were carried out under serum-free conditions.

2.3. Determination of PGE₂ and nitrite

The cultured medium of control and treated cells were collected, centrifuged, and stored at -70°C, until tested. The level of PGE₂ released into culture media was quantified using a specific enzyme immunoassay (EIA) according to the manufacturer's instructions (Amersham). The nitrite concentration in the cultured medium was measured as an indicator of NO production according to the Griess reaction [28].

2.4. Plasmids

The PPAR γ expression plasmid and AOX-TK reporter plasmid were generously provided by Professor Christopher K. Glass (California University)[13].

The mouse iNOS promoter plasmid was generously provided by Professor Charles J. Lowenstein (Johns Hopkins University)[31]. The mouse COX-2 promoter plasmid containing a 1035 bp fragment, -966 to +70 relative to the transcription start, and constructed as previously described [28]. To generate the pGEX-2T-PPAR γ LBD chimeric receptor expression plasmid, cDNA encoding the ligand binding domains (LBD) of the mouse PPAR γ 1(amino acids 174-475) were amplified by polymerase chain reaction and subcloned into the pGEX-2T expression plasmid. Transient cotransfection and luciferase activity assay using these plasmids were performed as described previously [28].

2.5. Ligand binding assay

The Gst-PPAR γ LBD was expressed in JM109 *Escherichia coli* [32] and the fusion proteins were bound to the Glutathione Sepharose-4B beads according to the manufacturer's instructions (Pharmacia Biotech). For competition binding assay, 10 μ l of Glutathione Sepharose-4B beads containing 0.1 μ g of Gst-PPAR γ LBD chimeric protein were incubated with or without unlabeled flavonoids at 4 $^{\circ}$ C for 12h in buffer containing 10 mM Tris (pH 7.4), 50 mM KCl, 10 mM dithiothreitol and proteinase inhibitors, then added [3 H]BRL49653 (specific activity, 60 Ci/mmol) for additional 8 h. Bound [3 H]BRL49653 was precipitated from free radioactivity by centrifugation, and washed three times with PBS. The beads containing [3 H]BRL49653 were collected and quantitated by liquid scintillation counting.

2.6. Limited protease digestion assay

The protease digestion assays were performed by the method of Allen et al. [33], with some modification. The PPAR γ expression plasmid [13] was used to synthesize [35 S]-radiolabeled PPAR γ in a coupled transcription/translation system according to the protocol of the manufacturer (Promega). Approximately 5 μ l of the transcription/translation reactions was preincubated with 1 μ l of tested compounds for 20 min at 25 $^{\circ}$ C. Trypsin was added and allowed to proceed for 10 min at 25 $^{\circ}$ C, then terminated by the addition of SDS sample loading buffer and boiling for 8 min. The products of the digestion were separated by electrophoresis through a 12% SDS-polyacrylamide gel. Labeled PPAR γ was visualized by autoradiography.

3. Results

3.1. Apigenin, chrysin, and kaempferol activated PPAR α in macrophages

A series of flavonoids including flavone, 5-methoxyflavone, 7,8-dihydroxyflavone, apigenin, 3-hydroxyflavone, kaempferol, morin, quercetin, myricetin, rutin, genistein, chrysin, luteolin, tangeretin, galangin, fisetin, pinocembrin, naringenin, isosakuranetin, eriodictyol, hesperetin, naringin, and biochanin A, tea polyphenols (Table 1), and indomethacin were first tested with

regard to their activation effects on PPAR γ in RAW264.7 cells. As RAW264.7 cells express very low levels of PPAR γ [13] and required transfection of a PPAR γ expression plasmid. The cells therefore allowed a direct assessment of the role of PPAR γ in mediating the effects of these flavonoids on macrophage gene expression. The PPAR γ expression plasmid was cotransfected into RAW264.7 cells with a reporter construct containing three copies of the acyl CoA oxidase PPAR responsive element (PPRE) upstream of the thymidine kinase (TK) promoter driving luciferase gene expression. In the absence of a cotransfected PPAR γ expression plasmid, treatment of RAW264.7 macrophages with the tested flavonoids at 10 μ M had little effect on activation of PPAR γ . However, when a PPAR γ expression plasmid was cotransfected into the cells, apigenin, chrysin, and kaempferol significantly induced the PPAR γ 8.13, 5.60 and 7.66-fold, respectively ($p < 0.05$) (Fig. 1A). The positive control of indomethacin (100 μ M) strongly induced the PPAR γ activity 13.34-fold which was compared with the mock experiment. Apigenin, chrysin, and kaempferol increased the PPAR γ activity in a dose-dependent manner, with the EC₅₀ of approximately 5 μ M, 10 μ M, and 10 μ M, respectively (Fig. 1B). However, there was a cytotoxic effect in RAW264.7 cells with 20 μ M of apigenin and a decrease in the activation of PPAR γ . Flavone, 7,8-dihydroxyflavone, 3-hydroxyflavone, luteolin, galangin, genistein, biochanin A also increased the PPAR γ activities when RAW264.7 cells were transfected with the expression plasmid of PPAR γ . However, the induction folds of these flavonoids showed no significant difference in activation of PPAR γ compared with control RAW264.7 cells that were transfected with PPAR γ expression plasmid (Fig. 1A, land 2, 2.74-fold). The other tested flavonoids were unable to activated the PPAR γ , and the data were not shown in Fig. 1A.

3.2. Apigenin, chrysin and kaempferol enhanced the inhibition of COX-2 and iNOS promoters' activities in a PPAR α -dependent manner

RAW264.7 cells was transiently transfected with the reporter plasmids of COX-2 or iNOS and both promoters' activities were markedly increased when RAW264.7 cells were treated with LPS (Fig. 2). Both promoters' activities were inhibited by concurrent treatment of the cells with apigenin, chrysin and kaempferol. Moreover, transfection of PPAR γ expression plasmid enhanced the inhibitory effects of these three flavonoids (Fig. 2A&B). The results suggested that apigenin, chrysin and kaempferol inhibited the promoter' activities of COX-2 and iNOS genes partially through PPAR γ pathways.

3.3. Apigenin, chrysin, and kaempferol bound with PPAR α and induced conformational change in PPAR α

We next sought to determine whether these three flavonoids activated PPAR γ through direct interaction with the PPAR γ receptor. The abilities of these flavonoids to bind to PPAR γ were assessed in a competition binding assay using [³H] BRL49653 and the Glutathione Sepharose beads containing Gst-PPAR γ LBD fusion protein. As shown in Fig. 3, [³H] BRL49653 bound specifically

and saturably to Gst-PPAR γ LBD beads with a K_d of 8 nM (Fig. 3A & B). No binding was detected in control Gst-PPAR γ LBD beads (Data not shown). Apigenin, chrysin, and kaempferol competed with [³H] BRL49653 for binding to the PPAR γ LBD in a dose-dependent manner, with an IC₅₀ of approximately 50 μ M. Proteolytic analysis have been used for several experiments to demonstrate that ligands of nuclear receptor can, upon binding, specifically alter the conformation of the receptor [33, 34]. This conformational change was reflected by the increased resistance of the receptor to partial digestion by proteases. To determine if there were conformational differences in PPAR γ bound to these three flavonoids, a limited trypsin digestion on a [³⁵S] methionine-labeled PPAR γ was performed. As shown in Fig. 4, incubation of PPAR γ with increasing concentrations of trypsin in the absence of ligand led to the complete digestion of PPAR γ . In contrast, BRL49653 induced a stronger protection of the 22-, 29- and 30-kDa fragments. These three flavonoids binding yielded 29- and 30-kDa protected fragments, especially 30-kDa band. These results indicated that these three flavonoids were able to bind to PPAR γ and flavonoids-bound PPAR γ had a distinct trypsin digestion pattern compared with the BRL49653-bound receptor.

4. Discussion

Flavonoids are naturally occurring plant polyphenols found in abundance in the diets rich in fruit, vegetables and plant-derived beverages such as tea. The PPAR γ ligands share certain structural characteristics including a lipophilic backbone and an acid moiety, usually a carboxylate. Although flavonoids only have a similar lipophilic backbone, several flavonoids also bind to PPAR γ in vitro (Fig. 4). Several reports have shown that treatment of various fibroblast and mesenchymal stem cells lines with PPAR γ ligands, including 15d-PGJ₂, the anti-diabetic drugs, and several NSAIDs, promotes their efficient conversion to adipocytes [3, 4, 35-37]. However, treatment of C3H10T1/2 stem cells with 10 μ M of apigenin, chrysin, and kaempferol failed to promote adipocytes differentiation as indicated by Oil Red O staining (Data not shown). Northern blot analysis indicated that PPAR γ expression levels were not induced in response to treatment with these three flavonoids in fibroblast cells (Data not shown). As we known, adipocyte differentiation required forced expression of PPAR γ and was significantly enhanced in the presence of PPAR γ activators [38]. For example, treatment of C3H10T1/2 cells with BRL49653 increased PPAR γ expression levels approximately 3-fold, and resulted in efficient adipocyte differentiation [3]. In addition, while flavonoids were potent inhibitors of several kinases involved in signal transduction, mainly protein kinase C [39] and tyrosine kinase [40, 41]. However, adipocyte differentiation is characterized by a coordinate increase in adipocyte-specific gene expression through activating of gene transcription. Therefore, they were insufficient to initiate the adipogenic signaling cascade in a mesenchymal stem cell line. Devchand et al. [11] proposed that leukotriene B₄, an agonist of the related receptor PPAR α , has anti-inflammatory activity. The flavonoids might also activate the PPAR α and have anti-inflammatory effects, but PPAR α is not expressed in activated macrophages.

Of the tested flavonoids (Table 1), the groups of flavanones and flavan-3-ol were inefficient on

activation of PPAR γ . This result suggested that the C2-C3 double bond of C ring was essential for their activation of PPAR γ . Some flavonoids of flavones, flavonols and isoflavones groups were able to activate PPAR γ . The activation of PPAR γ seems to be dependent on the number and the position of hydroxyl residues. The hydroxyl residues of the 5 and 7 positions of A ring and 4' position of B ring were important factors for activation of PPAR γ , such as apigenin, chrysin and kaempferol. However, additional hydroxyl residue of 3' position of B ring resulted in decrease the activation of PPAR γ , such as luteolin and quercetin.

These three flavonoids were able to activate PPAR γ in a transient reporter assay, with an EC₅₀ of approximately 5~10 μ M. However, they needed a higher concentration to bind to Gst-PPAR γ in vitro competitive binding assay (Fig. 3). The high concentration of IC₅₀ (50 μ M) suggested that these three flavonoids might not directly bind to PPAR γ or bind to PPAR γ in the other sites. In limited protease digestion assay (Fig. 4) indicated that PPAR γ has a conformational difference in the receptor bound to the three flavonoids versus natural ligand BRL49653. These results suggested that these three flavonoids might act as allosteric effectors, and were able to bind to PPAR γ and activate it, but its binding site might be different from the natural ligand BRL49653. Based on the different binding kinetics, it may interpret the fact that these three flavonoids did not promote differentiation of C3H10T1/2 stem cells to adipocyte. Our previous studies have shown that apigenin was able to inhibit I κ B kinase activity and prevent the activation of NF- κ B, and then suppress the promoter activities of COX-2 and iNOS [28]. Therefore, apigenin was a more stronger inhibitor of COX-2 and iNOS promoters activities than BRL49653 in the absence and presence of transfected PPAR γ expression plasmid (Fig. 2). However apigenin was more effective in the presence of PPAR γ than absence of PPAR γ in inhibition of COX-2 and iNOS promoters. These results indicated that these three flavonoids inhibited the expression of COX-2 and iNOS partially through activating PPAR γ . In RAW264.7 cells, the base level of PPAR γ protein was very low, so BRL49653 could not activate it and then to suppress the activities of COX-2 and iNOS promoters in the absence of transfected PPAR γ expression plasmid.

Our results suggested that certain flavonoids could activate PPAR γ , then to inhibit the protein expression of COX-2. Indomethacin was able to inhibit COX activity without affecting the protein levels of PPARs at lower concentration [42]. At higher concentration, we thought that anti-inflammatory activity of indomethacin might be also mediated through activation of PPAR γ then inhibition of COX expression. Since indomethacin also acts as a PPAR γ agonist, promoting adipocyte differentiation.

Our previous studies demonstrated that the anti-inflammatory properties of apigenin might be mediated through inhibition of I κ B kinase activity. In this study, we showed that apigenin was also an efficacious activator of PPAR γ which regulated inflammatory responses. These results suggested that apigenin, chrysin, and kaempferol were possible activators of PPAR γ , and might have therapeutic applications in inflammatory diseases, such as atherosclerosis and rheumatoid arthritis. These findings also provide a significant molecular basis for explaining how dietary flavonoids are







active in preventing cancer and inflammation.

5. References

- [1]. Schoonjans, K., Martin, G., Staels, B. and Auwerx, J. (1997) *Curr. Opin. Lipidology* 8, 159-166.
- [2]. Keller, H., Preyer, C., Medin, J., Mahfoudi, A., Ozato, K. and Wahli, W. (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90, 2160-2164.
- [3]. Lehmann, J. M., Moore, L. B., Smith-Oliver, T. A., Wilkison, W. O., Willson, T. M. and Kliewer, S. A. (1995) *J. Biol. Chem.* 270, 12953-12956.
- [4]. Lehmann, J. M., Lenhard, J. M., Oliver, B. B., Ringold, G. M. and Kliewer, S. A. (1997) *J. Biol. Chem.* 272, 3406-3410.
- [5]. Forman, B. M., Chen, J. and Evans, R. M. (1996) *Ann. NY Acad. Sci.* 804, 266-275.
- [6]. Kliewer, S. A., Forman, B. M., Blumberg, B., Ong, E. S., Borgmeyer, U., Mangelsdorf, D. J., Umesono, K. and Evans, R. M. (1994) *Proc. Natl. Acad. Sci. U.S.A.* 91, 7355-7359.
- [7]. Tontonoz, P., Hu, E., Graves, R. A., Budavari, A. I. and Spiegelman, B. M. (1994) *Genes Deve.* 8, 1224-1234.
- [8]. Chawla, A., Schwarz, E. J., Dimaculangan, D. D. and Lazar, M. A. (1994) *Endocrinology* 35, 798-800.
- [9]. Fajas, L., Auboeuf, D., Raspe, E., Schoonjans, K., Lefebvre, A. M., Saladin, R., Najib, J., Laville, M., Fruchart, J. C., Deeb, S., Vidal-Puig, A., Flier, J., Briggs, M. R., Staels, B., Vidal, H. and Auwerx, J. (1997) *J. Bio. Chem.* 272, 18779-18789.
- [10]. Spiegelman, B. M. (1998) *Diabetes* 47, 507-514.
- [11]. Devchand, P. R., Keller, H., Peters, J. M., Vazquez, M., Gonzalez, F. J. and Wahli, W. (1996) *Nature* 384, 39-43.
- [12]. Lemberger, T., Saladin, R., Vazquez, M., Assimacopoulos, F., Staels, B., Desvergne, B., Wahli, W. and Auwerx, J. (1996) *J. Biol. Chem.* 271, 1764-1769.
- [13]. Ricote, M., Li, A. C., Willson, T. M., Kelly, C. J. and Glass, C. K. (1998) *Nature* 391, 79-82.
- [14]. Elstner, E., Muller, C., Koshizuka, K., Williamson, E. A., Park, D., Asou, H., Shintaku, P., Said, J. W., Heber, D. and Koeffler, H. P. (1998) *Pro. Natl. Acad. Sci. U.S.A.* 95, 8806-8811.
- [15]. Kubota, T., Koshizuka, K., Williamson, E. A., Asou, H., Said, J. W., Holden, S., and Miyoshi, I. and Koeffler, H. P. (1998) *Cancer Res.* 58, 3344-3452.
- [16]. Sarraf, P., Mueller, E., Jones, D., King, F. J., DeAngelo, D. J., Partridge, J. B., Holden, S. A., Chen, L. B., Singer, S., Fletcher, C. and Spiegelman, B. M. (1998) *Nature Med.* 4, 1046-1052.
- [17]. Jiang, C., Ting, A. T. and Seed, B. (1998) *Nature* 391, 82-86.
- [18]. Konig, A., Schwartz, G. K., Mohammad, R. M., Al-Katib, A., and Gabilove, J. L. (1997) *Blood* 90, 4307-4312.
- [19]. Miller, A. B. (1990) *Rev. Oncol.* 3, 87-96.
- [20]. Middleton, E. Jr. and Kandaswami, C. (1986) In Harborne, J.H. and Liss, A.R. (ed.) *The flavonoids: Advances in Research Since 1969*. New York, pp. 619-652.

- [21]. Ferrandiz, M. L. and Alcaraz, M. J. (1991) *Agents Actions* 32, 283-288.
- [22]. Plaumann, B., Fritsche, M., Rimpler, H., Brandner, G. and Hess, R. D. (1996) *Oncogene* 13, 1605-1614.
- [23]. Cao, G., Sofic, E. and Prior, R. (1997) *Free Rad. Bio. Med.* 22, 749-760.
- [24]. Fotsis, T., Pepper, M. S., Aktas, E., Breit, S., Rasku, S., Adlercreutz, H., Wahala, K., Montesano, R. and Schweigerer, L. (1997) *Cancer Res.* 57, 2916-2921.
- [25]. Breinholt, V. and Larson, J. C. (1998) *Chem. Res. Toxicol.* 11, 622-629.
- [26]. Lu, Y. F., Santostefano, M., Cunningham, B. D. M., Threadgill, M. D. and Safe, S. (1995) *Arch. Biochem. Biophys.* 316, 470-477.
- [27]. Yin, M. J., Yamamoto, Y. and Gaynor, R. B. (1998) *Nature* 396, 77-80.
- [28]. Liang, Y. C., Huang, Y. T., Tsai, S. H., Lin-Shian, S. Y., Chen, C. F. and Lin, J. K. (1999) *Carcinogenesis* 20, 1945-1952.
- [29]. Liang, Y. C., Chen, Y. C., Lin, Y. L., Lin-Shiau, S. Y., Ho, C. T. and Lin, J. K. (1999) *Carcinogenesis* 20, 733-736.
- [30]. Lin, Y. L. and Lin, J. K. (1997) *Mol. pharmacol.* 52, 465-472.
- [31]. Lowenestein, C. J., Alley, E. W., Raval, P., Snowman, A. M., Snyder, S. H., Russell, S. W. and Murphy, W. J. (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90, 9730-9734.
- [32]. Liang, Y. C., Lin-Shiau, S. Y., Chen, C. F. and Lin, J. K. (1999) *J. Cell. Biochem.* 75, 1-12.
- [33]. Allen, G. F., Xiaohua, L., Tsai, S. Y., Weigel, N. L., Edwards, D. P., Tsai, M.-J. and O'Malley, B. W. (1992) *J. Biol. Chem.* 267, 19513-19520.
- [34]. McDonnell, D. P., Clemm, D. L., Hermann, T., Goldman, M. E. and Pike, J. W. (1995) *Mol. Endocrinol.* 9, 659-669.
- [35]. Kletzien, R. F., Clarke, S. D. and Ulrich, R. G. (1992) *Mol. Pharmacol.* 41, 393-398.
- [36]. Sparks, R. L., Strauss, E. E., Zygmunt, A. I. And Phelan, T. E. (1991) *J. Cell. Physiol.* 146, 101-109.
- [37]. Forman, B. M., Tontonoz, P., Cen, J., Brun, R. P., Spiegelman, B. M. and Evans, R. M. (1995) *Cell* 83, 803-812.
- [38]. Tontonoz, E. Hu. and B. M. (1994) *Cell* 79, 1147-1156.
- [39]. Ferriola, P. C., Cody, V. and Middleton, E. (1989) *Biochem. Pharmacol.* 38, 1617-1624.
- [40]. Cushman, M., Nagarathman, D., Burg, D. L. and Geahlen G. L. (1991) *J. Med. Chem.* 34, 798-806.
- [41]. Agullo, G. and Gamet-Payrastre, B. (1997) *Biochem. Pharmacol.* 53, 1649-1657.
- [42]. Vane, J. R. (1971) *Natl. New. Biol.* 231, 232-235.

Table 1
Chemical structures of the various flavonoids tested for the induction of PPAR γ activity in RAW246.7 cells

Chemical formula	Name	Substitution							
		5	6	7	8	2'	3'	4'	5'
Flavones									
	Flavone	H	H	H	H	H	H	H	H
	5-Methoxyflavone	OCH ₃	H	H	H	H	H	H	H
	7,8-Dihydroxyflavone	H	H	OH	OH	H	H	H	H
	Chrysin	OH	H	OH	H	H	H	H	H
	Apigenin	OH	H	OH	H	H	H	OH	H
	Luteolin	OH	H	OH	H	H	OH	OH	H
	Tangeretin	OCH ₃	OCH ₃	OCH ₃	OCH ₃	H	H	OCH ₃	H
Flavonols									
	3-Hydroxyflavone	H	H	H	H	H	H	H	H
	Galangin	OH	H	OH	H	H	H	H	H
	Kaempferol	OH	H	OH	H	H	H	OH	H
	Fisetin	H	H	OH	H	H	OH	OH	H
	Morin	OH	H	OH	H	OH	H	OH	H
	Quercetin	OH	H	OH	H	H	OH	OH	H
	Myricetin	OH	H	OH	H	H	OH	OH	OH
Flavonol glycoside									
	Rutin	OH	H	OH	H	H	OH	OH	H
(3: OR; R: Rhamnosylglucoside)									
Flavanones									
	Pinocembrin	OH	H	OH	H	H	H	H	H
	Naringenin	OH	H	OH	H	H	H	OH	H
	Isosakuranetin	OH	H	OH	H	H	H	OCH ₃	H
	Eriodictyol	OH	H	OH	H	H	OH	OH	H
	Hesperetin	OH	H	OH	H	H	OH	OCH ₃	H
Flavanone glycoside									
	Naringin	OH	H	OR	H	H	H	OH	H
(R: Rhamnosylglucoside)									
Isoflavones									
	Genistein	OH	H	OH	H	H	H	OH	H
	Biochanin A	OH	H	OH	H	H	H	OCH ₃	H
Flavan-3-ol									
	(-)-Epigallocatechin-3-gallate								
Theaflavins									
	Theaflavin-3,3'-digallate (R=Galloyl)								




Fig. 1. Effects of apigenin, chrysin, and kaempferol on the activation of PPAR γ in cells. RAW264.7 cells were cotransfected with the AOX-TK reporter plasmid and PPAR γ expression plasmid (open squares) or PPAR γ mock plasmid (closed squares) and β -galactosidase expression vector (pcDNA lacZ) as internal control. A: Transfected cells were treated with 10 μ M of flavonoids, or 100 μ M of indomethacin for 18 h, and cell extracts subsequently assayed for luciferase and β -galactosidase activities as described in Section 2. The values were represented as means \pm S.E.M. of triplicate tests. The numbers were obtained from ratios of the induction fold of transfected PPAR γ expression to mock expression and indicated in the figure. * $P < 0.05$ (Student's t -test) versus control (2.74). B: Transfected cells were treated with various concentrations of apigenin (closed circles), chrysin (open circles), and kaempferol (closed triangles) for 18 h, and cells extracts assayed for luciferase and β -galactosidase activities as described above. Con., control; Indo, indomethacin; Api, apigenin; Chr, chrysin; DHF, 7,8-dihydroxyflavone; Fla, flavone; HFl, 3-hydroxyflavone; Lut, luteolin; Gal, galangin; Kae, kaempferol; Bio, biochanin A; Gen, genistein.




Fig. 2. Apigenin, chrysin and kaempferol enhanced the inhibition of the promoter activities of COX-2 and iNOS in a PPAR γ -dependent manner. RAW264.7 cells were cotransfected with a COX-2 (A) or iNOS (B) promoter-luciferase reporter plasmid, pcDNA lacZ internal control vector, and either a PPAR γ expression plasmid or mock plasmid as indicated. Cells were treated with LPS (50 ng/ml) and BRL49653 (20 μ M), or various flavonoids (10 μ M) for 18 h, and extracts subsequently assayed for luciferase and β -galactosidase activities as described in Section 2. The values were represented as means \pm S.E.M. of triplicate tests. * $P < 0.05$, compared with the individual test of mock expression (Student's t -test).





Fig. 3. The binding capacity of apigenin, chrysin, and kaempferol to PPAR γ in vitro. A: Glutathione Sepharose beads containing the Gst-PPAR γ LBD fusion protein were incubated with increasing concentrations of [3 H]BRL49653. Non-specific binding was determined using 500-fold excess of unlabeled BRL49653. Closed triangles, total binding; closed circles, non-specific binding; open circles, specific binding. B: Scatchard analysis of specific binding data from A. Linear regression yielded a K_d of 8 nM. Similar results were obtained in three independent experiments performed in duplicate. C: Competition binding analysis was performed with Sepharose-GST-PPAR γ LBD beads and 5 nM [3 H]BRL49653 in the presence of various concentrations of apigenin (closed circles), chrysin (open triangles), or kaempferol (closed squares) as unlabeled competitors. Similar results were obtained in two independent experiments performed in duplicate.




Fig. 4. Apigenin, chrysin and kaempferol induced a protease-resistant conformation of PPAR γ . [35 S]PPAR γ was synthesized in vitro in a coupled transcription/translation system. The receptor was preincubated with ethanol (Veh), apigenin (Api), chrysin (Chr) or kaempferol (Kae), then added to dH $_2$ O or increasing concentration of trypsin. Digestion products were analyzed by SDS-PAGE followed by autoradiography. Asterisks and arrows indicate the full-length PPAR γ and trypsin-resistant fragment of PPAR γ , respectively.