ƒ ‹ F | Œfia‹ ·'e›ß• |–M^D‹ ª¤s›p e ƒ¤"G†ŁSi

‰w¿E朴／ oiNOS " †{／¸cytokines ~ 'æ／§ T, „¶ ·»…, fi|／¸

'..ƒ„¶¡／S‰ -- ¤ -• Q

›p e^ §¡G ª ¥«×p e

›p e‰s, „¡GNSC91-2314-B-038-010-

ɹf ·`¶¡¡G91 ƒ~08／°01／Ø 92 ƒ~07／°31／Ø

ɹf † ƒ ¡G» O ¥´ ·／j ·´ ¤ § †Mt

›p e¥ D ø／ĦG" L «•

†ŁSi^ «‹¡G" °††ŁSi

† B † zŁ„fG¡¥»›p eflA／¸-M§ Q ' ¸ ¤ ¥ L · …z2 ƒ†‹€ Æv¥Ai ／ ¶}‹d,

／／ ¥` Œ 92 ƒ~10／°30／Ø

ƒ ‹ F | Œ fi a ‹ ' e › ß • ┼M^D‹ª¤s›p eƒ¤"G†ł§i

† ˜ ⁄ T fl « , g ¤ t † ¸ „ "⁄—'I§l„D⁄§‰ --
‰ w ¿ E ®⁄iN©S " †{⁄¸cytokines ˜ 'æ⁄§ T fi § ¶ »…⁄¸fi|
' … ƒ ¶¡‰"– ¤ –· Q

Bradykinin-induced iNOS expression and cytokines release in human airway epithelial cells: the interaction mechanisms and signal transduction pathways

› pe‰ s „ ¡№SC 91-2314-B-038-010
ɪƒ · ››¡©91 ƒ~8 ⁄°1 ⁄ Ø 92 ƒ~7 ⁄°31 ⁄Ø
¥ D ⁄¡HG" L «• ɪƒ "⁄¸† ƒ ƒW¡"G¥ x ¥´ ·⁄j ·´ ¤ § †№t

⁄@¡B⁄⁄⁄ "K›n

¥»› pe¥D›nƒb– ·Q bradykinin ¤°¿E⁄H ^ "˝– ƒ ⁄ W ¥ † A549¡¸^cyclooxygenase-2 (COX-2) " †{" "T„¶ · » „. ƒi┤ƒb† ˜⁄@ƒ~ ⁄ † ˜⁄Gƒ~"˝‹ ª§⁄ o†{¡AƒbA549 † ›M⁄¡A bradykinin ¥i,g¥ ‹⁄˘p44/42 MAPK ⁄¸ p38 MAPK " " ,fi┤ oNF-κB "˝‹ ⁄˘¡A† « Æ ›£COX-2 "˝" †{¡CƒbƒbT ƒ~"˝‹ ª§⁄ § › №§⁄¶‡B– ·Q bradykinin ›P COX-2 " †{" " ,fi┤⁄¡Ap44/42 MAPK ⁄¸ p38 MAPK »P NF-κB ⁄§¶ ¡ " " T §¶·» fi┤¡A¥H⁄¸¤ ¥L⁄ _ p44/42 MAPK ⁄¸ p38 MAPK ‹ ⁄˘" " T fi § ¶»¡CRa§i┤§ ¤ fl manumycin A (0.3, 1 'M3 μM) ⁄¸Raf § ¤ flGW 5074 (0.3, 1 'M3 nM) ¥H f¶ q ‹ ^"˝ ⁄ ¡§ ¤ bradykinin⁄ o⁄§COX-2 " †{¡C Manumycin A (3 μM) »PGW 5074 (3 μM) ⁄]¥i¥H§ ¤ bradykinin⁄ op44/42 MAPK "˝‹ ⁄˘¡Aƒ ª⁄£┤… v̄T p38 MAPK "˝‹ ¡ ⁄˘¡CBradykinin ¥Hfi ¶¡‹ ^ "˝⁄ ¡¤ ˝o IKKα/β ‹ ¡ ' ° … W ¥ [A•PD¥ 98059 (30 μM)¡BSB 203580 (1 μM)'Mmanumycin A (3 μM) «Æ¡A ф{ PD 98059 »Pmanumycin A ¥ i § „ ¥§ bradykinin⁄ o IKKα/β " ˝¡ ⁄˘¡A¸ƒSB 203580 ¥ u • ┤ † ⁄˘ ¡§KKα/β "˝ ‹ ¡ ⁄˘¡Bradykinin ¤ ¿E A549 † › M £F«Pof˜Ras/Raf/p44/42MAPK and p38 MAPK is upstream of bradykinin-induced IKKα/β activation. IκB-α"˝⁄ , ¶i¤ƒ˜p65 'Mp50 NF-κB ¶i⁄J † ›Mfi ⁄ § ¥ ~ ¡ A ⁄] ┼¥"Hfi ¶¡ƒ ⁄˘ o

IκB-α »P p65 ` ß˜⁄ ¡C" ƒX¥H⁄ W †"G – ϑ ¡AƒbA549 † ›M⁄¡Abradykinin , ¥ ‹ ¡ ⁄˘Ras/Raf/p44/42 MAPK ⁄¸ p38 MAPK "˝ , fi┤¡A¤ ⁄ oIKKα/β " " ‹⁄¡A¤˘IκB-α `C »˜⁄˘»P⁄ , ƒ ¶i@B‹ ⁄˘ NF-κB ¡C £F bradykinin⁄ oIκB-α ` ß˜⁄˘⁄ § ¥ ~ A┼H⁄y• ┤ ƒ¤p65 ` ß˜⁄ ¡C

^ ` ¡ ©w¿E¤ O ¡ B fiæ2┤¡B p44/42MAPK¡Bp38MAPK¡B⁄W¥ † ›M

Abstract

The signal transduction pathway of bradykinin-induced COX-2 expression was studies in human pulmonary epithelial cell line (A549). The Ras inhibitor (manumycin A) and Raf inhibitor (GW 5074) concentration-dependently attenuated the bradykinin-induced COX-2 expression. The bradykinin-induced p44/42 MAPK but not p38 MAPK activation was inhibited by manumycin A and GW 5074. Bradykinin caused time-dependent increases in IKKα/β, IκB-α and p65 phosphorylation, IKKα/β activity in A549 cells. The IKKα/β activation was almost completely inhibited by manumycin A and PD 98059. However, SB 203580 partially inhibited IKKα/β activation. Taken together, these results suggest that bradykinin might activate IκB-α and p65 phosphorylation and IKKα/β activation in human airway epithelial cell line. The signal of Ras/Raf/p44/42MAPK and p38 MAPK is upstream of bradykinin-induced IKKα/β activation.

Keywords: Bradykinin, Cyclooxygenase-2, p44/42 MAPK, p38 MAPK, A549 cells.

⁄G¡B‰t¥ »P¥ "˝

• † ·¤' ¶¸ƒ ┼ o"˝Ψ˝¤° ° ¿ E fi‹ •┤ ⁄ o ƒ … '¸kallikrein "˝‹ ¡ ⁄˘ A┼¶i⁄@

¤ B Æ–Nkininogen ⁄ , ƒ ¤ ¤ ª ƒ ¥˝ ”«d ¡¡'Â⁄¥ < ª A I $„Ð ”i⁄O¡ B † ›M ‴W¥˝¡ ¥ kinins[1] ¡C Bradykinin ‹ O @ » ¤ ƒ+⁄ E › i … ”˝˝ ¥X¡ B ´ ¥G”˝ e¸¥ •P»P ˜ ¥•fP« »˜⁄$kinin¡ A ¥ « e‹w ƒ¤« ƒh¥˝† z ¸flf , g”˝‹ ¦ °¡Cƒ L‰ƒpƒ ¡A†o¤ ·” ª¡' †£¥˝ † ²˝' °‰ Ł ¡ A ¥] ‹ A ⁄ _¥ ×•˜ƒY ”¥ƒ ”⁄S«eƒ Ç¢fl ¡ Æ ' $ Œt⁄S¥\fl ¥D›n‹O¤ ¿ ” ” ´ X – i ¡ B … W ¥ƒ' ”¡ B”˝+¥G²˝ ¢B ¥ ƒ ƒbS ‰ † ¡˜”˝¿ @ ”” $C¡B»P‹O$_ƒ † … W ¥ [” ⁄j †hista¥mine ”˝˜ 'æ¡B« ¶i ¤ fl $$”˝¤ ¯Øf ¥b¥H⁄ ¤ ' ‹ ¡⁄˜”˝ T„¶· neutrophil 'M eosinophil ” ‴W¥˝⁄» ⁄ fik h» … ,fi⁄S⁄ £P¡Aƒ]ƒ †£¥˝⁄ £P”˝¥\fl ¡C ”˝†£¥˝[1] ¡C Bradykinin ¥ ¥ i ¤ ° Ł«Ð˚k˜P † ”æ”„ Œ † $i¥ « ¥X bradykinin ¥i , g ¤†”¸˝¶ · ⁄ J ¤,⁄gƒ _ neuropeptide ”˝ ¤ ¿E eicosanoid ”˝ƒX¤⁄¸ cytokines ”˝†£ ˜ 'æ¡Aƒ]ƒ ⁄ ofl ¥g'° o” ¢ › ¡C• M„L ¥˝¡Cƒb⁄ ”„¥˝” ª¡† ´·¡Abradykinin ¥ ¤° – › ¤ ¿Efi † flf⁄ fi ¡A¥i o†{¤ ' I $„Ð$ ¿E IL-1β, IL-2, IL-6 ”˝” †{[9] ¡Fƒb⁄H^ ”˝– kinin ”˝¿ @ ƒ+– ª ” ” W¥ Ł B2 ¤ – ¥œ$ ”ß¥ † ›M bradykinin ¥ i ¤¿E IL-8 ”˝†£¥˝ fl¥if ¤fi ˜ $¤„ L –›' » ⁄o late phase ⁄$ [10] ¡Fƒb⁄H^ ”˝fi ” ¥ › • ƒ bradykinin ¥i o” ¢ › ²[2] ¡C¥ ¥~¡Abradykinin ‹ O @ ×« ›« ¤ ¿E COX-2 ”˝” †{'M PGE2 ”˝˜ 'æ[11] ¡C¥t ›n”˝» o”¢(pro-inflammatory)⁄¶‰ Ł Æ¤¥ ¡g¥ ¥~ƒb⁄H^ ”˝– ”ß¥ † M¡¥Æi o†{PGE2 ¥i…W … ¥[•Lƒ ”ƒ… ”˝” ¥X¡A¶if › Ł ”˝‹ –j IL-1β' ⁄ o⁄$ IL-6 'M IL-8 ”˝†£¥˝¡Cƒ fl Ł~¡Aƒ ⁄ o⁄ fi ” ”˝„L« ⁄ ˇ ª(bronchial bradykinin ¤ ª”‹O$_• |»⁄ ' I $„ÐW¥ † hyperreactivity)[1] ¡C ›M' ¸COX-2 ” †{¡ A ¸¤« O, g¥ ƒ ” Tfi$¶· ` C »^ƒA2 (phospholipase A2, PLA2) »...–N T„¥ † ›M‰⁄¶·ƒ † ›Mfi ¤ »⁄ †o¤· ‹O¤ ›¯Ø†£¥˝⁄@¤ fl ‰Ł^ o”¢⁄¶‰ Ł”⁄ @ ›« o”¢' °ª ¥ ‰ Ł”˝ †{¡A«hƒ ⁄ ' ¥¸..ƒª$ „ª ›n”˝» fl ¡ Æ ¥D›n”˝$ ¥ ¸O–N† ›M‰ ⁄W”˝ ”˝ª $i¡Cƒ ' I $„ÐW¥ † ›M‹O› «n”˝ o”¢ ` C ›‰Ł¥ Ñ´ƒ¤”Æ¥˝´¥ ┼m»˜¡Aƒ fiæ» fl † ›M ÆA»P' I $„Ð” o” ¢ › ª†› «n”˝^ (cyclo-oxygenase, COX)ƒA¶i⁄@¤ BÑ”Æ¥˝´¥| ª†s' °¡Cƒ]ƒ„¡A¥ »ª¤s› pe–N¤˜¥¸' I $„ÐW †m»˜ (arachidonic acid) ¥ Ñ´ƒ¤«eƒ Ç¢fl ¥ † M¡ºA–·¤s bradykinin ⁄ o COX-2 ” (prostaglandin) ¡ A ¥‹ A prostaglandin E2 †{”˝ T„¶· » … ,fi¡C (PGE2), prostacyclin (PGI2) 'M thromboxane A2 (TxA2)¡C¥ « e‹w”˝ C »^fl ⁄T¡B †”G»P Q‰ A2 ¤ ƒ+ƒh˝ £ƒP˝« ƒ¡¡A»P o” ¢ › ,ª3ƒ+ ƒbƒ ˜⁄@ƒ~⁄¸†˜⁄ Œ~”˝‹ ª¤s⁄⁄ o†{¡Afb ^ «Y”˝‹O⁄ ⁄ ¶q14 kDa ⁄$ secretory PLA2 A549 † ›M⁄¡ bradykinin ¥ ¡g¥ ‹ ¡⁄˜p44/42 (sPLA2) 'M 85 kDa ⁄$ cytosolic PLA2 MAPK ⁄¸ p38 MAPK ”˝, fi|⁄ o NF-κB (cPLA2)¡Cƒ fiæ» fl «hƒ+¤ ˝ £ƒP˝ƒP¥\ ”˝‹ ¡⁄˜¡A+ «Æ ›P COX-2 ”˝” †{¡C¢– » fl (isoforms) ⁄ $O‹O COX-1 ⁄¸ radykinin ¥ ¡g¥ ‹ ¡⁄˜B2 ¤ ¯Ø¤ ⁄ op44/42 COX-2[3] ¡C COX-1 ” ‹ ' Twƒ«sƒb MAPK ⁄¸ p38 MAPK ‹ ¡⁄˜¡Cƒb†˜⁄Tƒ~”˝‹ ª (constitutive form) ¤ ƒ $b« ƒh˝ ” ” † ›M¡A¤s⁄⁄$ › –N$ ¶i⁄@¤B–· Q bradykinin ›P ¥ ¡A⁄ ¥ † ›M¸ƒ P˝O¡A¥ ƒ ' †£¥ ˝$«e COX-2 ” ‹ »⁄ ««‹ (inducible form)¡A• ¤ ƒ Ç¢fl ¥D›n$ Œ ” ˝ß†«ø»M« 'w”˝¥\fl [3] ¡C ¤ ⁄@¤· o”¢' ª«‰Łƒp lipopolysaccharide (LPS) '¸ cytokines ƒp IL-1β, TNF-α ¥”˝ ¤ ¿E• |»⁄ ¤ ” † {[4,5] ¡C¶i⁄@¤B”˝„ Œ ¯⁄] ^ „ŒCOX-2 ¥D›n‹O› ¢d†£¥˝⁄@¤·, o”¢ƒ+ ^ ” «eƒ Ç¢fl [3] ¡Cƒ ¥B⁄£˝ ƒb¥ ¸ ”˝¥˝†z‹ ˝A' ¸ƒbfl f ²”˝ ‹ ª⁄µª” ª¡”˝† ´·$¡• |˜ 'æ¥X« ƒ ª˝ƒ«eÇ¢fl ¤ ‰,” I $„Ð˝¥\

1. Ras ⁄¸ Raf ¥]$tƒb bradykinin ¤°¿E COX-2 ” †{”˝ T, „¶· »..., fi|

• „ ¥ ¥Ł$ ¤ fmanunycin A (0.3, 1 'M 3 μM) ⁄¸ Raf $ ¤ fGW 5074 (0.3, 1 'M 3 μM) « e ª†z30 ⁄ ˜ ` « Æ ªƒA¥[⁄J

bradykinin (10^{-8} M) ...
manunycin A ... GW 5074 ...
... COX-2 ... Ras ... Raf ... bradykinin ... COX-2 ...

2. Ras ... Raf ... bradykinin ... p44/42 MAPK ...

... manunycin A (3 µM) ... GW 5074 (3 µM) ... bradykinin (10^{-8} M) ... manunycin A ... GW 5074 ... bradykinin ... p44/42 MAPK ... p38MAPK ... bradykinin ... Ras ... Raf ... p44/42 MAPK ...

3. Bradykinin ... A549 ... IκB-α ... p65 ...

... anti-phospho-IκBα ... anti-phospho-p65 ... Western blot ... IκBα ... p65 ... bradykinin (10^{-8} M) ... A549 ... 5, 10, 20, 30 ... IκBα ... 20 ... p65 ... 10 ... A549 ... bradykinin ... IκBα ... p65 ...

4. Bradykinin ... IKKα/β ...

... anti-phospho-IKKα/β ... Western blot ... IKKα/β ... immunopreciptation ... IKKα/β ... anti-phospho-IκBα ... Western blot ... IKKα/β ... bradykinin (10^{-8} M) ... A549 ... 3, 5, 10, 20, 30 ... IKKα/β ... 5 ... 10 ...

5. Ras ... p44/42 ... p38 MAPK ... bradykinin ... IKKα/β ...

... manunycin A (3 µM) ... (30 µM) ... SB 203580 (1µM) ... bradykinin (10^{-8} M) ... PD 98059 ... manunycin A ... SB 203580 ... bradykinin ... IKKα/β ... Ras ... p44/42 MAPK ... p38 MAPK ... bradykinin ... IκB-α ... NF-κB ... bradykinin ... IκB-α ... p65 ...

...

... Bradykinin ... COX-2 ... bradykinin ... p44/42 ... p38 MAPK ... Ras/Raf/p44/42 MAPK ... p38 MAPK ... IKKα/β ... IκB-α ... NF-κB ... bradykinin ... IκB-α ... p65 ...

...

1.Proud, D. and Kaplan A.P. (1988) Kinin formation: mechanisms and role in inflammatory disorders Annu. Rev. Immunol. 6:49-83.

2.Abraham, W.M. (1992) The potential role of bradykinin antagonists in the treatment

of asthma. Agents Actions 38: 439-449.
3. Mitchell, J.A., Larkin, S. and Williams, T.J. (1995) Cyclooxygenase-2: regulation and revelance in inflammation. Biochem. Pharmacol. 50, 1535-1542.
4. Mitchell, J.A., Belvisi, M.G., Akaresereenont, P. Robbins, R.A., Kwon, O.J., Croxtall, J., Barnes, P.J. and Vane, J.R. (1994) Induction of cyclo-oxygenase-2 by cytokines in human pulmonary epithelial cells: regulation by dexamethasoner. Br. J. Pharmacol. 113, 1008-1014.
5. Vane, J.R., Mitchell, J.A., Appleton, I., Tomlinson, A., Bishop-Bailey, D.A., Croxtall, J. and Westwick, J. (1996) Inducible isoforms of cyclooxygenase and nitric oxide synthase in inflammation. Proc. Natl. Acad. Sci. U.S.A. 91, 2046-2050.
6. Moncada, S. Palmerm R.M.J. and Higgs, A. (1991) Nitric oxide: physiology, pathyology, and pharmacology. Pharmacol. Rev. 43: 109-142.
7. Moncada, S. Palmerm R.M.J. and Higgs, A. (1993) The L-arginin-nitric oxide pathway. New Eng. J. Med. 329: 2002-2012.
8. Forstermann, U. Gath, I., Schwartz, P., Closs, E.I. and Kleinert, H. (1995) Isoforms of nitric oxide synthase. Biochem Pharmacol. 50: 1321-1332.
9. Paegelow, I., Werner, H., Vietinghoff, G. and Modeer, T. (1995) Release of cytokines from isolated lung strips by bradykinin. Inflamm. Res. 44:306-311.
10. Pang, L. and Knox, A.J. (1998) Bradykinin stimulates IL-8 production in cultured human airway smooth muscle cells: Role of cyclooxygenase products. J. Immunol. 1661: 2509-2515.
11. Pang, L. and Knox, A.J. (1997) PGE2 release by bradykinin in human airway smooth muscle cells: involvement of cyclooxygenase-2 induction. Am. J. Physiol. 273: L1132-1140.