

• 系統編號	RN9511-2376
• 計畫中文名稱	革蘭氏陽性菌細胞壁成分 Peptidoglycan 刺激巨噬細胞前發炎物質釋放之訊息傳遞探討(I)
• 計畫英文名稱	Studies on the Proinflammatory Mediators Release Induced by Peptidoglycan from Gram-Positive Organisms in RAW264.7 Macrophages (I)
• 主管機關	行政院國家科學委員會
• 執行機構	臺北醫學大學醫事技術學系
• 本期期間	9208 ~ 9307
• 報告頁數	13 頁
• 研究人員	林建煌; 陳炳常 Lin, Chien-Huang; Chen, Bing-Chang
• 中文關鍵字	巨噬細胞;NF-kappaB 轉？因子;訊息傳遞;環氧化酶-2
• 英文關鍵字	Macrophages; NF-kappaB; Signal transduction; Cyclooxygenase-2 (COX-2); Peptidoglycan(PGN);Ras/Raf-1/ERK
• 中文摘要	<p>在本計劃中，我們將探討革？氏陽性菌細胞壁成分 peptidoglycan(PGN)誘導 RAW 264.7 巨噬細胞 cyclooxygenase-2(COX-2) 表現之訊息傳遞？徑。PGN 依劑？及時間相關曲線增加 COX-2 的表現，而這個作用可被 Ras 抑制劑(manumycin A)、Raf-1 抑制劑(GW 5074)及 mitogen-activated protein kinase kinase MEK)抑制劑(PD 098059)所抑制。在 RAW 264.7 巨噬細胞給予 PGN 可以依照時間曲線活化 Ras、Raf-1 及 ERK。PGN 所誘導增加 Ras 的活性可被 manumycin A 所抑制；PGN 所誘導增加 Raf-1 的 Ser-338 磷酸化可被 manumycin A 及 GW 5074 所抑制；PGN 所誘導增加 ERK 的活性可被 manumycin A、GW 5074 及 PD 098059 所抑制。RAW 264.7 巨噬細胞給予 PGN 可以增加 IkappaB kinase alpha/beta(IKKalpha/beta)和 IBalpha 磷酸化及 IBalpha 的？解及 kappaB-luciferase 的活性。巨噬細胞給予 NF-kappaB 抑制劑(pyrrolidine dithiocarbamate, PDTC)、IkappaBalpah 磷酸化抑制劑(Bay 117082)及 IkappaB protease 抑制劑(L-1-tosylamido-2-phenylethyl chloromethyl ketone (TPCK) and calpain inhibitorI)皆可抑制 PGN 誘導 COX-2 的表現。PGN 誘導 IKKalpha/beta 和 kappaB-luciferase 的活性也會被 Ras dominant negative mutant (RasN17)、manumycin A、GW5074 和 PD 098059 所抑制。？進一步探討發現在 PGN ？激可以依照時間作用使 p85alpha 及 Ras 結合至 Toll-like receptor 2 。因此在本計劃中首先發現 PGN 可以經由 Ras/Raf-1/ERK 的？徑？活化 IKKalpha/beta 及 NF-kappaB，進一步誘導巨噬細胞 COX-2 的表現。</p>
• 英文摘要	In this study, we investigated the signaling pathway involved in cyclooxygenase-2 (COX-2) expression caused by peptidoglycan

(PGN), a cell wall component of the gram-positive bacterium, *Staphylococcus aureus*, in RAW 264.7 macrophages. PGN caused dose- and time-dependent increases in COX-2 expression, which was attenuated by a Ras inhibitor (manumycin A), a Raf-1 inhibitor (GW 5074), and a mitogen-activated protein kinase kinase (MEK) inhibitor (PD 098059). Treatment of RAW 264.7 macrophages with PGN caused time-dependent activations of Ras, Raf-1, and extracellular signal-regulated kinase (ERK). The PGN-induced increase in Ras activity was inhibited by manumycin A. Raf-1 phosphorylation at Ser338 by PGN was inhibited by manumycin A and GW 5074. The PGN-induced increase in ERK activity was inhibited by manumycin A, GW 5074, and PD 098059. Stimulation of cells with PGN activated IkappaB kinase alpha/beta (IKK α/β), IkappaB α phosphorylation, IkappaB α degradation, and kappaB-luciferase activity. Treatment of macrophages with a NF-kappaB inhibitor (pyrrolidine dithiocarbamate, PDTC), an IkappaB α phosphorylation inhibitor (Bay 117082), and IkappaB protease inhibitors (L-1-tosylamido-2-phenylethyl chloromethyl ketone (TPCK) and calpain inhibitor I) all inhibited PGN-induced COX-2 expression. The PGN-mediated increase in the activities of IKK α/β and kappaB-luciferase were also inhibited by the Ras dominant negative mutant (RasN17), manumycin A, GW 5074, and PD 098059. Further studies revealed that PGN induced the recruitment of p85 α and Ras to Toll-like receptor 2 (TLR2) in a time-dependent manner. Our data demonstrate for the first time that PGN activates the Ras/Raf-1/ERK pathway, which in turn initiates IKK α/β and NF-kappaB activation, and ultimately induces COX-2 expression in RAW 264.7 macrophages.