系統編號	RN9704-2527
------	-------------

• 計畫中文名稱 生物可分解之專一性奈米黃金殼的開發及其在近紅外線熱治療腫瘤上之應用

• 計畫英文名稱 Switch Activation of PI-PLC Downstream Signals in Activated Macrophages with Wortmannin

• 主管機關	行政院國家科學委員會	• 計畫編號	NSC95-2221-E038-011
• 執行機構	台北醫學大學醫事技術學系		
• 本期期間	9508 ~ 9607		
• 報告頁數	12 頁	• 使用語言	英文
• 研究人員	梁有志 Liang, Yu-Chih		
• 中文關鍵字			
• 英文關鍵字	Phosphatidylinositol 3-kinase; Phosphoinositide-specific phospholipase C; Inducible nitric oxide synthase; Wortmannin; Lipopolysaccharide		
• 中文摘要	查無中文摘要		
• 英文摘要	Phosphatidylinositol (4,5)-bisphosphate (PtdIns(4,5)P2) has been known to serve as a substrate for phosphatidylinositol 3-kinase (PI3K) and phosphoinositide-specific phospholipase C (PI-PLC), which can produce PtdIns(3,4,5)P3 and inositol 1,4,5-trisphosphate (Ins(1,4,5)P3) and diacylglycerol (DAG), respectively. In this study, we elucidated the role of PI-PLC during the LPS-activated mouse macrophages RAW264.7 treated with PI3K inhibitor wortmannin. First, wortmannin treatment enhanced Ins(1,4,5)P3 production and iNOS expression in LPS-activated macrophages. Inhibition of PI3K by p85 siRNA also showed an enhancement of iNOS expression. On the other hand, overexpression of PI3K by ras-p110 expression plasmid significantly decreased iNOS expression in LPS-activated macrophages. Second, treatment of PI-PLC inhibitor U73122 reversed the enhancement of iNOS expression in LPS-activated macrophages. However, NF-kappaB activity determined by EMSA assay and reporter plasmid assay did not change during LPS-activated macrophages with or without wortmannin. We propose that the inhibition of PI3K by wortmannin in mouse macrophages enhances the PI-PLC downstream signals, and subsequently increases the LPS induction of iNOS		

expression independently of Akt pathway.