

題名:NF- κ B-activated tissue transglutaminase is involved in ethanol-induced hepatic injury and the possible role of propolis in preventing fibrogenesis

作者:何元順

Ching-Shyang Chen; Chih-Hsiung Wu; Yen-Chun Lai; Wen-Sen Lee; Hsiu-Min Chen; Rong-Jane ChenLi-Ching Chen; Yuan-Soon Ho(correspo

貢獻者:醫學檢驗暨生物技術學系

上傳時間:2009-08-25T02:38:43Z

摘要:Abstract

The increased expression and cross-linking activity of tissue transglutaminase (tTG) have been demonstrated in acute liver injury and fibrosis.

We focused on the molecular mechanisms that contribute to ethanol-induced tTG expression and investigated the efficacy of propolis components

in preventing both the tTG expression in vitro and fibrogenesis in vivo. We demonstrate herein that both ERK1/2 and PI3K/Akt pathways can

regulate the effects of ethanol on NF- κ B-dependent transcription and these signaling pathways may be involved in activation of ethanol-mediated

tTG expression. We also found that administration of pinocembrin (PIN), one of the major components of propolis, inhibited tTG activation and

significantly prevented the development of thioacetamide (TAA)-induced liver cirrhosis. The present study suggests that tTG may be an important

member of the cascade of factors necessary for ethanol-induced liver fibrogenesis and PIN could serve as an anti-fibrogenic agent.