

The protective role of heme oxygenase-1 induction on testicular tissues after testicular torsion and detorsion

黃俊仁

Yang S;Shih HJ;Chow YC;Tsai PS;Huang CJ

摘要

Abstract

PURPOSE: Testicular torsion-detorsion has been identified as an ischemia-reperfusion type of injury. We elucidated the protective role of heme oxygenase-1 super induction on testicular torsion-detorsion injury. **MATERIALS AND METHODS:** Adult male Sprague-Dawley rats were randomly allocated to undergo testicular torsion-detorsion, immediately followed by injection of normal saline, the heme oxygenase-1 inducer hemin or hemin plus the heme oxygenase-1 inhibitor tin protoporphyrin. Another set of rats that underwent sham operation, immediately followed by injection of normal saline, hemin or hemin plus tin protoporphyrin, served as controls. Testes were harvested 4 and 24 hours after detorsion, respectively, in the experimental groups or at comparable time points in the control groups. **RESULTS:** Histological evaluation confirmed that torsion-detorsion caused significant testicular tissue injury. Torsion-detorsion also caused significant increases in the testicular levels of nitric oxide, malondialdehyde, myeloperoxidase activity and heme oxygenase-1. The heme oxygenase-1 inducer hemin significantly enhanced the heme oxygenase-1 expression induced by torsion-detorsion and in turn attenuated testicular injury, and increases in nitric oxide, malondialdehyde and myeloperoxidase activity. In addition, the protective effects of hemin were significantly offset by the heme oxygenase-1 inhibitor tin protoporphyrin. **CONCLUSIONS:** Super induction of heme oxygenase-1 protects testes from torsion-detorsion injury.