

授課講師學經歷

姓名	趙振瑞
課程名稱(主題)	母乳的營養與嬰兒副食品
服務單位(包含職稱)	臺北醫學大學保健營養學系教授
學歷	美國俄亥俄州立大學人體營養與食品管理系博士
經歷	臺北醫學大學保健營養學系教授、國際學術交流中心主任


授課內容摘要

母乳依產後分泌時間分為初乳、過渡乳，及成熟乳，特別是初乳具有許多較高濃度與生長發育有關之調節因子，如乳鐵蛋白、分泌型免疫球蛋白 A (secretary IgA; sIgA)、生長因子，而這些調節因子會隨產後時間增長而減少於母乳中之濃度。母乳中有許多調節因子，如抗微生物因子、抗發炎因子、免疫調節因子，這些調節因子可能會受飲食因素而影響其濃度。

哺乳期間母親的營養狀態會影響母乳品質，進而影響餵補母乳寶寶的營養狀態。研究發現母體攝取的營養素會影響母乳品質與分泌量，母乳中的某些營養素含量，會因母親在飲食中增加攝取而增加，像是如蛋白質、維生素 A、維生素 B 群、維生素 E、錳、碘、鎂等。哺乳期間適當減重婦女，每天平均減少 538 大卡，並不會影響母乳分泌量與品質。但對中度營養不良的哺乳婦女每天增加 500 大卡熱量攝取者較每天僅增加 140 kcal 熱量攝取者明顯增加母乳中 sIgA 含量，對乳鐵蛋白含量則無影響。降低肉類、蛋、乳製品，及高油脂食物攝取量，而增加米飯、蔬菜、豆類，及海藻攝取量會降低 2~5 個月產後婦女母乳中總蛋白質含量。

雞精為一種傳統高蛋白補充品，由於其所含蛋白質分子量較小，因此較易被人體吸收。本研究目的探討懷孕後期(37 週)至產後 3 天補充雞精對哺乳婦女乳汁蛋白組成的影響。於婦產科門診篩選，並經婦產科主治醫師確定為健康產婦共 30 位。依其接受雞精意願分為控制組($n=15$)和雞精組($n=15$)。實驗期間，雞精組受試者每天分別於早、中，及晚餐後 30 分鐘內飲用一瓶雞精(70 mL/瓶)，每日 3 瓶。雞精組平均補充雞精為 18 ± 5 天。控制組於研究進行期間，則限制其高蛋白飲食攝取。所有受試者均維持平日正常飲食。雞精組初乳中乳鐵蛋白、上皮生長因子、轉形生長因子- $\beta 2$ 含量分別顯著較控制組高 34%、62%，及 196% ($p < 0.05$)。但雞精組初乳中總蛋白、酪蛋白，及乳白蛋白含量與控制組相較則無差異。綜合上述，補充雞精可提高哺乳婦女乳汁中乳鐵蛋白、上皮生長因子，及轉形生長因子- $\beta 2$ 含量，可能對胎兒改善免疫功能和刺激生長發育有所助益。

母乳的營養與嬰兒副食品



趙振瑞 教授
臺北醫學大學保健營養學系

1

Contents

- Nutrient requirements for infants
- Human milk
- Effects of nutritional factors on milk proteins
- Starting your baby on solid food
- Conclusion



2

Nutrient Requirements for Infants

- Energy requirement is determined by size, rate of growth, activity, and energy needed for metabolic activities
- Protein requirement:
 - 50% for growth in the first 2 mo
 - declines to 11% by 2-3 years of age
- Fat provides 40-50% calorie supplies (EFA)
- Carbohydrate: primarily lactose
- Water requirement: adequate human milk or formula

3

Dietary Reference Intakes

Age	Energy (kcal/kg)	Protein (g/kg)	Calcium (mg)
0 mo~	110-120	2.4	200
3 mo~	110-120	2.2	300
6 mo~	100	2.0	400
9 mo~	100	1.7	400

DOH, Taiwan, 2002

4

Water Requirements for Infants

Age	Body weight (kg)	Water (ml/d)	Water (ml/kg/d)
3 days	3.0	250-300	80-100
10 days	3.2	400-500	125-150
3 mo	5.4	750-850	140-160
6 mo	7.3	950-1,100	130-155
9 mo	8.6	1,100-1,250	125-145
12 mo	9.5	1,150-1,300	120-135

Barness, 1987

5

Lactogenesis

- colostrum
6-12 h~2-3 d postpartum
- transitional milk
1-2 wk postpartum
- mature milk
> 3 wk postpartum

6

Human Milk Composition

- Immunoglobulins
- lactoferrin
- proline-rich polypeptide
- growth factors
- growth hormone
- leukocytes
- enzymes
- cytokines and lymphokines
- glycoproteins
- carbohydrate, fat, vitamins, and minerals

7

Human Milk Composition

Table 1 Chemical composition of human milk compared with milk from various species (values per 100 ml)

Constituents	Human milk	Cow milk	Buffalo milk
Protein (g)	1.2	3.3	3.8
Casein (g)	0.4	2.8	3.0
Lactalbumin (g)	0.3	0.4	0.4
Lactoglobulin (g)	0.2	0.2	0.2
Fat (g)	3.8	3.7	7.5
Lactose (g)	7.0	4.8	4.4
Calorie (Kcal)	71	69	100
Mineral matter (g)	0.21	0.72	0.80
Calcium (mg)	33	125	210
Phosphorus (mg)	15	96	130
Chlorine (mg)	43	103	112
Magnesium (mg)	4	12	15
Potassium (mg)	55	138	142
Sodium (mg)	15	58	65
Iron (mg)	0.15	0.10	0.20
Copper (mg)	0.04	0.03	0.02
Manganese (mg)	0.7	2.0	—
Zinc (mg)	0.53	0.38	—
Iodine (mg)	0.007	0.021	0.004
Vitamin A (IU)	160	152	206
Vitamin D (IU)	1.4	2.0	—
Thiamine (mg)	0.017	0.04	0.05
Riboflavin (mg)	0.04	0.18	0.10
Nicotinic acid (mg)	0.17	0.08	0.28
Pantothenic acid (mg)	0.20	0.35	—
Vitamin B ₆ (mg)	0.001	0.035	—
Folic acid (μg)	1.3	5.6	3.3
Biotin (μg)	0.4	2.0	—
Vitamin B ₁₂ (μg)	0.03	0.50	0.30
Vitamin C(mg)	4.0	2.0	2.5

Source: Swaminathan (1998)

<http://www.emeraldinsight.com/fig/0170340402001.png>

8


Various Milk Compositions

	Protein (g/100 g)	Fat (g/100 g)	Carbohydrate (g/100 g)	Energy (kcal/100 g)
Human	1.1-1.2	3.8-4.5	6.8-7.0	66-73
Cow	3.1-3.9	3.5-4.9	4.6-5.1	62-80
Goat	2.9-3.4	3.5-4.1	4.6-4.7	62-69


Webb *et al.* 1974, Jensen 1995

9


Various Milk Compositions




Various Milk Compositions in Fatty Acids



Various Milk Compositions in Fatty Acids




Changes in Milk Lactose



<http://www.unu.edu/unupress/food/8F174e/8F174E03.GIF>

13


Changes in Human Milk Compositions



<http://mammary.nih.gov/Reviews/lactation/Neville001/index.html#lactogenesis>

14

Changes in Human Milk Compositions



<http://mammary.nih.gov/Reviews/lactation/Neville001/index.html#lactogenesis>

15

Proteins in Human Milk

• whey/casein (100:0, 86:14, 76:24, 70:30, 65:35, 60:40, 50:50 vs cow 18:82)

• whey

- α-lactalbumin
- lactoferrin
- immunoglobulins
- lysozyme
- lactoperoxidase
- haptocorrin
- lactadherin
- bile salt-dependent lipase

16

Proteins in Human Milk



• casein (~30% of protein fraction)

α s1-, β-, κ-casein

cow: α-/β-/κ-casein = 50:36:13

Casein Micelle

Casein Submicelle



17

Hormones in Human Milk

Table 1. Some hormones and their concentrations found in human milk.*

Hormone	Concentration
Pituitary	
Prolactin	20-90 ng/mL
Growth hormone	5-30 μU/mL
Thyroid-stimulating hormone	2.7-5.0 μU/mL
Hypothalamus	
Thyroid-releasing hormone	0.025-1.5 ng/mL
Luteinizing hormone-releasing hormone	Not available
Somatostatin	23-113 pg/mL
Gonadotropin-releasing hormone	0.1-4.0 ng/mL
Growth hormone-releasing hormone	23-430 pg/mL
Thyroid	
Thyroxine (T ₄)	0.3-12 ng/mL
Triiodothyronine (T ₃)	0.2-0.4 ng/mL
Reverse T ₃	0.008-0.15 ng/mL
Parathyroid	
Parathormone	15 pg/mL
Parathormone-related peptide	30-50 ng/mL
Calcitonin/calcitonin inhibiting protein	0-5 ng/mL
Steroid	
Estrogen	15-840 ng/mL
Progesterone	10-40 ng/mL
Adrenal steroids	0.2-32 ng/mL

*Data from Hemosh (2001).
<http://www.ehponline.org/members/2003/6023/table1.gif>

18

Growth Factors in Human Milk

- insulin-like growth factor (IGF)-I, -II
- epidermal growth factor (EGF)
- nerve growth factor (NGF)
- transforming growth factors (TGF- α , - β 1, - β 2)
- platelet-derived growth factor (PDGF)
- hepatocyte growth factor (HGF)
- colony-stimulating growth factor (CSGF)
- vascular endothelial growth factor (VEGF)

19

Factors in Human Milk

- antimicrobial factors
 - antibacterial factors
 - antiviral factors
 - antiparasite factors
- anti-inflammatory factors
- immunomodulators
 - nucleotides, cytokines
 - anti-idiotypic antibodies, lactoferrin

20

Antibacterial Factors in Human Milk

- secretory IgA, IgG, IgM, IgD
- analogues of epithelial cell receptors (oligosaccharides and sialylated oligosaccharides)
- *Bifidobacterium bifidum* growth factors (oligosaccharides, glycopeptides)
- other bifidobacteria growth factors (α -lactoglobulin, lactoferrin, sialyllactose)
- carbohydrate

21

Antibacterial Factors in Human Milk

- cathelicidin
- casein, κ -casein, phosphorylated β -casein
- complement C1-C9 (mainly C3 and C4)
- α -, β -defensins
- factor binding proteins (zinc, vitamin B₁₂, folate)
- ganglioside
- glycolipid Gb3
- glycoproteins
- α -lactalbumin

22

Antibacterial Factors in Human Milk

- lactoferrin
- lactoperoxidase
- lipids
- lysozyme
- milk cells (80% macrophages, 15% neutrophils, 0.3% B and 4% T lymphocytes)
- mucin
- phosphatidylethanolamine
- sialyllactose

23

Antibacterial Factors in Human Milk

- sialyloligosaccharides on sIgA
- CD14
- sulphatide (sulphogalactosylceramide)
- xanthine oxidase
- secretory leukocyte protease inhibitor
- heparin

24

Antiviral Factors in Human Milk

- secretory IgA, IgG, IgM
- *Bifidobacterium bifidum*
- chondroitin sulphate
- α -, β -defensins
- haemagglutinin inhibitors
- lactadherin (mucin-associated glycoprotein)
- histo-blood group carbohydrates
- lactoferrin

25

Antiviral Factors in Human Milk

- lipid (unsaturated fatty acids and monoglycerides)
- lysozyme
- α 2-macroglobulin
- milk cells
- neutrophil-derived α -defensin-1
- ribonuclease
- secretory leukocyte protease inhibitor
- sialic acid-glycoproteins

26

Antiviral Factors in Human Milk

- soluble intracellular adhesion molecule 1 (sICAM-1)
- soluble vascular cell adhesion molecule 1 (sVCAM-1)
- sulphatide (sulphogalactosylceramide)
- vitamin A
- prostaglandins E1, E2, F2 α
- gangliosides
- glycolipid Gb4
- heparin

27

Antiparasite Factors in Human Milk

- secretory IgA, IgG
- gangliosides
- lipid (free fatty acids and monoglycerides)
- lactoferrin
- macrophages

28

Anti-inflammatory Factors in Human Milk


- secretory IgA
- lactoferrin
- lysozyme
- κ -casein
- antioxidants (β -carotene, α -tocopherol, ascorbate, uric acid)
- EGF
- platelet activating factor-acetylhydrolase
- antiproteases (α 1-antichymotrypsin, α 2-trypsin)
- prostaglandins

29

Nutritional Factors Affecting Milk Proteins



Weight loss
2303 kcal \rightarrow \downarrow 538 kcal
 \downarrow 4.8 \pm 1.2 kg
In 10 weeks




× Milk quantity and quality

Dusdieker *et al.* 1994

30

Nutritional Factors Affecting Milk Proteins

Moderately
undernourished mother
caloric supplement
500 kcal/d vs 140 kcal/d



↑ secretory IgA × lactoferrin


Herias *et al.* 1993

31

Nutritional Factors Affecting Milk Proteins

↓ meat, egg, dairy
products, food
prepared with oil

↑ rice,
vegetables,
beans, seaweed



↓ total protein at 2-5 mo postpartum

Yoneyama *et al.* 1994

32

Manufacture of Chicken Extract

高級雞茸、平取雞油

採用最新科學的酶法提煉方法，以超導磁場及0度低溫的技術，提煉出品質優良、且無任何化學添加劑。

雞油去油

雞油在提煉前經高心率在攝氏零下，經由急凍及解凍的多次循環，成爲不含油質的雞油。


此口爲高級沙律油與手製，因此單層的口味不似普通油質，且含有豐富的自然滋養成分，且能與各種調味料完美結合，使您的料理更添風味。

FATS

<http://www.brands.com.tw/cms.www/main.aspx?sid=415>

33

Experimental Designs



Human Milk Collection

- no breastfeeding 1 h before milk collection
- twice a day
- the interval at least 4 h
- electronic suction pump
- 15-min suction of each breast
- from 1 to 3 d postpartum
- 5-10 mL for protein analyses

38

Demographic and Clinical Characteristics

	Control	Chicken extract
Gestational weight gain, kg	16.4 ± 5.2	15.8 ± 4.9
Gestational age, weeks	38.9 ± 1.0	38.9 ± 0.9
Parity, n	1.4 ± 0.5	1.1 ± 0.4
Primiparous, n	9 (60.0%)	13 (86.7%)
Infant sex	9M, 6F	6M, 9F
Infant birth length, cm	51.3 ± 2.5	54.4 ± 2.3
Infant birth weight, g	3438 ± 205	3250 ± 397
Infant birth head circumference, cm	33.7 ± 0.9	33.4 ± 1.1

Chao *et al.* 2004 39

Daily Dietary Intake

	Control		Chicken extract			
	Pregnancy	Postpartum	Exclude	Include	Exclude	Include
			Pregnancy		Postpartum	
Energy						
kcal/d	1986 ± 293 [‡]	2081 ± 373	2097 ± 449	2168 ± 449	1923 ± 206 [‡]	1994 ± 206
Carbohydrate						
g/d	257 ± 63	257 ± 78	246 ± 77	246 ± 77	228 ± 52	228 ± 52
% energy	51.5 ± 8.1 [‡]	49.0 ± 9.9	46.4 ± 9.0 [*]	44.8 ± 8.8	47.5 ± 9.9	45.8 ± 9.6
Protein						
g/d	82 ± 18 [‡]	101 ± 18 ^{‡#}	91 ± 20	109 ± 20 [‡]	107 ± 30 [#]	125 ± 30 [‡]
% energy	16.7 ± 3.4 [‡]	20.1 ± 5.7 [#]	17.7 ± 4.3	20.6 ± 4.6 [‡]	22.4 ± 6.4 [#]	25.2 ± 6.3
Fat						
g/d	71 ± 16 [‡]	73 ± 25 [‡]	85 ± 24 [*]	85 ± 24	61 ± 17 [#]	61 ± 17
% energy	32.4 ± 6.8	31.3 ± 7.7	36.3 ± 7.0 [*]	35.0 ± 6.8	28.3 ± 7.1 [#]	27.3 ± 6.8

^{*}vs control, [‡]vs exclude, [#]vs include, ^{*}vs pregnancy (P<0.05) Chao *et al.* 2004 40

Milk Protein Concentrations



	Control	Chicken extract
Total protein, g/L	64.9 ± 26.1	75.6 ± 47.5
Lactoferrin, g/L	7.3 ± 3.0	9.8 ± 3.5 [*] ↑ 34%
Epidermal growth factor, µg/L	146.8 ± 73.5	237.6 ± 144.6 [*] ↑ 62%
Transforming growth factor-β2, µg/L	7.8 ± 5.8	23.1 ± 17.4 [*] ↑ 196%
Secretory immunoglobulin A, g/L	4.1 ± 1.3	4.8 ± 1.5

^{*}vs control (P<0.05)

Chao *et al.* 2004

41

Milk Casein and Lactalbumin



Chao *et al.* 2004 42

Effects of Chicken Extract on Human Milk


Protein compositions of human milk can be altered by diet. Supplementation with chicken extract increases colostrum levels of lactoferrin, EGF, and TGF- β 2, which are important for the growth and immune functions of the infants, in lactating women.



43

Breast Milk Intake for Infants

Figure 1:
Usual Breast Milk Intake by Age



WHO 1998¹

http://media.gerber.com/nirt/cm2/upload/074FF5FE-6DA9-4E8F-A5C9-DF3A0DD4AB96/107_02-fig1.jpg

44

Solid Food for Infants

- WHO:

- 0-6 mo → exclusive breastfeeding
 - > 6 mo → solid foods

- Other organizations:

- 0-4 mo → breastfeeding or bottle-feeding
not needed nutritionally, extrusion reflex
 - 4-6 mo → solid foods

45

Solid Food for Infants

- non-allergenic, easily digested, and rich in iron
- one new, "single-ingredient" food introduced per week
- gradually introduced by increasingly coarser textures



46

Solid Food for Infants

行政院衛生署針對4至6個月嬰兒每天飲食建議：

類別	五穀根莖類	蔬菜類	水果類	奶類
食物	米糊或麥糊	胡蘿蔔、菠菜、青江菜、小白菜、空心菜等蔬菜湯	橘子、柳丁、西瓜、蕃石榴、葡萄等自榨稀釋(1:1)果汁	母乳或嬰兒配方奶粉
份量	4湯匙	1~2茶匙	1~2茶匙	5次 170-200 cc/次

47

Solid Food for Infants

行政院衛生署針對7至9個月嬰兒每天飲食建議：

類別	五穀根莖類	蔬菜類	水果	奶類	蛋豆魚肉類
食物	1份：稀飯、麵條、麵線 1/2碗、薄片土司麵包1片、饅頭1/3個、米粉、麥粉4湯匙	菜泥	自榨果汁或果泥	母乳或嬰兒配方奶粉	1份：蛋黃泥2個、豆腐1個四方塊或半盒豆漿 240 cc、魚泥、肉泥、肝泥1兩
份量	2.5-4份	1~2湯匙	1~2湯匙	4次 200-250 cc/次	1-1.5份

48

Solid Food for Infants



行政院衛生署針對9至12個月嬰兒每天飲食建議：

類別	五穀根莖類	蔬菜類	水果	奶類	蛋豆魚肉類
食物	1份：稀飯、麵條、麵線 1/2碗、乾飯1/4碗、薄片土司 麵包1片 饅頭 1/3個、米粉、麥粉4湯匙	切碎蔬菜	自榨果汁或果泥	母乳或嬰兒配方食品	1份：蒸全蛋 1個 豆腐 1個四方塊或半盒 豆漿 240 cc 魚泥、肉泥、肝泥 1兩
份量	4-6份	2~4湯匙	2~4湯匙	1(2)-3次 200-250 cc/次	1.5-2份

49


Solid Food for Infants

- iron-fortified rice cereal
- juice: limited to 120 to 180 mL/d
- vegetables:



50


Baby's First Solid Food



White 2006
<http://www.blisstree.com/breastfeeding123/files/2006/12/food-poll.jpg>

51

Solid Food for Caution



52

Conclusion

The American Academy of Pediatrics (AAP) recommends exclusive breastfeeding for a minimum of 6 mo and introduction of appropriate solid food from 6 mo to 1 yr.

First
BABY'S
Food

53
