

- 79-94.
20. Furukawa, S., Matsubara, T., Tsuji, K., Motohashi, T., Okumura, K., Yabuta, K. Serum soluble CD4 and CD8 levels in Kawasaki disease. *Clin. Exp. Immunol.* 1991; **86**: 134-139.
 21. Pui, C.H., Ip, S.H., Iflah, S., Behm, F.G., Grose, B.H., Dodge, R.K., Crist, W.M., Furman, W.L., Murphy, S.B., Rivera, G.K. Serum interleukin 2 receptor levels in childhood acute lymphoblastic leukemia. *Blood* 1988; **71**: 1135-1137.
 22. Semenza, G., Foa, R., Agostini, C., Zambello, R., Trentin, L., Vinante, F., Benedetti, F., Chilosi, M., Pizzolo, G. High serum levels of soluble interleukin 2 receptor in patients with chronic lymphocytic leukemia.
 23. Pui, C.H., Schell, M.J., Vodian, M.A., Kline, S., Mirro, J., Crist, W.M., Behm, F.G. Serum CD4, CD8, and interleukin-2 receptor levels in childhood acute myeloid leukemia. *Leukemia* 1991; **5**: 249-254.
 24. Kangro, H.O., Ward, A., Argent, S., Heath, R.B. Detection of specific IgM in varicella and herpes zoster by antibody-capture radioimmunoassay. *Epidem. Infect.* 1988; **101**: 187-195.
 25. Vitale, G., Mocciaro, C., Malta, R., Gambino, G., Spinelli, A., Giordano, C., Stassi, G., Arcoleo, F., Milano, S., Cillari, E. Evaluation of serum levels of soluble CD4, CD8 and β 2-microglobulin in visceral human leishmaniasis. *Clin. Exp. Immunol.* 1994; **97**: 280-283.