

題名:Molecular mechanisms in apoptosis induced by magnolol in liver cancer cells

作者:陳盛?

LIN SY; CHANG YT; LIU TD; CHEN SH; PAN S; HO YS; LEE YH; LEE WS

貢獻者:醫學系內科學科

上傳時間:2009-10-06T04:12:18Z

摘要:Magnolol has been reported to have anticancer activity.

In this study we found that treatment with 100 microm magnolol induced apoptosis in cultured human hepatoma (Hep G2) and colon cancer (COLO 205) cell lines but not in human untransformed gingival fibroblasts and human umbilical vein endothelial cells. Our investigation of apoptosis in Hep G2 cells showed a sequence of associated intracellular events that included (a) increased cytosolic free  $Ca^{2+}$ ; (b) increased translocation of cytochrome c (Cyto c) from mitochondria to cytosol; (c) activation of caspase 3, caspase 8, and caspase 9; and (d) downregulation of bcl-2 protein. Pretreatment of the cells with the phospholipase C inhibitor 1-[6-[[[(17 beta)-3-methoxyestra-1,3,5(10)-trien-17-yl]amino]hexyl]-1 H-pyrrole-2,5-dione (U73122) or the intracellular chelator of  $Ca^{2+}$  1,2-bis(2-aminophenoxy)ethane-N,N,N~,N~-tetraacetic acid acetoxymethyl ester (BAPTA/AM) inhibited the subsequent magnolol augmentation of  $[Ca^{2+}]_i$  and also the activation of caspase-8 and caspase-9, so that the occurrence of apoptosis in those cells was greatly reduced. Pretreatment of the cells with ZB4 (which disrupts the Fas response mechanism) also decreased the subsequent magnolol-induced caspase-8 activation and reduced the occurrence of apoptosis. We interpreted these findings to indicate that the above-listed sequence of intracellular events led to the apoptosis seen in Hep G2 cells and that  $[Ca^{2+}]_i$ , Cyto c, and Fas function as intracellular signals to coordinate

those events.