

Yu-Wen Cheng · Ching-Hao Li · Chen-Chen Lee
Jaw-Jou Kang

Alpha-naphthoflavone induces vasorelaxation through the induction of extracellular calcium influx and NO formation in endothelium

Received: 10 June 2003 / Accepted: 5 September 2003 / Published online: 15 October 2003

© Springer-Verlag 2003

Abstract The effect of α -naphthoflavone (α -NF) on vascular function was studied in isolated ring segments of the rat thoracic aorta and in primary cultures of human umbilical vein endothelial cells (HUVECs). α -NF induced concentration-dependent relaxation of the phenylephrine-precontracted aorta endothelium-dependently and -independently at lower and higher concentrations, respectively. The cGMP, but not cAMP, content was increased significantly in α -NF-treated aorta. Pretreatment with *N*^ω-nitro-L-arginine methyl ester (L-NAME) or methylene blue attenuated both α -NF induced vasorelaxation and the increase of cGMP content significantly. The increase of cGMP content induced by α -NF was also inhibited by chelating extracellular Ca²⁺ with EGTA. These results suggest that the endothelium-dependent vasorelaxation induced by α -NF is mediated most probably through Ca²⁺-dependent activation of NO synthase and guanylyl cyclase. In HUVECs, α -NF induced concentration-dependent formation of NO and Ca²⁺ influx. α -NF-induced NO formation was abolished by removal of extracellular Ca²⁺ and by pretreatment with the Ca²⁺ channel blockers SKF 96365 and Ni²⁺, but not by the L-type Ca²⁺ channel blocker verapamil. The Ca²⁺ influx, as measured by ⁴⁵Ca²⁺ uptake, induced by α -NF was also inhibited by SKF 96365 and Ni²⁺. Our data imply that α -NF, at lower concentrations, induces endothelium-dependent vasorelaxation by promoting extracellular Ca²⁺ influx in endothelium and the activation of the NO-cGMP pathway.

Keywords Naphthoflavone · Endothelium · NO · Ca²⁺

Introduction

The endothelium plays an important role in the vascular system. Not only is it a barrier between the blood and tissue, it also produces a variety of vasoactive agents that are important in controlling the body's homeostasis under normal and many pathological conditions (Jaffe 1985; Vanhoutte et al. 1986). In the vascular system, the endothelium, when stimulated by neurotransmitters, hormones, substances derived from platelets and the coagulation system, can evoke vasorelaxation (Furchgott and Zawadzki 1980; Luscher et al. 1988) through the production of NO (Palmer et al. 1988). NO is synthesized from the amino acid L-arginine by oxidation of its terminal guanidine nitrogen by the endothelial cell enzyme nitric oxide synthase (eNOS) (Cobb et al. 1993; Moncada and Higgs 1993) in a calcium-dependent manner (Moncada et al. 1991). Besides its vasorelaxing property, NO released from endothelium is also important in preventing the aggregation of platelets (Furchgott et al. 1984; Furchgott 1984) and inhibiting smooth muscle proliferation (Ignarro et al. 2002; Gewaltig and Kojda 2002).

Flavonoids are substances occurring naturally in fruit, vegetables, grains, barks, roots, stems, flowers, tea and wine (Middleton 1998). More than 4,000 different flavonoids have been identified, many of which are responsible for the attractive colours of flowers, fruits, and leaves (De Groot and Rauen 1998). These natural products are known for their beneficial effects on health, especially for protection against vascular disease and cancer (Birt et al. 1986; Wei et al. 1990; Heo et al. 1992). One of their therapeutically relevant effects on the vascular system may be attributed to their ability to interact with the NO-generating pathway in vascular endothelium (Balestrieri et al. 2003; Duffy and Vita 2003; Youdim et al. 2002). The most frequently studied flavonoid, quercetin, has biological properties consistent with its protective effect on the vascular system (Lanza et al. 1987; Gryglewski et al. 1987; Tzeng et al. 1991; Frankel et al. 1993a, 1993b). The specific effects on vascular system might also come from the broad

C.-H. Li · C.-C. Lee · J.-J. Kang (✉)
Institute of Toxicology, College of Medicine,
National Taiwan University, 1 Jen-Ai Road, Section 1,
Taipei, Taiwan
Tel.: +886-2-23123456 ext 8603, Fax: +886-2-23410217,
e-mail: jjkang@ha.mc.ntu.edu.tw

Y.-W. Cheng
School of Pharmacy, Taipei Medical University,
250 Wu Hsing Street, Taipei, Taiwan

spectrum of modulating effects of flavonoids as antioxidants (Hanasaki et al. 1994; Kerry and Abbey 1997), and inhibitors of ubiquitous enzymes such as lipoxygenase (Alcaraz and Hoult 1985; Moroney et al. 1988), cyclooxygenase (Moroney et al. 1988), phospholipase A2 (Alcraz and Hoult 1985; Fawzy et al. 1988) and protein kinase C (Ferriola et al. 1989). They also inhibit LDL oxidation (De Whalley et al. 1990; Rankin and Leake 1988) and platelet aggregation (Gryglewski et al. 1987; Lanza et al. 1987) and promote vasodilation (Duarte et al. 1993a, 1993b). All these findings have led researchers to use flavonoids as the starting material for drug or health-food development aimed at reducing the risk factor for vascular disease (Formica and Regelson 1995). In addition to the beneficial effect on the vascular system, flavonoids also have antiviral and carcinostatic properties (Buening et al. 1981; Guengerich and Kim 1990; Cholbi et al. 1991; Li et al. 1994; Siess et al. 1995; Sousa et al. 1985). The anti-carcinogenicity of some flavonoids has been attributed to modulation of the cytochrome P450 enzymes that metabolize procarcinogens to their activated form (Benson et al. 1980; Gordon et al. 1991; Kanazawa et al. 1998; Nijhoff et al. 1993; Rodgers and Grant 1998).

α -Naphthoflavone (α -NF) is a prototype flavone that belongs to a group of phytochemicals and is a normal component of human diets (Shou et al. 1994; Das et al. 1994). Their ability to modulate P450-mediated activities was first reported over three decades ago (Diamond and Gelboin 1969). Most of these studies have assessed the effect of α -NF on P450-mediated hydroxylation of benzo(a)pyrene (BP), an environmental pollutant present in cigarette smoke and polluted air that is carcinogenic in experimental animals (Kinoshita and Gelboin 1972). In addition to its inhibition on P450s, α -NF is also an antagonist at the aromatic hydrocarbon receptor (AhR), the cellular receptor of BP and other polycyclic aromatic hydrocarbons (PAHs) (Dong et al. 2001; Jeon et al. 2002). The α -NF isomer, β -naphthoflavone (β -NF), on the other hand, is a strong agonist at the AhR (Staples et al. 1998; Jeon et al. 2002).

The aim of the present study was to investigate the mechanism underlying α -NF-induced vasorelaxation. This question was addressed in thoracic aorta ring segments isolated from rats and in primary cultures of human umbilical vein endothelial cells (HUVECs). We found that α -NF induced endothelium-dependent vasorelaxation in a Ca^{2+} -dependent manner.

Materials and methods

Chemicals. α -NF, β -NF, phenylephrine (PE), acetylcholine (ACh), sodium nitroprusside, trichloroacetic acid (TCA), EGTA and 3-isobutylmethylxanthine (IBMX) were obtained from Sigma (St. Louis, Mo., USA). Cell culture reagents including M-199 medium, L-glutamine, penicillin, streptomycin and fetal bovine serum (FBS) were obtained from Gibco BRL (Grand Island, N.Y., USA). cAMP and cGMP enzyme immunoassay kits were purchased from Cayman Chemical (Ann Arbor, Mich., USA). ^3H -labelled L-arginine and $^{45}\text{Ca}^{2+}$ were purchased from Amersham Life Sciences (Arlington Heights, Ill., USA). All other chemicals were from Sigma. When drugs were dissolved in dimethylsulphoxide (DMSO), the final con-

centration of DMSO in the bathing solution did not exceed 0.1%, a concentration not interfering with muscle contraction or other measurements.

Aortic ring studies. Male Wistar rats (250–300 g) were purchased from the Animal Centre of the College of Medicine, National Taiwan University, Taipei, Taiwan. The thoracic aorta was removed carefully after the rat had been killed by stunning followed by exsanguination. Fat and connective tissues were dissected away in normal Krebs' solution (in mM: NaCl 118.5, KCl 4.8, MgSO_4 1.2, KH_2PO_4 1.2, NaHCO_3 25, glucose 11.1 and CaCl_2 2.5; pH 7.4). The aortae were then cut into rings about 5 mm long in a 10-ml organ bath gassed continuously with 95% O_2 /5% CO_2 at $37 \pm 0.5^\circ\text{C}$. Two L-shaped stainless-steel hooks were inserted into the aortic lumen; one was fixed at the bottom of the bath and the other connected to a force transducer (Hu et al. 2001). The aortic rings were equilibrated in Krebs' solution and maintained under an optimal tension of 1 g for 45 min. During this period the organ baths were perfused with fresh (37°C) buffer solution for 45 min. Once at their optimal length, the segments were allowed to equilibrate for 30 min before experimentation. Contractions were recorded isometrically via a force-displacement transducer (Grass FT.03) connected to a MacLab/8e recorder (ADInstruments, Castle Hill, NSW, Australia). The presence of functional endothelium was assessed by determining the ability of $10 \mu\text{M}$ ACh to induce more than 80% relaxation of rings precontracted with $3 \mu\text{M}$ PE. The endothelium was removed by rubbing the luminal surface gently with a cotton ball. Successful removal of endothelium was confirmed by the absence of ACh-induced relaxation. The denuded aorta was also challenged with PE and a high $[\text{K}^+]$ (60 mM) to ensure that the vessel had not been damaged during denudation. Aortic rings with a normal contractile response were then used for experiments.

α -NF and β -NF concentration/response curves were obtained by adding increasing concentrations of these substances (0.1–100 μM) to rings precontracted with $3 \mu\text{M}$ PE after the response to the previous concentration had stabilized. To examine the effect of NOS inhibition, N^{ω} -nitro-L-arginine methyl ester (L-NAME, 300 μM) was added 10 min before PE.

Rat aorta cAMP and cGMP measurement. Rat aorta cyclic nucleotide contents were determined according to Kauffman et al. (1987). Depending on the purpose of the experiment, the aorta rings were either placed in Ca^{2+} -free Krebs' (containing EGTA 2.5 mM) buffer or pretreated with the inhibitors L-NAME (300 μM) or methylene blue (10 μM) for 10 min. Test compounds, such as sodium nitroprusside (10 μM), ACh (10 μM), forskolin (10 μM), α - or β -NF (10–100 μM) were added and the ring incubated for 5 min. After incubation with test compounds, the rat aortic rings were frozen rapidly in liquid N_2 and stored at -70°C . For assay, the tissue was homogenized in 0.5 ml 10% TCA in a Potter glass homogenizer. The homogenates were centrifuged at 10,000 g for 5 min and supernatants removed and extracted 4 times with 3 vol ether. cAMP and cGMP contents were then assayed using enzyme immunoassay kits. The precipitates were used for protein determination (Lowry et al. 1951).

HUVEC isolation and culture. Human umbilical cords were obtained from the Hospital of the National Taiwan University, Taipei, Taiwan. HUVECs were isolated by enzymatic digestion from 20-cm-long umbilical cord vein segments filled with 0.1% collagenase (Rosenkranz-Weiss et al. 1994). After 15 min incubation at 37°C , the vein segments were perfused with 30 ml medium 199 containing 10 U/ml penicillin and 100 $\mu\text{g}/\text{ml}$ streptomycin to collect the cells. After centrifugation for 8 min at 900 g, the cell pellet was re-suspended in the same medium supplemented with 20% heat-inactivated FBS, 30 $\mu\text{g}/\text{ml}$ endothelial cell growth supplement (ECGS) and 90 $\mu\text{g}/\text{ml}$ heparin. Confluent primary cells were detached using trypsin-EDTA (0.05:0.02% v/v), and HUVECs from passage 2 were used in the present study. Cultures had typical cobblestone morphology and stained uniformly for human von Willebrand factor (vWF) (Janel et al. 1997) as assessed by indirect immunofluorescence.

NO determination. HUVECs cultured in 12-well plates were washed twice with in a HEPES buffer (in mM: HEPES 10, NaCl 145, KCl 5, CaCl₂ 1, MgCl₂ 1, Na₂HPO₄ 1, glucose 10, pH 7.4) and then incubated at 37 °C in the same buffer for 30 min with various concentrations of α -NF (1–100 μ M) or ACh (30 μ M) as positive control. Supernatants were collected and then injected into the nitrogen-purge chamber containing vanadium (III) chloride in HCl at 91 °C. All NO metabolites are liberated as gaseous NO and react with ozone to form activated nitrogen dioxide that luminesces in the red and far-red spectrum. The chemiluminescent signals were detected by a nitric oxide analyser (NOA280, Sievers Instruments, Boulder, Colo., USA) accordingly (Ewing and Janero 1998). The cells were detached and homogenized for protein determination. For calculation of concentrations, the area under the curve was converted to nanomolar NO using an NaNO₃ standard curve and the final data was expressed in picomoles/milligram protein.

⁴⁵Ca²⁺ uptake. The ⁴⁵Ca²⁺ influx measurement was modified from Cheng and Kang (1997). Confluent HUVECs cultured in 6-well plates were washed twice with HEPES buffer and then incubated in the same buffer containing ⁴⁵Ca²⁺ (10 μ Ci/ml) and treated with test compounds with or without the blockers for 5 min. Two concentrations of α -NF (50 and 100 μ M) and blockers, including the receptor-operated Ca²⁺ channel blocker, SKF96365 (30 μ M), the non-specific Ca²⁺ channel blocker Ni²⁺ 1 mM and the L-type Ca²⁺ channel blocker verapamil (2 μ M) were used. After incubation, the supernatants were aspirated and the cells washed 3 times with cold HEPES buffer containing 10 mM LaCl₃. Cells were lysed with 0.01 N NaOH and the cell ⁴⁵Ca²⁺ content measured by scintillation counter (Model 2200; Beckman, Palo Alto, Calif., USA).

Statistical analysis. Data are expressed as means \pm SEM from *n* experiments. The significance of differences between means was established using Student's *t*-test, with *P* < 0.05 being considered significant. EC₅₀ values were calculated from five regression lines. Each regression line was constructed with between three and five points. These points corresponded to response magnitudes of 20–80%.

Results


Vasorelaxant effect of α -naphthoflavone in rat aorta

A transient phasic contraction followed by a tonic contraction was induced by PE (Fig. 1A) and the contracted aorta could be relaxed by addition of ACh (10 μ M) through the induction of NO formation in endothelium-intact but not in the denuded (endothelium removed) aortic rings. As seen in Fig. 1, 1 and 50 μ M α -NF induced vasorelaxation of intact aorta precontracted with PE (3 μ M) by 48.58 ± 10.7 and $74.72 \pm 6.2\%$, respectively. The vasorelaxation induced by α -NF was greatly attenuated in denuded aorta, suggesting that most of the relaxation caused by α -NF was endothelium dependent. Pretreatment with the NOS inhibitor L-NAME (Fig. 1B) or the guanylyl cyclase inhibitor methylene blue (data not shown) also attenuated the α -NF-induced vasorelaxation, suggesting that this vasorelaxation effect of α -NF might be due to the activation of NO synthesis in the endothelium.

Both α -naphthoflavone and β -naphthoflavone induce vasorelaxation in the rat aorta

Increasing concentrations of α -NF or β -NF were added cumulatively (0.1–100 μ M) to induce relaxation of the precontracted, intact or denuded aorta or the aorta pretreated with L-NAME (Fig. 2). α -NF induced vasorelaxation in the intact aorta concentration dependently with an EC₅₀ of 0.95 ± 0.13 μ M. The α -NF-induced vasorelaxation was largely prevented (~80%) in denuded or L-NAME-treated aortae. α -NF also induced vasorelaxation in denuded or L-NAME-treated aortae, however at much higher con-

Fig. 1A, B Effect of α -naphthoflavone (α -NF) on contractility of isolated rat aortic rings. **A** Phenylephrine (PE, 3 μ M)-precontracted, intact (left) or denuded (right) rings. **B** *N*^ω-nitro-L-arginine methyl ester (L-NAME, 300 μ M)-pretreated rings. In denuded aorta, the endothelium was removed by rubbing with a cotton ball, and the absence of 10 μ M acetylcholine (ACh)-induced relaxation was taken as an indicator of successful denudation. The experiments were repeated with at least three different preparations






Fig. 2 Concentration/response curves for α -NF (A) and β -NF (B)-induced vasorelaxation of PE-precontracted rat thoracic aortic rings with (●) or without (■) endothelium, or in the presence of L-NAME 300 μ M (▲). Means \pm SEM, $n=6$

centrations (Fig. 2A). Compared with α -NF, the isomer β -NF had a much weaker effect vasorelaxant effect ($EC_{50} > 100 \mu$ M, Fig. 2B).

Effect of α -naphthoflavone on cAMP and cGMP content in rat aorta

The above data suggest that the endothelium-dependent vasorelaxation induced by α -NF at lower concentrations was due to activation of eNOS. NO, once generated, can activate guanylyl cyclase, with subsequent generation of cGMP in many cells, including smooth muscle (Moncada and Higgs 1993). The effects of α -NF and β -NF on cyclic nucleotide formation in aortic rings were thus investigated and data are summarized in Table 1. Sodium nitroprusside, an NO donor, and forskolin, an adenylyl cyclase activator, increased cGMP and cAMP contents in aorta, respectively. α -NF concentration-dependently increased cGMP but not cAMP content. β -NF also increased cGMP content, but to a lesser degree.

ACh (10 μ M) increased cGMP formation significantly in intact (2.15 ± 0.09 pmol/mg protein) but not in denuded aorta (0.09 ± 0.02 pmol/mg protein) relative to control (0.11 ± 0.03 pmol/mg protein) (Table 2). ACh-induced cGMP formation was inhibited in aorta pretreated with L-NAME (300 μ M) (0.15 ± 0.02 pmol/mg protein) or methylene blue (0.53 ± 0.10 pmol/mg protein). Induction of cGMP forma-

Table 1 Effects of α -naphthoflavone and β -naphthoflavone on cAMP and cGMP contents of isolated rat aortic rings. cAMP and cGMP contents were measured as described in Methods. Means \pm SEM, $n=6$ individual experiments (ND not determined)

Treatment		Cyclic GMP (pmol/mg protein)	Cyclic AMP (pmol/mg protein)
Control		0.14 ± 0.06	1.33 ± 0.25
Sodium nitroprusside	10 μ M	$1.72 \pm 0.56^{***}$	ND
Forskolin	10 μ M	ND	$9.57 \pm 1.56^{***}$
α -Naphthoflavone	10 μ M	0.31 ± 0.07	1.73 ± 0.28
	50 μ M	$1.09 \pm 0.37^*$	2.18 ± 0.26
	100 μ M	$1.48 \pm 0.23^{**}$	1.98 ± 0.32
β -Naphthoflavone	10 μ M	0.13 ± 0.08	1.00 ± 0.21
	50 μ M	0.36 ± 0.11	1.18 ± 0.28
	100 μ M	0.55 ± 0.09	1.73 ± 0.31

* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs. control (Student's t -test)

tion by α -NF (100 μ M, 1.48 ± 0.23 pmol/mg protein) was also abolished in the denuded aorta (0.21 ± 0.17 pmol/mg protein) and in the aorta treated with 300 μ M L-NAME (0.25 ± 0.11 pmol/mg protein) or 10 μ M methylene blue (0.53 ± 0.22 pmol/mg protein). Interestingly, the increase of cGMP content induced by α -NF was also diminished when extracellular Ca^{2+} was chelated with EGTA (Table 2), suggesting that α -NF-induced cGMP formation was dependent on both Ca^{2+} and the endothelium.

Effect of α -naphthoflavone on NO formation in HUVECs

The direct effect of α -NF on endothelial cells was investigated further in primary cultures of HUVECs and the data are summarized in Fig. 3. α -NF concentration-dependently induced NO formation in HUVECs, with maximal induction at 100 μ M (Fig. 3A). ACh (30 μ M) also induced significant NO formation in HUVECs both in normal and Ca^{2+} -free (5 mM EGTA) HEPES buffer (8.79 ± 0.68 and 8.07 ± 1.58 pmol/mg protein) (Fig. 3B). The NO formation in HUVECs induced by α -NF (50 μ M: 8.90 ± 1.26 , 100 μ M: 10.72 ± 1.74 pmol/mg protein) was attenuated significantly in Ca^{2+} -free HEPES buffer (2.60 ± 0.66 and 2.30 ± 0.32 pmol/mg protein respectively), suggesting that α -NF induced NO formation was dependant on extracellular Ca^{2+} . α -NF-induced NO formation was blocked by pretreatment with the receptor-operated Ca^{2+} channel blocker SKF96365 (30 μ M) and the non-specific Ca^{2+} channel blocker, Ni^{2+} (1 mM), but not the L-type Ca^{2+} channel blocker verapamil (2 μ M) (Fig. 3C). In contrast, β -NF did not induce NO formation at concentrations up to 100 μ M (data not shown).


α -Naphthoflavone induces $^{45}Ca^{2+}$ influx in HUVECs

Both α -NF-induced cGMP (Table 2) and NO (Fig. 3B) formation were attenuated in the absence of extracellular Ca^{2+} , suggesting that α -NF might exert its effect through

Table 2 Effects of blockers on acetylcholine- and α -naphthoflavone-induced cGMP formation. Means \pm SEM, $n=6$ individual experiments. Intact aorta was used in all experiments unless specified

Treatment	Cyclic GMP (pmol/mg protein)
Control	0.11 \pm 0.03
ACh (10 μ M)	2.15 \pm 0.09***
ACh (Denuded, 10 μ M)	0.09 \pm 0.02
L-NAME (300 μ M)	0.07 \pm 0.07
Methylene blue (10 μ M)	0.13 \pm 0.04
Ca ²⁺ -free (2.5 mM EGTA)	0.05 \pm 0.01
ACh (10 μ M)	2.15 \pm 0.09***
+L-NAME (300 μ M)	0.15 \pm 0.02
+Methylene blue (10 μ M)	0.53 \pm 0.10
α -NF (100 μ M)	1.48 \pm 0.23***
+L-NAME (300 μ M)	0.25 \pm 0.11
+Methylene blue (10 μ M)	0.53 \pm 0.22
+Ca ²⁺ -free (2.5 mM EGTA)	0.18 \pm 0.02
α -NF (Denuded, 100 μ M)	0.21 \pm 0.17

*** $P<0.001$ vs. control (Student's t -test)



induction of extracellular Ca²⁺ influx. The effect of α -NF on Ca²⁺ flux was investigated using the ⁴⁵Ca²⁺ loading method in adherent HUVECs. As seen in Fig. 4, a 30~40% increase of Ca²⁺ flux was observed in α -NF treated HUVECs. This was inhibited by SKF96365 and Ni²⁺, but not verapamil. As seen with NO formation, β -NF did not induce ⁴⁵Ca²⁺ influx at concentrations up to 100 μ M (data not shown).

Discussion

In the vascular system, NO is synthesized by eNOS after the latter's activation in endothelial cells and stimulates cGMP production by activating soluble guanylyl cyclase in the adjacent smooth muscle (Palmer et al. 1987). Increased cGMP causes contracted muscle to relax, possibly through lowering the intracellular [Ca²⁺], most likely by increasing Ca²⁺ efflux to the extracellular space and Ca²⁺ reuptake into intracellular stores (Lincoln et al. 1990; Lincoln and Cornwell 1991; Ganitkevich et al. 2002) or by dephosphorylation of myosin light-chain kinase (Ganitkevich et al. 2002; Silveira et al. 1998). Flavonoids exert physiological actions on various biological systems including the vascular system. Studies into the correlation between the low mortality rate due to cardiovascular disease and the red wine consumption in Mediterranean populations (Renaud and DeLorgeril 1992) indicates that the flavonoids in red wine are responsible, at least in part, for this effect (Formica and Regelson 1995).

The present study showed that α -NF relaxed the endothelium-intact rat aorta with an EC₅₀ of 0.95 \pm 0.13 μ M,

Fig. 3A–C Effect of α -NF on NO formation in human umbilical vein endothelial cells (HUVECs). **A** HUVECs were treated with 50 or 100 μ M α -NF. **B** HUVECs were treated with ACh (30 μ M) or α -NF (50 or 100 μ M) for 30 min in normal HEPES buffer with 1 mM CaCl₂ or in Ca²⁺-free HEPES buffer (containing 5 mM EGTA with no added CaCl₂). **C** Effect of pretreatment with various substances: acetylcholine (ACh, 30 μ M); SKF96365 (30 μ M); Ni (1 mM); verapamil (2 μ M) as indicated. Means \pm SEM, $n=3$ independent experiments. * $P<0.05$ vs. respective control




Fig. 4 Effect of α -NF on $^{45}\text{Ca}^{2+}$ -influx. HUVECs were incubated in HEPES buffer containing $^{45}\text{Ca}^{2+}$ (10 $\mu\text{Ci}/\text{ml}$) and treated with test compounds for 5 min: α -NF (50 or 100 μM) with or without blockers (SKF96365, 30 μM ; Ni, 1 mM and verapamil, 2 μM) as indicated. Means \pm SEM, $n=3$ independent experiments. * $P<0.05$ vs. respective control

a value lower than flavonoid EC_{50} s reported previously (Fitzpatrick et al. 1993; Jimenez et al. 1999; Kim et al. 2000). Endothelial denudation significantly attenuated α -NF-induced relaxation, suggesting that most of the vasorelaxant effect of α -NF was endothelium dependent. Pretreatment with L-NAME, a NOS inhibitor, or methylene blue, an inhibitor of guanylyl cyclase, inhibited the α -NF-induced relaxation to an extent similar to that seen in the endothelium-denuded preparations. These findings suggest that activation of NOS might be responsible for the endothelium-dependent vasorelaxation induced by α -NF. Consistent with this is the increased cGMP content in the α -NF-treated aorta. Whilst α -NF at 1 μM induced nearly 50% relaxation of the isolated aorta, a significant increase in cGMP content was achieved only in aortae exposed to α -NF concentrations exceeding 10 μM . Several possibilities might explain this discrepancy. First, although the increase in cGMP content in the aorta treated with 10 μM α -NF was not significant, the trend was still apparent compared with the basal level. Without knowing the exact correlation between intracellular cGMP content and the relaxation effect, it is difficult to compare concentration/responses relationships from the two measurements. It is possible that only a small amount of cGMP is needed to induce vasorelaxation. Second, cGMP content was measured using an enzyme immunoassay kit, the sensitivity of which for cGMP is limited, especially with respect to the latter's extraction from whole tissue. Finally, we have shown also that α -NF induces endothelium-independent relaxation. The EC_{50} obtained from muscle relaxation measurement can therefore be expected to be lower than that obtained from cGMP measurement. The steric isomer β -NF also induced vasorelaxation, however, at much higher concentrations with an $\text{EC}_{50}>100 \mu\text{M}$. The endothelium-de-

pendent vasorelaxation induced by β -NF is most probably through the NO-cGMP pathway, as for α -NF, since β -NF treatment also augmented the cGMP formation in aorta.

Three NO synthase (NOS) isoforms have been identified to date (Bredt et al. 1991; Sessa et al. 1992; Xie et al. 1992). eNOS and neural NOS (nNOS) have been shown to be Ca^{2+} /calmodulin-dependent and expressed are constitutively mainly in endothelial cells and neurons, respectively. eNOS and nNOS are activated mainly by an increase of cytosolic $[\text{Ca}^{2+}]$ (Mayer et al. 1989). Another NOS isoform, inducible NOS (iNOS), can be induced by endotoxin and cytokines in immune cells and is independent of Ca^{2+} (Kerwin and Heller 1994; Iyengar et al. 1985). In the present study, the α -NF-induced increase of cGMP content was attenuated when extracellular Ca^{2+} was chelated with EGTA. This implies that the activation of NOS may have been due to an α -NF-induced influx of extracellular Ca^{2+} into the endothelium. The results from the experiments with HUVECs support this contention. First, α -NF treatment augmented NO release from HUVECs, a response that was abolished when extracellular Ca^{2+} was chelated. Second, Ca^{2+} influx, as evident by the increase of $^{45}\text{Ca}^{2+}$ uptake, in HUVECs was induced by α -NF.

Both the increase of NO formation and Ca^{2+} influx induced by α -NF in HUVECs were inhibited by the Ca^{2+} channel blockers Ni^{2+} and SKF96365. These results suggest that α -NF treatment induced Ca^{2+} influx through an SKF96365- and Ni^{2+} -sensitive Ca^{2+} channel in the endothelium. SKF 96365 inhibits the Ca^{2+} influx through the non-selective cationic channel activated by internal Ca^{2+} store depletion by endoplasmic reticulum Ca^{2+} -ATPase inhibitors or receptor agonists (Low et al. 1996; Millanvoye-Van Brussel et al. 1999). However, the SKF 96365-sensitive Ca^{2+} channel can also be activated directly without depletion of internal Ca^{2+} store (Inazu et al. 1995). In addition, SKF 96365 inhibits the unidentified Ca^{2+} channel activated by mechanical stress in endothelial cells (Yao et al. 2000). Ni^{2+} , on the other hand, is a non-specific Ca^{2+} channel blocker and inhibits the Ca^{2+} channel activated in endothelial cells by the agonist ACh (Wang et al. 1996), the Ca^{2+} pump inhibitor cyclopiazonic acid (Li and van Breemen 1996) and blood flow (Yao et al. 2000). It is interesting to note that β -NF treatment also induces (Xie et al. 2002) or potentiates (Graier et al. 1995) capacitative Ca^{2+} influx in endothelial cells, possibly through the formation of P450 metabolite 5,6-epoxyeicosatrienoic acid (5,6-EET), a calcium influx factor (CIF). However, our data showed that α -NF, a P450 inhibitor, is a more potent inducer of vasorelaxation and Ca^{2+} influx in endothelial cells. These results imply that formation of CIF through P450 metabolism might not be involved in the induction of Ca^{2+} influx by naphthoflavone compounds. A previous study has also shown that flavonoids can induce Ca^{2+} influx in endothelium and hence vasorelaxation through activation of tetraethylammonium-sensitive K^{+} -channels (Kim et al. 2000). The inability of tetraethylammonium to inhibit the α -NF induced vasorelaxation (data not shown) also precludes this possibility. Moreover, the L-type voltage-gated Ca^{2+} channel blocker verapamil did not inhibit the α -NF in-

duced NO formation and Ca²⁺ influx. This implies that either the Ca²⁺ influx induced by α -NF is not through L-type voltage-gated Ca²⁺ channel or that there are no L-type voltage-gated Ca²⁺ channels in HUVECs. The presence of L-type voltage-gate Ca²⁺ channels in endothelial cells is controversial. Although lack of voltage-gated Ca²⁺ channels has been reported in endothelial cells isolated from the porcine coronary artery (Uchida et al. 1999) and the rabbit and rat aorta (Muraki et al. 2000), the voltage-gated Ca²⁺ channels have been demonstrated in freshly isolated capillary endothelial cells from bovine adrenal glands (Bossu et al. 1992a, 1992b) and cerebral microvascular endothelial cells from newborn pig brain cortex (Yakubu et al. 2002). However, to our knowledge, L-type Ca²⁺ channels have not been demonstrated in human endothelial cells (Ding and Vaziri 2000). Further investigation will be needed to understand the exact mechanism of how or which Ca²⁺ channel was activated by α -NF in endothelium.

Our study showed also that higher concentrations of α -NF also induced vasorelaxation in the endothelium-denuded aorta. The EC₅₀ for this endothelium-independent effect was estimated to be 200-fold higher than that of endothelium-dependent effect. Flavonoids such as dioclein (Trigueiro et al. 2000) induce endothelium-independent vasorelaxation by inhibiting voltage-dependent Ca²⁺-influx and the release of intracellular Ca²⁺ store in rat aorta. Eriodictyol, a flavonoid obtained from *Satureja obovata* also induces vasodilatation by inhibiting Ca²⁺ influx in rat aorta (Ramon Sanchez de Rojas et al. 1999). α -NF's endothelium independent vasorelaxation may be mediated by the same mechanism. This however, awaits further investigation.

Flavonoids, including genistein (3–100 μ M); kaempferol (3–60 μ M) and quercetin (1–100 μ M), increase intracellular cAMP content in uterine smooth muscle (Revuelta et al. 1999). An increase of cAMP content in smooth muscle also induces vasorelaxation (Lincolin et al. 1990; Lincolin and Cornwell 1991). In the present study, however, the cAMP content, unlike cGMP, was not increased in α -NF treated aorta. This suggests that cAMP might not play an important role in α -NF induced vasorelaxation. In addition to the effect on vascular endothelial cells, preliminary findings show that α -NF also inhibits the platelet aggregation induced by collagen, arachidonic acid, platelet activation factor and ADP (Y.-W. Cheng, C.-H. Li, C.-C. Lee, J.-J. Kang, unpublished data).

In conclusion, the present study demonstrates that the flavonoid compound α -NF promotes the influx of extracellular Ca²⁺ and release of NO by vascular endothelium. In addition to its vasorelaxant effect, NO is considered an important anti-atherogenic factor by virtue of its inhibitory effect on platelet aggregation (Furchgott 1984) and smooth muscle proliferation (Ignarro et al. 2002). α -NF also exerts anti-carcinogenic effects by virtue of its inhibition of P450 (Andries et al. 1990; Shimada et al. 1998; Tassaneeyakul et al. 1993) and antagonism at the AhR (Dong et al. 2001; Jeon et al. 2002). The results of this study provide evidence for a further beneficial effect of α -NF on the vascular system.

Acknowledgements This study was supported in part by a grant from the National Science Council, Taiwan.

References

- Alcaraz MJ, Hoult JRS (1985) Actions of flavonoids and the novel anti-inflammatory flavone, hypolaetin-8-glucoside on prostaglandin biosynthesis and inactivation. *Biochem Pharmacol* 34: 2477–2482
- Andries MJ, Lucier GW, Goldstein J, Tompson CL (1990) Involvement of cytochrome P-450c in α -naphthoflavone metabolism by rat liver microsomes. *Mol Pharmacol* 37:990–995
- Balestrieri M.L., Castaldo D, Balestrieri C, Quagliuolo L, Giovane A, Servillo L (2003) Modulation by flavonoids of PAF and related phospholipids in endothelial cells during oxidative stress. *J Lipid Res* 44:380–387
- Benson AM, Hunkeler MJ, Talalay P (1980) Increase of NAD(P)H:quinone reductase by dietary antioxidants: possible role in protection against carcinogenesis and toxicity. *Proc Natl Acad Sci USA* 77:5216–5220
- Birt DF, Walker D, Tibbles MB, Bresnick K (1986) Anti-mutagens and anti-promotion by apigenin, robinetin and indole-3-carbinol. *Carcinogenesis* 7:959–963
- Bossu JL, Elhamdani A, Feltz A (1992a) Voltage-dependent calcium entry in confluent bovine capillary endothelial cells. *FEBS Lett* 299:239–242
- Bossu JL, Elhamdani A, Feltz A, Tanzi F, Aunis D, Thierse D (1992b) Voltage-gated Ca entry in isolated bovine capillary endothelial cells: evidence of a new type of BAY K8644-sensitive channel. *Pflugers Arch* 420:200–207
- Bredt DS, Hwang PM, Glatt CE, Lowenstein C, Reed RR, Snyder SH (1991) Cloned and expressed nitric oxide synthase structurally resembles cytochrome O-450 reductase. *Nature* 351:714–718
- Buening MK, Chang RL, Huang ML, Fortner JC, Wood AW, Conney AH (1981) Activation and inhibition of benzo(a)pyrene and aflatoxin B1 metabolism in human liver microsomes by naturally occurring flavonoids. *Cancer Res* 41:67–72
- Cheng YW, Kang JJ (1997) Mechanism of vasorelaxation of thoracic aorta caused by xanthone. *Eur J Pharmacol* 336:23–28
- Cholbi MR, Paya M, Alcaraz MJ (1991) Inhibitory effects of phenolic compounds on CCl₄ induced microsomal lipid peroxidation. *Experientia* 47:195–199
- Cobb JP, Cunnion RE, Danner RL (1993) Nitric oxide as a target for therapy in septic shock. *Crit Care Med* 21:1261–1263
- Das A, Wang JH, Lien EJ (1994) Carcinogenicity, mutagenicity and cancer preventing activities of flavonoids: a structure-system-activity relation (SSAR) analysis. *Prog Drug Res* 42:133–166
- De Groot H, Rauen U (1998) Tissue injury by reactive oxygen species and the protective effects of flavonoids. *Fundam Clin Pharmacol* 12:249–255
- De Whalley CV, Rankin SM, Hoult JRS, Jessup W, Lerke DE (1990) Flavonoids inhibit the oxidative modification of low density lipoproteins by macrophages. *Biochem Pharmacol* 39: 1743–1750
- Diamond L, Gelboin HV (1969) Alpha-naphthoflavone: an inhibitor of hydrocarbon cytotoxicity and microsomal hydroxylase. *Science* 166:1023–1025
- Ding Y, Vaziri ND (2000) Nifedipine and diltiazem but not verapamil up-regulate endothelial nitric oxide synthase expression. *J Pharmacol Exp Ther* 292:606–609
- Dong W, Teraoka H, Kondo S, Hiraga T (2001) 2,3,7,8-Tetrachlorodibenzo-p-dioxin induces apoptosis in the dorsal mid-brain of zebrafish embryos by activation of aryl hydrocarbon receptor. *Neurosci Lett* 303:169–172
- Duarte J, Perez-Vizcaino F, Utrilla P, Jimenez J, Tamargo J, Zarzuelo A (1993a) Vasodilatory effects of flavonoids in rat aortic smooth muscle: structure-activity relationships. *Gen Pharmacol* 24:857–862

- Duarte J, Perez-Vizcaino F, Zarzuelo A, Jimenez J, Tamargo J (1993b) Vasodilator effects of quercetin in isolated rat vascular smooth muscle. *Eur J Pharmacol* 239:1–7
- Duffy SJ, Vita JA (2003) Effects of phenolics on vascular endothelial function. *Curr Opin Lipidol* 14:21–27
- Ewing JF, Janero DR (1998) Specific S-nitrosothiol (thionitrite) quantification as solution nitrite after vanadium (III) reduction and ozone-chemiluminescent detection. *Free Radic Biol Med* 25:621–628
- Fawzy AA, Vishwanath BS, Franson RC (1988) Inhibition of human non-pancreatic phospholipases A2 by retinoids and flavonoids. Mechanism of action. *Agents Actions* 25:394–400
- Ferriola PC, Cody V, Middleton E Jr (1989) Protein kinase C inhibition by flavonoids. Kinetic mechanisms and structure-activity relationships. *Biochem Pharmacol* 38:1617–1624
- Fitzpatrick DF, Hirschfield SL, Coffey RG (1993) Endothelium-dependent vasorelaxing activity of wine and other grape products. *Am J Physiol* 265:H774–H778
- Formica JV, Regelson W (1995) Review of the biology of quercetin and related bioflavonoids. *Food Chem Toxicol* 33:1061–1080
- Frankel EN, Kanner J, German JB, Parks E, Kinsella JE (1993a) Inhibition of oxidation of human low-density lipoprotein by phenolic substances in red wine. *Lancet* 34:454–457
- Frankel EN, Waterhouse AL, Kinsella JE (1993b) Inhibition of human LDL oxidation by resveratrol. *Lancet* 341:1103–1104
- Furchgott RF (1984) The role of endothelium in the responses of vascular smooth muscle to drugs. *Annu Rev Pharmacol Toxicol* 24:175–197
- Furchgott RF, Zawadzki JV (1980) The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature* 288:373–376
- Furchgott RF, Cherry PD, Zawadzki JV, Jothianandan D (1984) Endothelial cells as mediators of vasodilation of arteries. *J Cardiovasc Pharmacol* 6 (Suppl 2):S336–S343
- Ganitkevich V, Hasse V, Pfister G (2002) Ca²⁺-dependent and Ca²⁺-independent regulation of smooth muscle contraction. *J Muscle Res Cell Motil* 23:47–52
- Gewaltig MT, Kojda G (2002) Vasoprotection by nitric oxide: mechanisms and therapeutic potential. *Cardiovasc Res* 55:250–260
- Gordon GB, Prochaska HJ, Yang LYS (1991) Inhibition of NAD(P)H:quinone reductase in human peripheral blood lymphocytes. *Carcinogenesis* 12:2393–2396
- Graier WF, Simecek S, Sturek M (1995) Cytochrome P450 monooxygenase regulated signalling of Ca²⁺ entry in human and bovine endothelial cells. *J Physiol (Lond)* 482:259–274
- Gryglewski RJ, Korbut R, Robak J, Swies J (1987). On the mechanism of anti-thrombotic action of flavonoids. *Biochem Pharmacol* 36:317–322
- Guengerich FP, Kim DH (1990) In vitro inhibition of dihydropyridine oxidation and AFB1 activation in human liver microsomes by naringenin and other flavonoids. *Carcinogenesis* 11:2275–2279
- Hanasaki Y, Ogawa S, Fukui S (1994) The correlation between active oxygen scavenging and antioxidative effects of flavonoids. *Free Radic Biol Med* 16:845–850
- Heo MY, Yu KS, Kim H, Kim HP, Au WW (1992) Anticlastogenic effect of flavonoids against mutagen-induced micronuclei in mice. *Mutat Res* 284:243–249
- Hu CM, Cheng HW, Cheng YW, Kang JJ (2001) Mechanisms underlying the induction of vasorelaxation in rat thoracic aorta by sanguinarine. *Jpn J Pharmacol* 85:47–53
- Ignarro LJ, Sisodia M, Trinh K, Bedrood S, Wu G, Wei LH, Buga GM (2002) Nebivolol inhibits vascular smooth muscle cell proliferation by mechanisms involving nitric oxide but not cyclic GMP. *Nitric Oxide* 7:83–90
- Inazu M, Zhang H, Daniel EE (1995) Different mechanisms can activate Ca²⁺ entrance via cation currents in endothelial cells. *Life Sci* 56:11–17
- Iyengar R, Stuehr DJ, Marletta MA (1985) Macrophage synthesis of nitrite, nitrate and N-nitrosamines: precursors and role of respiratory burst. *Proc Natl Acad Sci USA* 84:6369–6373
- Jaffe EA (1985) Physiologic functions of normal endothelial cells. *Ann NY Acad Sci* 454:279–291
- Janel N, Dosne AM, Obert B, Meyer D, Kerbirou-Nabias D (1997) Functional characterization of bovine von Willebrand factor gene promoter in bovine endothelial cells demonstrates species-specific properties. *Gene* 198:127–134
- Jeon YJ, Youk ES, Lee SH, Suh J, Na YJ, Kim HM (2002) Polychlorinated biphenyl-induced apoptosis of murine spleen cells is aryl hydrocarbon receptor independent but caspases dependent. *Toxicol Appl Pharmacol* 181:69–78
- Jimenez R, Anderiambeloso E, Duarte J, Andriantsitohaina R, Jimenez J, Perez-Zicaino F, Zarzuelo A, Tamargo J (1999) Involvement of thromboxane A2 in the endothelium-dependent contraction induced by myricetin in rat isolated aorta. *Br J Pharmacol* 127:1539–1544
- Kanazawa K, Yamashita T, Ashida H, Danno G (1998) Antimutagenicity of flavones and flavonols to heterocyclic amines by specific and strong inhibition of the cytochrome P450 1A family. *Biosci Biotechnol Biochem* 62:970–977
- Kauffman RF, Schenck KW, Utterback BG, Crowe VG, Cohen MK (1987) In vitro vascular relaxation by new inotropic agents: relationship to phosphodiesterase inhibition and cyclic nucleotides. *J Pharmacol Exp Ther* 242:864–872
- Kerry NL, Abbey M (1997) Red wine and fractionated phenolic compounds prepared from red wine inhibit low density lipoprotein oxidation in vitro. *Atherosclerosis* 135:93–102
- Kerwin JF Jr, Heller M (1994) The arginine-nitric oxide pathway: a target for new drugs. *Med Res Rev* 14:23–74
- Kim SH, Kang KW, Kim KW, Kim ND (2000) Procyanidins in *Crataegus* extract evokes endothelium-dependent vasorelaxation in rat aorta. *Life Sci* 67:121–131
- Kinoshita N, Gelboin HV (1972) Aryl hydrocarbon hydroxylase and polycyclic hydrocarbon tumorigenesis: effect of the enzyme inhibitor 7,8-benzoflavone on tumorigenesis and macromolecular binding. *Proc Natl Acad Sci USA* 69:824–828
- Lanza F, Bereta A, Stierle A, Corre G, Cazenave JP (1987) Cyclic nucleotide phosphodiesterase inhibitors prevent aggregation of human platelets by raising cyclic AMP and reducing cytoplasmic free calcium mobilization. *Thromb Res* 45:477–484
- Li L, van Breemen C (1996) Agonist- and CPA-induced elevation of cytoplasmic free Ca²⁺ in intact valvular endothelium from rabbits. *Am J Physiol* 270:H837–H848
- Li Y, Wang E, Patten CJ, Chen L, Yang CS (1994) Effects of flavonoids on cytochrome P-450 dependent acetaminophen metabolism in rat and human liver microsomes. *Drug Metab Dispos* 22:566–571
- Lincoln TM, Cornwell TL (1991) Towards an understanding of the mechanism of action of cyclic AMP and cyclic GMP in smooth muscle relaxation. *Blood Vessels* 28:129–137
- Lincoln TM, Cornwell TL, Taylor AE (1990) cGMP-dependent protein kinase mediates the reduction of Ca²⁺ by cAMP in vascular SMC. *Am J Physiol* 258:C399–C407
- Low AM, Berdik M, Sormaz L, Gataianese S, Buchanan MR, Kwan CY, Daniel EE (1996) Plant alkaloids, tetrandrine and dernandezine, inhibit calcium-depletion stimulated calcium entry in human and bovine endothelial cells. *Life Science* 58:2327–2335
- Lowry OH, Rosebrough NJ, Farr AL, Randall RL (1951) Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265–275
- Luscher TF, Diederich D, Siebenmann R, Lehmann K, Stulz K, Stulz P, Von SL, Yang ZH, Turina M, Gradel E, Weber E (1988) Difference between endothelium-dependent relaxation in arterial and in venous coronary bypass grafts. *N Engl J Med* 319:462–467
- Mayer B, Schmidt K, Humbert P, Bohme E (1989) Biosynthesis of endothelium-derived relaxation factor: a cytosolic enzyme in porcine aortic endothelial cells Ca²⁺-dependently converts L-arginine into an activator of soluble guanylyl cyclase. *Biochem Biophys Res Commun* 164:678–685
- Middleton EJ (1998) Effect of plant flavonoids on immune and inflammatory cell function. *Adv Exp Med Biol* 439:175–182

- Millanvoye-Vanbrussel E, David-Duflho M, Pham TD, Iouzalén L, Aude Devynck M (1999). Regulation of arachidonic acid release by calcium influx in human endothelial cells. *J Vasc Res* 36:235–244
- Moncada S, Higgs A (1993) The L-arginine-nitric oxide pathway. *N Engl J Med* 329:2002–2012
- Moncada S, Palmer RM, Higgs EA (1991) Nitric oxide: physiology, pathophysiology, and pharmacology. *Pharmacol Rev* 43:109–142
- Moroney MA, Alcaraz MJ, Forder RA, Carey F, Hoult JRS (1988) Selectivity of neutrophil 5-lipoxygenase and cyclo-oxygenase inhibition by an anti-inflammatory flavonoid glycoside and related aglycone flavonoids. *J Pharm Pharmacol* 40:787–792
- Muraki K, Watanabe M, Imaizumi Y (2000) Nifedipine and nisoldipine modulate membrane potential of vascular endothelium via a myo-endothelial pathway. *Life Sci* 67:3163–3170
- Nijhoff WA, Groen GM, Peter HM (1993) Induction of rat hepatic and intestinal glutathione S-transferases and glutathione by dietary naturally occurring anticarcinogens. *Int J Oncol* 3:1131–1139
- Palmer RM, Ferrige AG, Moncada S (1987) Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 327:524–526
- Palmer RM, Ashton DS, Moncada S (1988) Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* 333:664–666
- Ramon Sanchez de Rojas V, Somoza B, Ortega T, Villar AM, Tejerina T (1999) Vasodilatory effect in rat aorta of eriodictyol obtained from *Satureja obovata*. *Planta Med* 65:234–238
- Rankin SM, Leake LS (1988) The modification of low-density lipoprotein by macrophage, by oxidation or proteolysis. *Agents Actions Suppl* 26:233–239
- Renaud S, DeLorgeril M (1992) Wine, alcohol, platelets and the French paradox for coronary heart disease. *Lancet* 339:1523–1526
- Reuelta MP, Hidalgo A, Cantabrana B (1999) Involvement of cAMP and beta-adrenoceptors in the relaxing effect elicited by flavonoids on rat uterine smooth muscle. *J Auton Pharmacol* 19:353–358
- Rodgers EH, Grant MH (1998) The effect of the flavonoids, quercetin, myricetin and epicatechin on the growth and enzyme activities of MCF-7 human breast cancer cells. *Chem Biol Interact* 116:213–228
- Rosenkranz-Weiss P, Sessa WC, Millstien S, Kaufman S, Watson CA, Pober JS (1994) Regulation of nitric oxide synthesis by proinflammatory cytokines in human umbilical vein endothelial cells. Elevations in tetrahydrobiopterin levels enhance endothelial nitric oxide synthase specific activity. *J Clin Invest* 93:2236–2243
- Sessa WC, Harrison JK, Barber CM (1992) Molecular cloning of a cDNA encoding endothelial cell nitric oxide synthase. *J Biol Chem* 267:15274–15276
- Shimada T, Yamazaki H, Foroozesh M, Hopkins NE, Alworth WL, Guengerich FP (1998) Selectivity of polycyclic inhibitors for human cytochrome P450s 1A1, and 1B1. *Chem Res Toxicol* 11:1048–1056
- Shou M, Grogan J, Mancewicz, JA, Krausz KW, Gonzalez FJ, Gelboin HV, Korzekwa KR (1994) Activation of CYP3A4: evidence for the simultaneous binding of two substrates in a cytochrome P450 active site. *Biochemistry* 33:6450–6455
- Siess MH, Leclerc J, Canivenc-Lavier MC, Rat P, Suschetet M (1995) Heterogenous effects of natural flavonoids on mono-oxygenase activities in human and rat liver microsomes. *Toxicol Appl Pharmacol* 130:73–78
- Silveira LA, Smith JL, Tan JL, Spudich JA (1998) MLCK-A, an unconventional myosin light chain kinase from *Dictyostelium*, is activated by a cGMP-dependent pathway. *Proc Natl Acad Sci USA* 95:13000–13005
- Sousa LR, Marletta MA (1985) Inhibition of cytochrome P450 activity in rat liver microsomes by the naturally occurring flavonoid quercetin. *Arch Biochem Biophys* 240:345–357
- Staples JE, Murante FG, Fiore NC, Gasiewicz TA, Silverstone AE (1998) Thymic alterations induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin are strictly dependent on aryl hydrocarbon receptor activation in hemopoietic cells. *J Immunol* 160:3844–3854
- Tassaneeyakul W, Birkett DJ, Veronese ME, Mcmanus ME, Tukey RH, Quattrochi LC, Gelboin HV, Miners JO (1993) Specificity of substrate and inhibitor probes for human cytochromes P450 1A1 and 1A2. *J Pharmacol Exp Ther* 265:401–407
- Trigueiro F, Cortes SF, Almeida RN, Lemos VS (2000) Endothelium-independent vasorelaxant effect of dioclein, a new flavonoid isolated from *Dioclea grandiflora*, in the rat aorta. *J Pharm Pharmacol* 52:1431–1434
- Tzeng SH, Ko WC, Ko FN, Teng CM (1991) Inhibition of platelet aggregation by some flavonoids. *Thromb Res* 64:91–100
- Uchida H, Tanaka Y, Ishii K, Nakayama K (1999) L-type Ca²⁺ channels are not involved in coronary endothelial Ca²⁺ influx mechanism responsible for endothelium-dependent relaxation. *Res Commun Mol Pathol Pharmacol* 104:127–144
- Vanhoutte PM, Rubanyi GM, Miller VM, Houston DS (1986) Modulation of vascular smooth muscle contraction by endothelium. *Annu Rev Physiol* 48:307–330
- Wang X, Chu W, van Breemen C (1996) Potentiation of acetylcholine-induced responses in freshly isolated rabbit aortic endothelial cells. *J Vasc Res* 33:414–424
- Wei H, Tye L, Bresnick E, Birt DF (1990) Effect of apigenin a plant flavonoid on epidermal ornithine decarboxylase and skin tumor promotion in mice. *Cancer Res* 50:499–502
- Xie Q, Zhang Y, Zhai, C, Bonanno JA (2002) Calcium influx factor from cytochrome P450 metabolism and secretion-like coupling mechanisms for capacitative calcium entry in corneal endothelial cells. *J Biol Chem* 277:16559–16566
- Xie QW, Cho HJ, Calaycay J (1992) Cloning and characterization of inducible nitric oxide synthase from mouse macrophages. *Science* 256:225–228
- Yakubu MA, Leffler CW (2002) L-type voltage-dependent Ca²⁺ channels in cerebral microvascular endothelial cells and ET-1 biosynthesis. *Am J Physiol* 283:C1687–C1695
- Yao X, Kwan HY, Chan FL, Chan NW, Huang Y. (2000) A protein kinase G-sensitive channel mediates flow-induced Ca²⁺ entry into vascular endothelial cells. *FASEB J* 14:932–938
- Youdim KM, McDonald J, Kalt W, Joseph JA (2002) Potential role of dietary flavonoids in reducing microvascular endothelium vulnerability to oxidative and inflammatory insults. *J Nutr Biochem* 13:282–288